

RESEARCH ONE-PAGER

NAD⁺

Cellular coenzyme · mitochondrial and DNA-repair research

500 mg and 1000 mg vials

OVERVIEW

NAD⁺ is not a peptide. It is a coenzyme every cell uses to make energy and to run DNA-repair tools. Levels are studied as they change with age, stress, and metabolic load.

IN PLAIN LANGUAGE

Think of NAD⁺ as cellular fuel-and-repair currency. People look at it for energy, recovery, and aging-biology research: mitochondria (the cell's engines) and PARP / sirtuin DNA-repair pathways.

WHAT RESEARCHERS LOOK AT

- Mitochondrial energy production
- DNA-repair and sirtuin-pathway research
- Cellular aging and stress-load models
- Mental and physical energy questions in the literature
- A cornerstone coenzyme, not a receptor-agonist peptide

LABORATORY RECONSTITUTION

Reconstitute aseptically with bacteriostatic water or the diluent specified in the laboratory SOP. Invert gently until dissolved; do not shake vigorously. Larger milligram vials need more diluent than a 10 mg peptide — confirm the volume for the vial on the bench before starting. Aliquot if the design requires repeated sampling. Offered as 500 mg and 1000 mg. Confirm diluent — NAD⁺ handling differs from a 10 mg peptide vial.

STORAGE

Lyophilized: refrigerate, protected from light and moisture. After reconstitution: refrigerate and follow laboratory SOP (commonly 14–30 days with bacteriostatic water). Avoid repeated freeze–thaw cycles.

RESEARCH USE ONLY

The language above is a plain-English reading of published research interests. It is not medical advice and not a claim that this product treats, prevents, or diagnoses any condition. This material is not a drug or cosmetic, and it is not intended for human or veterinary use. HarmonyX supplies it to qualified research purchasers for laboratory work. Certificates of analysis are available on request.

Questions on this vial? Paul and Thabby review every request.

research@harmony-x.com · 360-910-6274