

ORBIFOLD HILBERT SCHEMES AND A GENERALIZATION OF CORES AND QUOTIENTS

PAUL JOHNSON

1. INTRODUCTION

[1]

2. HILBERT SCHEMES OF POINTS ON SURFACES

2.1. Hilbert schemes of points in the plane.

Theorem 2.1 (Ellingsrud and Strømme, 1987).

$$\sum_{k,n \geq 0} b_k(\text{Hilb}_n(\mathbb{C}^2)) t^k q^n = \prod_{\ell=1}^{\infty} \frac{1}{1 - t^{2\ell-2} q^\ell}$$

2.1.1. *Białynicki-Birula decomposition.* Proved using the Białynicki-Birula decomposition. The Białynicki-Birula decomposition should be understood in analogy with Morse theory.

The role of the Morse flow will be played by the flow $x \mapsto \varepsilon x$ as $\varepsilon \in \mathbb{C}^*$ tends toward zero. We assume that the \mathbb{C}^* action on X is such that this limit point exists for all $x \in X$.

Let p be a fixed point of the \mathbb{C}^* action. Then linearizing the \mathbb{C}^* action on X gives a \mathbb{C}^* action on $T_p X$, and so $T_p X$ is not just a vector space but a \mathbb{C}^* representation, and hence decomposes into a direct sum of irreducible representations. Let V_a denote the irreducible representation of \mathbb{C}^* where $\varepsilon \in \mathbb{C}^*$ acts as ε^a .

Let $T_p^+ X$ (respectively $T_p^- X$) denote the subspace of $T_p X$ where the \mathbb{C}^* action acts with a positive (respectively negative) exponent. I.e., if

$$T_p X = \bigoplus_{n \in \mathbb{Z}} V_n^{e_n}$$

then

Let

$$\mathcal{S}_p = \{x \in X \mid \lim_{\varepsilon \rightarrow 0} \varepsilon x = p\}$$

Then clearly we have $X = \sqcup_p \mathcal{S}_p$; the point is that \mathcal{S}_p is a subvariety isomorphic to $\mathbb{C}^{l(p)}$.

We first observe that V_0 cannot occur in $T_p X$, as

We saw in Section [REFER BACK TO] that if X is a variety with a \mathbb{C}^* action with k isolated fixed points, then $\chi(X) = k$. If we know the weights of the \mathbb{C}^* action

on the tangent spaces of the fixed points, the Białynicki-Birula decomposition leverages this result to give the betti numbers of X , or even further the class of X in the Grothendieck ring of varieties.

2.1.2. *Tangent space to a monomial ideal.* To apply the Białynicki-Birula decomposition to $\mathrm{Hilb}_n(\mathbb{C}^2)$, we need to calculate the weights of the \mathbb{C}^* action on $T_\lambda \mathrm{Hilb}_n(\mathbb{C}^2)$.

[Ellingsrud and Strømme, Cheah]

$$T_\lambda \mathrm{Hilb}_n(\mathbb{C}^2) = \sum_{\square \in \lambda} \left(x^{-\ell(\square)} y^{a(\square)+1} + x^{\ell(\square)+1} y^{-a(\square)} \right)$$

Proof. First, we use $T_\lambda \mathrm{Hilb}_n(\mathbb{C}^2) =_R (\mathcal{I}_\lambda, R/\mathcal{I}_\lambda)$.

Analogy with the Grassmannian makes this intuitively plausible; indeed, if $V \subset W$ is a k dimensional subspace, then $T_V \mathrm{Gr}_k(W) = (V, V^\perp) = (V, W/V)$. Thus, we understand the deformations of \mathcal{I}_λ as a vector space; for the deformation to remain an ideal it is plausible that we should require the deformation to be a map of R modules and not just vector spaces.

More formally, it is a general fact that first order deformations of objects are given by ${}^1(\mathcal{F}, \mathcal{F})$, and obstructions to these deformations are given by ${}^2(\mathcal{F}, \mathcal{F})$; starting from this fact and considering the long exact sequences by taking (\mathcal{I}) to

$$0 \rightarrow \mathcal{I} \rightarrow R \rightarrow R/\mathcal{I} \rightarrow 0$$

gives the result. □

REFERENCES

- [1] S. M. Gusein-Zade, I. Luengo, and A. Melle-Hernández. A power structure over the Grothendieck ring of varieties. *Math. Res. Lett.*, 11(1):49–57, 2004.

UNIVERSITY OF SHEFFIELD

E-mail address: paul.johnson@sheffield.ac.uk