

# AN INTRODUCCION TO FJRW THEORY

TYLER JARVIS

.Our starting point is

**Theorem 1.** *Intersection theory of  $\psi$  classes on  $\overline{\mathcal{M}}_{g,n}$  is governed by the KdV hierarchy*

Immediately after this, Witten gave two generalizations of  $\overline{\mathcal{M}}_{g,n}$ . The first, are the moduli space of stable maps  $\overline{\mathcal{M}}_{g,n}(X, \beta)$ .

The second is  $\overline{\mathcal{M}}_{g,n}^{1/r}$ , the moduli space of  $r$ -spin curves, that is, curves  $\mathcal{C}$  together with a line bundle  $\mathcal{L}$  so that  $\mathcal{L}^{\otimes r} \cong \mathcal{K}_{\log}$ .

FJRW is a further generalization of  $r$ -spin curves, parallel to Gromov-Witten theory.

	GW-theory	FJRW
Input	$X$	$W \in \mathbb{C}[x_1, \dots, x_N]; G \text{ finite}$
State Space	$H^*(X)$	$\mathcal{A}^{W,G}$
Moduli space	$\overline{\mathcal{M}}_{g,n}(X, \beta) \rightarrow \overline{\mathcal{M}}_{g,n}$	$\overline{\mathcal{M}}_{g,n}^{W,G} \rightarrow \overline{\mathcal{M}}_{g,n}$
Virtual Class	$X$	$X$
CohFT	$\Lambda^{GW}$	$\Lambda^{FJRW}$
Larger picture		
	A	B
CY	GW	BK, BCOV, Costello-Li, etc.
LG	FJRW	Saito-Givental, Dubrovin-Zhang

FJRW theory requires the following inputs:

- $\mathbb{C}^*$  action on  $\mathbb{C}^N$  with weights  $c_1, \dots, c_N$ ; that is  $\lambda \cdot (x_1, \dots, x_N) = (\lambda^{c_1} x_1, \dots, \lambda^{c_N} x_N)$ .
- Polynomial  $W \in \mathbb{C}[x_1, \dots, x_N]$  that is quasihomogenous of weight  $d$  with respect to  $\mathbb{C}^*$  action – that is  $W(\lambda \cdot x) = \lambda^d W(x)$ .
- Quasihomogeneity forces  $W$  to have a singularity at the origin, we require this singularity to be isolated; that is,  $X_W = \{W = 0\} \subset WP(c_1, \dots, c_N)$  is smooth.
- $W$  should have no terms of form  $x_i x_j$ , which implies that the weights  $c_i/d \leq 1/2$  and weights are uniquely determined by  $W$ .

**Example 1.**  $W = x^r$  is the  $A_{r-1}$  singularity.

**Example 2.**  $W = x^5 + x^y2$  is the  $D_5$  singularity; we have  $d = 5, c_1 = 1, c_2 = 2$ .

The  $\mathbb{C}^*$  action and  $W$  together give the exponential grading element

$$J = (\exp(2\pi i c_1/d), \dots, \exp(2\pi i c_N/d))$$

with  $J$  fixing  $W$ .

$G$  is a finite abelian group acting diagonally on  $\mathbb{C}^N$  that fixes  $W$  and contains  $J$ .

Let  $G^{max} = \{(\alpha_1, \dots, \alpha_N) | W(\alpha_1 x_1, \dots, \alpha_N x_N) = W(x_1, \dots, x_N)\}$  denote the maximal diagonal symmetry group of  $W$ .

**Example 3.**

$$G_{D_6}^{max} = \{\zeta^{-2}, \zeta\} | \zeta^1 0 = 1\}$$

and  $J_{D_6} = (\exp(2\pi i/5), \exp(2\pi i 2/5))$ , so that  $\langle J \rangle G^{max}$ .

For all  $g \in G^{max}$ , we can write  $g = (\exp(2\pi i \theta_1), \dots, \exp(2\pi i \theta_N))$ , s.t.  $\theta_i \in [0, 1) \cap \mathbb{Q}$ . We call  $\theta_i$  the *phases* of  $g$ .

$$\text{Age}(g) = \sum \theta_i$$

$$N_g = \dim \text{Fix}(g)$$

Note:  $\text{Age}(g) + \text{Age}(g^{-1}) = N - N_g$ .

The idea is that, given the data described above, FJRW theory will produce a CohFT. What is a CohFT? It is a collection of tautological classes that behave like you would expect them to if you've been trained in Gromov-Witten theory.

More specifically, we have two gluing maps  $\rho_{\text{tree}}, \rho_{\text{loop}}$ , and the forgetting marked point map  $\tau$ .

A CohFT with flat identity consists of a state space  $\mathcal{A}$ , that has a pairing  $\langle, \rangle : \mathcal{A} \otimes \mathcal{A} \rightarrow \mathbb{C}$ , and a family  $\{\Lambda_{g,n}\}$ , with  $\Lambda_{g,n} \in H^*(\overline{\mathcal{M}}_{g,n}) \otimes (\mathcal{A}^*)^n$  and a vector  $1 \in \mathcal{A}$  that behave well with respect to the gluing maps, that is:

$$\rho_{\text{tree}}^* \Lambda_{g_1+g_2, k_1+k_2}(\alpha_{k_1}, \dots, \alpha_{k_1+k_2}) = \sum_{r,s} \Lambda_{g_1, k_1+1}(\alpha_1, \dots, \alpha_{k_1}, r) \eta^{rs} \Lambda_{g_2, k_2+1}(s, \alpha_{k_1+1}, \dots, \alpha_{k_1+k_2})$$

here  $r, s$  run over a basis of  $\mathcal{A}$ , and  $(\eta^{rs})$  is the inverse of the metric  $\langle, \rangle$  in terms of a basis of  $\mathcal{A}$ .

We have analogous requirements for  $\rho_{\text{loop}}^*$ :

$$\rho_{\text{loop}}^* \Lambda_{g+1, k}(\alpha_1, \dots, \alpha_k) = \sum_{r,s} \Lambda_{g, k+2}(\alpha_1, \dots, \alpha_k, r, s) \eta^{rs}$$

We want  $\Lambda_{g,k}$  to be symmetric; that is,  $S_k$  invariant, as an element of  $H^*(\overline{\mathcal{M}}_{g,n}) \otimes (\mathcal{A}^*)^k$ .

The flat identity condition states that y

$$\tau^*(\Lambda_{g,k}(\alpha_1, \dots, \alpha_k) = \Lambda_{g, k+1}(\alpha_1, \dots, \alpha_k, 1)$$

and  $\Lambda_{0,3}(\alpha_1, \alpha_2, 1) = \langle \alpha_1, \alpha_2 \rangle$ .

To construct the CohFT, we will use a moduli space (stack) consisting of curves with extra structure.

The moduli space will be  $\overline{\mathcal{M}}_{g,k}^{W,G}$  with maps  $e_i$  to  $\mathcal{IBG}$ , and a finite stabilization map st to  $\overline{\mathcal{M}}_{g,k}$ .

The moduli space  $\overline{\mathcal{M}}_{g,k}^{W,G}$  is smooth and proper.

This will be based off

$$\mathcal{W}_{g,k}^{W,G} = \{\mathcal{C}, p_1, \dots, p_k, \mathcal{L}_1, \dots, \mathcal{L}_N, \phi_i\}$$

where

- $(\mathcal{C}, p_1, \dots, p_k)$  is a stable pointed orbicurve.
- $\mathcal{L}_i$  are orbifold line bundles on  $\mathcal{C}$
- $W = \sum i = 1^s a_i W_i$ , with  $w_i$  monomials in the  $x_i$
- $\phi_i : W_i(\mathcal{L}_\infty, \dots, \mathcal{L}_N) \rightarrow \mathcal{K}_{\log}$
- The group action at each marked point and node factors through  $G$ .

Here  $\mathcal{K}_{\log=\mathcal{K}\otimes\mathcal{O}(\sum p_i)}$  has sections locally given by  $dz/z$  around the marked points.

**Example 4.** *r-spin curves* Let  $W = x^r, J = \exp(2\pi i/r), G^{max} = \langle J \rangle$  and

$$\overline{\mathcal{M}}_{g,k}^{x^r, G^{max}} = (\mathcal{C}, p_1, \dots, p_k$$

*Polischchuck-Vaintrob have an alternate definition.*

Let  $\Gamma = \mathbb{C}^*G \subset GL(N)$  MORE IN NOTES

## 1. STATE SPACE

First we will give the high-brow definition; then we will give a more down-to-earth version.

$W : \mathbb{C}^N \rightarrow \mathbb{C}$  is  $G$ -invariant, and so defines a map  $W : [\mathbb{C}^N/G] \rightarrow \mathbb{C}$ . -  $W^\infty = \text{Re}W^{-1}((M, \infty))$ ,  $M \gg 0$ ; that is, we want to consider those points in  $[\mathbb{C}^N/G]$  that map to something with very large real part under  $W$ . Then:

$$\mathcal{A} = H_{CR}^{*+2\sum c_i/d}([\mathbb{C}^N/G], W^\infty, \mathbb{C})$$

In another way, this is

$$\bigoplus_{\gamma \in G} H^*(\text{Fix}(\gamma); W_\gamma^\infty, \mathbb{C})^G$$

Note that  $\text{Fix}(\gamma) = \mathbb{C}^{N_\gamma}$ .

If  $\gamma = J = (\exp(2\pi i c_1/d), \dots, \exp(2\pi i c_N/d))$ , then  $\text{Fix}(J) = \{0\}$ , and  $\mathcal{A}_J = H^*(\{0\}, \mathbb{C}) = \mathbb{C}$ , and  $1 \in \mathcal{A}$  is defined to be  $1 \in \mathcal{A}_J$ .

Note that this is a bit surprising for those used to Chen-Ruan cohomology, as the identity lives in a twisted sector.

Another, more computational description:

**Theorem 2.** Wall, Orlik-Solomon, Seabastrani

$$H^*(\mathbb{C}^n, W^\infty; \mathbb{C}) \cong \Omega^N / (dW \wedge \Omega^{N-1})$$

as  $G$ -modules.

This is germs of  $N$ -forms on  $\mathbb{C}^N$  at 0, modulo  $dW = 0$ .

Another way to put this is  $\mathbb{C}[x_1, \dots, x_N]dx_1 \wedge \dots \wedge dx_N / (\frac{\partial W}{\partial x_1}dx_1 + \dots + \frac{\partial W}{\partial x_N}dx_N) \wedge dx_1 \wedge \dots \wedge dx_N$ .

$\cong \mathbb{C}[x_1, \dots, x_N] / (\frac{\partial W}{\partial x_1}, \dots, \frac{\partial W}{\partial x_N})dx_1 \wedge \dots \wedge dx_N$  This is the Milnor ring, or local algebra, of  $W$ .

Fact:  $\Omega^N / dW \wedge \Omega^{N-1}$  is finite dimensional if and only if  $W$  has an isolated singularity at 0. The dimension is called the milnor number, and is denoted  $\mu$ .

The Milnor ring is powerful: for quasihomogenous singularities, that is a natural class of equivalence of singularities, so that two singularities are equivalent if and only if their Milnor rings are isomorphic.

The pairing on  $H^*(\mathbb{C}^N, W^\infty, \mathbb{C})$  matches the residue pairing on  $\Omega^N / dW \wedge \Omega^{N-1}$ .

Residue pairing:

$$\langle f dx_1 \wedge \dots \wedge dx_N, g dx_1 \wedge \dots \wedge dx_N \rangle = \frac{1}{(2\pi i)^N} \int \frac{f g dx_1 \wedge \dots \wedge dx_N}{\frac{\partial W}{\partial x_1} \dots \frac{\partial W}{\partial x_N}}$$

Fact: in  $\mathbb{C}[x_1, \dots, x_N] / (\frac{\partial W}{\partial x_1}, \dots, \frac{\partial W}{\partial x_N})$  there is an obvious grading. The Hessian of  $W$  spans the part of highest degree. If  $\langle f, g \rangle = \alpha \mu$ , then  $fg = \alpha \text{Hess} + \text{lower order terms}$ .

*Fact:*

$$H^{N-p,p}(\mathbb{C}^n, W^\infty \cdot \mathbb{C})^{\langle J \rangle} \rightarrow \{\phi dx_1 \wedge \dots \wedge dx_N \mid \deg \phi = p\}$$

*Highest degree in Milnor ring is the degree of the Hessian, which is denoted  $\hat{c} = \sum (1 - 2(c_i/d))$ .*

*Then  $\langle, \rangle : \mathcal{A}^{p,q} \otimes \mathcal{A}^{\hat{c}-p, \hat{c}-q} \rightarrow \mathbb{C}$ .*

*So that  $\mathcal{A}$  behaves like the cohomology of a manifold of dimension  $\hat{c}$ , which may be fractional –  $\mathcal{A}$  may have fractional grading anyway. Bu if  $\hat{c} = N - 2$ , then  $X_W = \{W = 0\}$  is CY, and  $\mathcal{A}$  agrees with the cohomology of  $X_W$ .*

*MORE ON BIGRADING LATER*

*We have a moduli space, and a state space, the idea is that we will construct a virtual fundamental class  $[\overline{\mathcal{M}}_{g,k}^{W,G}] \in H_*(\overline{\mathcal{M}}_{g,l}^{W,G}) \otimes (\mathcal{A}^*)^k$  such that the classes we get by pushing down the virtual classes to  $\overline{\mathcal{M}}_{g,k}$  we get a CohFT.*

*More specifically, define:*

$$\Lambda_{g,k} = \frac{PDst_*[\overline{\mathcal{M}}_{g,k}^{W,G}]}{|G|^g \deg st}$$

*$\Lambda_{g,k}$  forms a CohFT*

*$\dim \Lambda_{g,k}(\alpha_1, \dots, \alpha_k) = (\hat{c} - 3)(1 - g) + k - \sum \deg \alpha_i$   $G^{max}$  – invariant.*

*Deformation invariance: If  $W_t$  is a 1-parameter family of polynomials, then  $\Lambda_{g,k}^{W_t, G}$  is independent of  $t$ .*

*Decomposition:  $\Lambda_{g,k}^{W_1+W_2, G_1 \times G_2} = \Lambda_{g,k}^{W_1, G_1} \otimes \Lambda_{g,k}^{W_2, G_2}$ .*

*Additional properties that facilitate computation.*