

TOBACCO RELATED DISORDER

Ifra Muskan

DIAGNOSTIC CRITERIA

The diagnostic criteria for Tobacco Use Disorder (TUD) are outlined in the Diagnostic and Statistical Manual of Mental Disorders, 5th Edition (DSM-5).

To diagnose TUD, a healthcare professional will assess the patient's behavior and experiences related to tobacco use, looking for at least two of the following criteria within a 12-month period:¹

- . _Tobacco is often used in larger amounts or over a longer period than intended_
- .2. _Desire or unsuccessful efforts to cut down or control tobacco use_.

CONTINUED...

3. _A great deal of time is spent in activities necessary to obtain, use, or recover from the effects of tobacco_
- .4. _Craving, or strong desire to use tobacco_
- .5. _Recurrent tobacco use resulting in a failure to fulfill major role obligations at work, school, or home_.
6. _Continued tobacco use despite having persistent or recurrent social or personal problems caused or worsened by tobacco use_
- .7. _Giving up important activities in order to use tobacco_.
8. _Recurrent tobacco use in situations in which it is physically hazardous_.
9. _Continued tobacco use despite knowledge of having a persistent or recurrent physical or psychological problem that is likely to have been caused or worsened by tobacco use_.

TOLERANCE AND WITHDRAWAL

10. _Tolerance (needing to use increasingly larger amounts to achieve the desired effect)_

.11. _Withdrawal, as manifested by the presence of two or more of the following symptoms when tobacco use has stopped or been reduced: irritability, frustration, or anger; anxiety; difficulty concentrating; increased appetite; insomnia; depression; or restlessness