

Institute of Applied Psychology
Govt. Graduate College for Women Baghbanpura, Lahore
Biological basis of behaviour- Assignment



Submitted by

Maida Naeem 429

Kashish kashif 414

Semester-6

Session: 2021-2025

Submitted to

Miss adeela

Course Code: APSY-365

Topic: structure and function of nervous system, ganglia
cell, neuron (types, structure, function) neural transmission, termination of
psp, neurological directional term

Nervous system

The nervous system is a complex network that coordinates the body's actions and sensory information by transmitting signals to and from different parts of the body. It is divided into two main parts: the central nervous system (CNS) and the peripheral nervous system (PNS).

Structure of the Nervous System

1. Central Nervous System (CNS):

- **Brain:** The control center of the body, responsible for processing sensory information, regulating bodily functions, and enabling thought and consciousness. It consists of different parts such as the cerebrum, cerebellum, and brainstem.
- **Spinal Cord:** A long, thin bundle of nerves that extends from the brain down through the spine, transmitting signals between the brain and the rest of the body.

2. Peripheral Nervous System (PNS):

- **Somatic Nervous System:** Controls voluntary movements by relaying information between the CNS and skeletal muscles.
- **Autonomic Nervous System (ANS):** Regulates involuntary functions such as heart rate, digestion, and respiratory rate. It is further divided into the sympathetic (fight or flight response) and parasympathetic (rest and digest) systems.
- **Nerves:** Bundles of axons that carry signals to and from the CNS and PNS.

Function of the Nervous System

1. **Sensory Input:** The nervous system receives information from sensory receptors that detect changes in the environment (internal and external). This information is sent to the CNS for processing.

2. **Integration:** The CNS processes and interprets the sensory input, deciding what action, if any, should be taken.

3. **Motor Output:** The CNS sends signals through the PNS to effector organs (muscles and glands) to initiate a response or action.

4. **Homeostasis:** The nervous system helps maintain homeostasis by regulating body functions such as temperature, blood pressure, and fluid balance through the ANS.

5. **Mental Activity:** The brain is involved in higher-order functions such as thinking, memory, learning, and emotions.

6. **Coordination:** The nervous system coordinates voluntary and involuntary actions, ensuring that the body functions as a coherent unit.

Key Components

1. **Neurons:** The basic functional units of the nervous system, responsible for transmitting nerve impulses. They consist of a cell body, dendrites (receive signals), and an axon (sends signals).
2. **Synapses:** Junctions between neurons that allow them to communicate with each other through neurotransmitters.
3. **Glial Cells:** Support cells in the nervous system that provide structural and metabolic support to neurons, maintain homeostasis, form myelin, and participate in signal transmission.

The nervous system's intricate design enables it to perform its critical role in regulating and coordinating bodily functions, responding to stimuli, and enabling complex behaviors.

Electrical and Chemical Signaling

1. Action Potential:

- A rapid change in the electrical charge across the neuron's membrane.
- Generated when the neuron is sufficiently stimulated, leading to the propagation of an electrical signal along the axon.

2. Neurotransmitters:

- Chemical messengers released by neurons at synapses.
- Bind to receptors on the adjacent neuron or effector cell, influencing its activity.

Neurons are integral to the functioning of the nervous system, enabling communication between different parts of the body and facilitating complex processes such as thought, sensation, and movement.

Ganglia cell

Ganglia (singular: ganglion) are clusters of neuron cell bodies located outside the central nervous system (CNS). Ganglia serve as relay points and processing centers for the transmission of neural signals.

Types of Ganglia

1. Sensory Ganglia:

- Contain the cell bodies of sensory neurons.
- Located on the dorsal roots of spinal nerves and on some cranial nerves.

- Examples: Dorsal root ganglia, trigeminal ganglia.

2. Autonomic Ganglia:

- Part of the autonomic nervous system (ANS).
- Contain the cell bodies of autonomic (motor) neurons.
- Divided into sympathetic and parasympathetic ganglia.
 - Sympathetic Ganglia: Located along the sympathetic trunk or prevertebral ganglia. They relay signals from the CNS to various organs to prepare the body for "fight or flight" responses.
 - Parasympathetic Ganglia: Located near or within the walls of target organs. They relay signals to promote "rest and digest" activities.

Structure of Ganglia

- Neuron Cell Bodies: The core of the ganglion, containing the cell bodies of neurons.
- Satellite Cells: Glial cells that surround the neuron cell bodies within the ganglion, providing support and nutrients, and helping regulate the extracellular environment.
- Axons: Nerve fibers that enter and leave the ganglion, carrying signals to and from the neuron cell bodies.

Function of Ganglia

1. Relay Points:

- Ganglia act as relay stations where signals are transmitted from one neuron to another.
- Sensory ganglia relay sensory information from the peripheral nervous system (PNS) to the CNS.
- Autonomic ganglia relay motor information from the CNS to various organs and tissues.

2. Signal Processing:

- Some ganglia can modulate and integrate incoming signals before passing them on to the next neuron or target organ.
- This processing helps fine-tune bodily responses to various stimuli.

Role in the Nervous System

- Sensory Pathways: Sensory ganglia are essential for transmitting sensory information such as touch, pain, and temperature from the body to the brain.

- Autonomic Control: Autonomic ganglia play a crucial role in regulating involuntary bodily functions, such as heart rate, digestion, and respiratory rate.

Ganglia are key components of the peripheral nervous system, ensuring efficient transmission and processing of neural signals between the CNS and the rest of the body.

Neuron

A neuron, or nerve cell, is the fundamental unit of the nervous system responsible for transmitting information throughout the body. Neurons communicate via electrical and chemical signals and play a crucial role in sensing stimuli, processing information, and controlling bodily functions.

Structure and characteristics of neuron

Structure of a Neuron

1. Cell Body (Soma):

- Nucleus: Contains the cell's genetic material (DNA) and regulates the cell's activities.
- Cytoplasm: Surrounds the nucleus and contains various organelles, including mitochondria (energy production), ribosomes (protein synthesis), and endoplasmic reticulum (processing and transport of proteins).
- Nissl Bodies: Rough endoplasmic reticulum in neurons, involved in protein synthesis.

2. Dendrites:

- Branch-like extensions from the cell body.
- Receive signals from other neurons or sensory receptors and transmit these signals to the cell body.
- Have numerous branches, increasing their surface area for synaptic connections.

3. Axon:

- A single, long projection that conducts electrical impulses away from the cell body.
- Axon Hillock: The region where the axon originates from the cell body, often the site of action potential initiation.
- Myelin Sheath: A fatty layer that insulates the axon, formed by glial cells (Schwann cells in the PNS and oligodendrocytes in the CNS). Myelination increases the speed of electrical signal transmission.
- Nodes of Ranvier: Gaps in the myelin sheath where the axon membrane is exposed, facilitating rapid signal conduction via saltatory conduction.

- Axon Terminals: The ends of the axon that form synapses with other neurons or effector cells, where neurotransmitters are released to transmit signals.

Characteristics of Neurons

1. Excitability:

- Neurons can respond to stimuli (e.g., chemical, electrical, or mechanical) and convert these into neural signals (action potentials).

2. Conductivity:

- Neurons can transmit electrical impulses along their membrane.

3. Secretion:

- Neurons release neurotransmitters at synapses to communicate with other neurons or effector cells.

4. Longevity:

- Neurons have an extended lifespan and, in many cases, last the entire life of an organism. Unlike many other cell types, neurons do not undergo frequent cell division.

5. Amitotic:

- Most neurons do not divide after they have fully matured. Once they are damaged or die, they are not typically replaced.

6. Metabolic Rate:

- Neurons have a high metabolic rate and require a constant supply of oxygen and glucose to function properly.

Types of Neurons (Based on Function)

1. Sensory Neurons (Afferent Neurons):

- Transmit sensory information from receptors to the CNS.

2. Motor Neurons (Efferent Neurons):

- Transmit signals from the CNS to muscles and glands to elicit responses.

3. Interneurons (Association Neurons):

- Found primarily in the CNS, they connect neurons within the CNS and play a role in processing information and determining responses.

Types of Neurons (Based on Structure)

1. Unipolar Neurons:

- Have a single process extending from the cell body, which divides into two branches: one acting as a dendrite and the other as an axon. Common in sensory neurons.

2. Bipolar Neurons:

- Have one axon and one dendrite extending from the cell body. Found in sensory organs such as the retina of the eye.

3. Multipolar Neurons:

- Have one axon and multiple dendrites. The most common type of neuron in the CNS and are primarily involved in motor and interneuron functions.

Neurons are highly specialized cells with unique structures and characteristics that enable them to perform their vital roles in transmitting, processing, and responding to information throughout the nervous system.

Types of neuron

Neurons can be classified based on their function, structure, and location. Here are the primary classifications:

Types of Neurons by Function

1. Sensory Neurons (Afferent Neurons):

- Function: Transmit sensory information from sensory receptors (e.g., skin, eyes, ears) to the central nervous system (CNS).
- Location: Found in sensory organs and throughout the body, connecting to the spinal cord and brain.
- Example: Neurons in the retina of the eye that detect light.

2. Motor Neurons (Efferent Neurons):

- Function: Transmit signals from the CNS to effector organs (muscles and glands) to initiate a response or action.
- Location: Found in the CNS and extend to muscles and glands.
- Example: Neurons that control muscle contractions.

3. Interneurons (Association Neurons):

- Function: Connect neurons within the CNS and process information. They play a crucial role in reflexes and higher functions like learning and memory.

- Location: Found entirely within the CNS (brain and spinal cord).

- Example: Neurons in the spinal cord that mediate reflexes.

Types of Neurons by Structure

1. Unipolar Neurons:

- Structure: Have a single process extending from the cell body, which divides into two branches: one functioning as a dendrite and the other as an axon.

- Function: Mainly sensory neurons that transmit sensory information to the CNS.

- Location: Common in sensory neurons of the peripheral nervous system (PNS).

2. Bipolar Neurons:

- Structure: Have one axon and one dendrite extending from opposite ends of the cell body.

- Function: Primarily involved in sensory functions.

- Location: Found in sensory organs such as the retina of the eye, the olfactory epithelium in the nose, and the cochlea in the ear.

3. Multipolar Neurons:

- Structure: Have one axon and multiple dendrites extending from the cell body.

- Function: Most common type of neuron in the CNS, involved in integrating information from other neurons and transmitting it to muscles and glands.

- Location: Found throughout the CNS.

- Example: Motor neurons and interneurons in the brain and spinal cord.

4. Anaxonic Neurons:

- Structure: Have multiple dendrites but no apparent axon.

- Function: Do not generate action potentials but are involved in local signal integration.

- Location: Found in some parts of the CNS and in sensory organs.

Types of Neurons by Location

1. CNS Neurons:

- Function: Involved in processing information, memory, thought, and controlling voluntary actions.
- Location: Brain and spinal cord.
- Example: Pyramidal cells in the cerebral cortex.

2. PNS Neurons:

- Function: Transmit signals between the CNS and the rest of the body.
- Location: Throughout the body, connecting to the spinal cord and brain.
- Example: Dorsal root ganglion neurons.

Specialized Neurons

1. Pyramidal Neurons:

- Function: Involved in motor control, cognitive function, and neural communication.
- Location: Cerebral cortex and hippocampus.
- Structure: Triangular-shaped cell body with a long apical dendrite and several basal dendrites.

2. Purkinje Cells:

- Function: Regulate motor movements and coordination.
- Location: Cerebellum.
- Structure: Large cell body with a highly branched dendritic arbor.

3. Mirror Neurons:

- Function: Fire both when performing an action and when observing the same action performed by another.
- Location: Various brain regions, including the premotor cortex.
- Role: Believed to be involved in learning through imitation and understanding others' actions.

These classifications illustrate the diversity of neurons in terms of structure, function, and location, highlighting their specialized roles in the nervous system.

Kashish 414

Neural Transmission

Neural transmission, also known as synaptic transmission, is the process by which neurons communicate with each other or with effector cells (muscles or glands). This involves both electrical and chemical signals. Here's an overview of the steps involved:

Steps in Neural Transmission

1. Resting Potential:

- Neurons maintain a resting membrane potential, typically around -70 millivolts (mV), due to the difference in ion concentrations inside and outside the cell. This is achieved by the sodium-potassium pump, which actively transports Na^+ ions out of the cell and K^+ ions into the cell.

2. Depolarization:

- When a neuron receives a strong enough stimulus, sodium channels open, allowing Na^+ ions to flow into the cell. This influx of positive ions causes the membrane potential to become less negative, leading to depolarization. If the depolarization reaches a certain threshold (usually around -55 mV), an action potential is generated.

3. Action Potential:

- An action potential is a rapid, temporary change in the membrane potential that travels along the axon. It follows an all-or-nothing principle: once the threshold is reached, the action potential fires completely.

- During the action potential, Na^+ channels open rapidly, causing a spike in membrane potential (up to +30 mV). This is followed by the opening of potassium channels, allowing K^+ ions to flow out of the cell, repolarizing the membrane back to its resting potential.

4. Propagation of the Action Potential:

- The action potential travels down the axon as a wave of depolarization. In myelinated axons, this occurs via saltatory conduction, where the action potential jumps from one node of Ranvier (gaps in the myelin sheath) to the next, increasing the speed of transmission.

5. Synaptic Transmission:

- When the action potential reaches the axon terminals, it triggers the opening of voltage-gated calcium channels, allowing Ca^{2+} ions to enter the terminal.

- The influx of Ca^{2+} ions causes synaptic vesicles filled with neurotransmitters to fuse with the presynaptic membrane and release their contents into the synaptic cleft (the gap between the neurons).

6. Neurotransmitter Release:

- Neurotransmitters diffuse across the synaptic cleft and bind to specific receptors on the postsynaptic membrane (the neuron receiving the signal).

- This binding can either depolarize (excite) or hyperpolarize (inhibit) the postsynaptic neuron, depending on the type of neurotransmitter and receptor involved.

7. Post-Synaptic Potential:

- If the postsynaptic potential is excitatory (excitatory postsynaptic potential or EPSP), it brings the membrane potential closer to the threshold for firing an action potential.

- If the postsynaptic potential is inhibitory (inhibitory postsynaptic potential or IPSP), it makes the membrane potential more negative, moving it further from the threshold and reducing the likelihood of firing an action potential.

8. Termination of Signal:

- The neurotransmitter's action is terminated by various mechanisms, including reuptake into the presynaptic neuron, enzymatic degradation, or diffusion away from the synapse.

- This ensures that the neurotransmitter does not continuously stimulate the postsynaptic neuron.

Types of Neurotransmitters

1. Excitatory Neurotransmitters:

- Promote the generation of action potentials in the postsynaptic neuron.

- Examples: Glutamate, acetylcholine.

2. Inhibitory Neurotransmitters:

- Inhibit the generation of action potentials in the postsynaptic neuron.

- Examples: GABA (gamma-aminobutyric acid), glycine.

3. Modulatory Neurotransmitters:

- Modify the overall activity of neurons and can have both excitatory and inhibitory effects depending on the context.

- Examples: Dopamine, serotonin, norepinephrine.

Importance of Neural Transmission

- Communication: Enables rapid and precise communication between different parts of the nervous system and the body.

- Coordination: Coordinates complex processes such as movement, sensation, thought, and emotion.

- Adaptation: Allows the nervous system to adapt to changes in the environment and maintain homeostasis.

Neural transmission is a fundamental process that underlies all neural activities, from basic reflexes to complex cognitive functions.

Neural communication

The communication system within the nervous system is highly specialized and involves intricate processes to transmit information rapidly and accurately between neurons and to other cells. Here's an overview of the communication system within the nervous system:

Components of the Neural Communication System

1. Neurons:

- The primary cells responsible for transmitting electrical and chemical signals.

2. Synapses:

- Junctions where neurons communicate with other neurons, muscles, or glands.

3. Neurotransmitters:

- Chemicals that transmit signals across a synapse from one neuron to another.

4. Receptors:

- Proteins on the postsynaptic cell membrane that bind to neurotransmitters and initiate a response.

5. Glial Cells:

- Support cells that provide structural and metabolic support to neurons, maintain homeostasis, and facilitate signal transmission.

Steps in Neural Communication

1. Signal Reception:

- Dendrites and Cell Body: Neurons receive incoming signals from other neurons at synapses located on their dendrites and cell body. These signals can be excitatory or inhibitory.

2. Signal Integration:

- Summation: The neuron integrates all incoming excitatory and inhibitory signals. If the cumulative signal exceeds a certain threshold, an action potential is generated at the axon hillock.

3. Action Potential Generation:

- Depolarization: If the threshold is reached, voltage-gated sodium channels open, causing a rapid influx of Na^+ ions and a reversal of the membrane potential.

- Repolarization: Voltage-gated potassium channels open, allowing K^+ ions to exit the cell, restoring the negative membrane potential.

4. Action Potential Propagation:

- Propagation: The action potential travels down the axon as a wave of depolarization.

- Saltatory Conduction: In myelinated axons, the action potential jumps between nodes of Ranvier, speeding up transmission.

5. Synaptic Transmission:

- Arrival at Axon Terminal: When the action potential reaches the axon terminal, it causes voltage-gated calcium channels to open.

- Neurotransmitter Release: The influx of Ca^{2+} ions triggers the fusion of synaptic vesicles with the presynaptic membrane, releasing neurotransmitters into the synaptic cleft.

6. Signal Transmission Across the Synapse:

- Binding to Receptors: Neurotransmitters diffuse across the synaptic cleft and bind to specific receptors on the postsynaptic membrane.

- Postsynaptic Response: Binding of neurotransmitters to receptors causes ion channels to open or close, leading to a change in the membrane potential of the postsynaptic cell.

7. Termination of Signal:

- Reuptake: Neurotransmitters are taken back into the presynaptic neuron for reuse.

- Enzymatic Degradation: Enzymes break down neurotransmitters in the synaptic cleft.

- Diffusion: Neurotransmitters diffuse away from the synapse.

Types of Synapses

1. Electrical Synapses:

- Gap Junctions: Direct physical connections between neurons that allow ions to flow directly from one cell to another.

- Speed: Very fast transmission.

Example: Found in certain brain regions and in the heart.

2. Chemical Synapses:

- Synaptic Cleft: A small gap between the presynaptic and postsynaptic neurons.
- Neurotransmitter Release: Involves the release of neurotransmitters that bind to receptors on the postsynaptic cell.
- Versatility: Allows for complex modulation of signals.

Types of Receptors

1. Ionotropic Receptors:

- Ligand-Gated Ion Channels: Open or close in response to neurotransmitter binding, allowing ions to flow across the membrane.
- Speed: Fast response.
- Example: Nicotinic acetylcholine receptors.

2. Metabotropic Receptors:

- G-Protein Coupled Receptors: Activate second messenger systems in the cell when a neurotransmitter binds.
- Speed: Slower but allows for amplification and prolonged effects.
- Example: Muscarinic acetylcholine receptors.

Role of Glial Cells

1. Astrocytes:

- Maintain the extracellular environment, provide metabolic support, and help form the blood-brain barrier.

2. Oligodendrocytes and Schwann Cells:

- Form the myelin sheath around axons in the CNS and PNS, respectively.

3. Microglia:

- Act as immune cells in the CNS, clearing debris and protecting against pathogens.

4. Ependymal Cells:

- Line the ventricles of the brain and the central canal of the spinal cord, involved in the production and circulation of cerebrospinal fluid (CSF).

Importance of Neural Communication

- Coordination: Ensures coordinated functioning of various body systems.
- Response to Stimuli: Enables rapid response to environmental changes and stimuli.
- Homeostasis: Maintains internal stability by regulating physiological processes.
- Cognition and Emotion: Underlies complex processes such as thought, memory, learning, and emotions.

The nervous system's communication system is essential for maintaining the body's overall function and adapting to internal and external changes.

Process of NT

The process of neurotransmission (NT) is the mechanism by which neurons communicate with each other and with other types of cells, such as muscle cells or gland cells. This process involves the release of neurotransmitters, which are chemical messengers that transmit signals across synapses. Here's a detailed look at the steps involved in neurotransmission:

Steps in the Process of Neurotransmission

1. Synthesis and Storage of Neurotransmitters:

- Synthesis: Neurotransmitters are synthesized from precursor molecules in the neuron's cell body or axon terminals.
- Storage: Once synthesized, neurotransmitters are stored in synaptic vesicles within the axon terminals.

2. Arrival of Action Potential:

- Depolarization: When an action potential reaches the axon terminal, it causes depolarization of the presynaptic membrane.
- Calcium Influx: The depolarization opens voltage-gated calcium channels, allowing Ca^{2+} ions to enter the axon terminal.

3. Release of Neurotransmitters:

- Vesicle Fusion: The influx of Ca^{2+} ions triggers synaptic vesicles to move towards and fuse with the presynaptic membrane.
- Exocytosis: The vesicles release their neurotransmitter contents into the synaptic cleft through exocytosis.

4. Binding to Receptors:

- Diffusion: Neurotransmitters diffuse across the synaptic cleft.
- Receptor Binding: Neurotransmitters bind to specific receptors on the postsynaptic membrane. These receptors can be ionotropic (ligand-gated ion channels) or metabotropic (G-protein coupled receptors).

5. Postsynaptic Response:

- Ionotropic Receptors: Binding of neurotransmitters to ionotropic receptors causes ion channels to open, leading to a rapid change in the postsynaptic membrane potential. This can result in either an excitatory postsynaptic potential (EPSP) or an inhibitory postsynaptic potential (IPSP).
- Metabotropic Receptors: Binding of neurotransmitters to metabotropic receptors activates intracellular signaling pathways (second messenger systems), leading to slower but longer-lasting effects.

6. Termination of Signal:

- Reuptake: Neurotransmitters are taken back into the presynaptic neuron via transporter proteins. They can be repackaged into vesicles for reuse.
- Enzymatic Degradation: Enzymes in the synaptic cleft break down neurotransmitters into inactive components.
- Diffusion: Neurotransmitters diffuse away from the synapse and are eventually absorbed by surrounding cells or degraded.

7. Postsynaptic Integration:

- The postsynaptic neuron integrates the various excitatory and inhibitory signals it receives. If the combined effect of these signals depolarizes the membrane to the threshold level, an action potential is generated in the postsynaptic neuron.

Key Neurotransmitters and Their Functions

1. Acetylcholine (ACh):

- Function: Involved in muscle activation, attention, learning, and memory.
- Receptors: Nicotinic (ionotropic) and muscarinic (metabotropic).

2. Glutamate:

- Function: The main excitatory neurotransmitter in the CNS, involved in learning and memory.
- Receptors: NMDA, AMPA, and kainate receptors (all ionotropic).

3. Gamma-Aminobutyric Acid (GABA):

- Function: The main inhibitory neurotransmitter in the CNS, involved in reducing neuronal excitability.
- Receptors: GABA_A (ionotropic) and GABA_B (metabotropic).

4. Dopamine:

- Function: Involved in reward, motivation, and motor control.
- Receptors: D1 and D2 families (metabotropic).

5. Serotonin (5-HT):

- Function: Involved in mood regulation, appetite, and sleep.
- Receptors: Multiple types, mainly metabotropic.

6. Norepinephrine (Noradrenaline):

- Function: Involved in attention, arousal, and the fight-or-flight response.
- Receptors: Alpha and beta adrenergic receptors (metabotropic).

7. Endorphins:

- Function: Involved in pain relief and feelings of euphoria.
- Receptors: Opioid receptors (metabotropic).

Summary

Neurotransmission is a highly orchestrated process involving the synthesis, release, binding, and termination of neurotransmitter signals. This complex communication system enables neurons to transmit information rapidly and precisely, facilitating everything from basic reflexes to complex cognitive functions.

Termination of PSP

The termination of postsynaptic potentials (PSPs) is essential to ensure that neural signals are transient and do not persist indefinitely. This allows neurons to reset and be ready to respond to new signals. The termination of PSPs involves several mechanisms:

Mechanisms of Terminating Postsynaptic Potentials

1. Reuptake of Neurotransmitters:

- Process: Specialized transporter proteins in the presynaptic membrane or nearby glial cells actively transport neurotransmitters back into the presynaptic neuron or glial cells.

- Example: The serotonin transporter (SERT) reabsorbs serotonin, and the dopamine transporter (DAT) reabsorbs dopamine.

2. Enzymatic Degradation:

- Process: Enzymes in the synaptic cleft or associated with the postsynaptic membrane break down neurotransmitters into inactive metabolites.

- Example: Acetylcholinesterase breaks down acetylcholine into acetate and choline.

3. Diffusion Away from the Synapse:

- Process: Neurotransmitters diffuse out of the synaptic cleft and away from the receptors, reducing their concentration at the synapse.

- Example: This can occur with any neurotransmitter but is especially significant for those not rapidly reabsorbed or degraded.

4. Desensitization of Receptors:

- Process: Receptors on the postsynaptic membrane become less responsive to neurotransmitters after prolonged exposure, temporarily reducing their activity.

- Example: Some receptors, such as certain glutamate receptors, can become temporarily inactivated or desensitized after continuous stimulation.

Detailed Steps in Each Mechanism

1. Reuptake:

- Transporter Binding: Neurotransmitters bind to specific transporter proteins on the presynaptic neuron or glial cells.

- Transport: The transporter proteins use energy (typically from ATP) to move neurotransmitters across the cell membrane back into the cell.

- Recycling: Once inside the cell, neurotransmitters can be repackaged into synaptic vesicles for reuse or broken down by intracellular enzymes.

2. Enzymatic Degradation:

- Enzyme Action: Specific enzymes break down neurotransmitters into non-active metabolites.

- Example of Enzymes:

- Acetylcholinesterase: Breaks down acetylcholine into choline and acetate.

- Monoamine Oxidase (MAO) and Catechol-O-Methyltransferase (COMT): Break down monoamines like dopamine, norepinephrine, and serotonin.

3. Diffusion:

- Concentration Gradient: Neurotransmitters diffuse away from the synaptic cleft down their concentration gradient.

- Extracellular Space: Neurotransmitters may be taken up by nearby cells or degraded by extracellular enzymes in the surrounding tissue.

4. Receptor Desensitization:

- Temporary Inactivation: Prolonged exposure to a neurotransmitter can cause receptors to become temporarily inactive.

- Receptor Conformation: Changes in receptor conformation can reduce their binding affinity for the neurotransmitter or block their ion channels.

- Phosphorylation: Certain receptors undergo phosphorylation, which can reduce their activity.

Importance of Termination Mechanisms

- Signal Precision: Ensures that neural signals are discrete and precisely timed, allowing for accurate communication between neurons.

- Prevention of Overstimulation: Prevents continuous activation of the postsynaptic neuron, which could lead to excitotoxicity or desensitization.

- Resetting the Synapse: Allows the synapse to be ready for the next neurotransmitter release, maintaining efficient neural communication.

- Homeostasis: Maintains the balance of neurotransmitter levels in the synaptic cleft and surrounding areas, ensuring proper neural function.

These termination mechanisms are crucial for maintaining the proper functioning of the nervous system, allowing neurons to communicate effectively and respond dynamically to changing signals.

Neuroanatomical directional terms and planes of reference are essential for accurately describing the locations and orientations of structures within the nervous system. Here's an overview:

Directional Terms

1. Anterior (Rostral):

- Towards the front or head end.

- Example: The frontal lobe is anterior to the parietal lobe.

2. Posterior (Caudal):

- Towards the back or tail end.

- Example: The occipital lobe is posterior to the parietal lobe.

3. Superior (Dorsal):

- Towards the top or back side.

- Example: The parietal lobe is superior to the temporal lobe.

4. Inferior (Ventral):

- Towards the bottom or belly side.

- Example: The temporal lobe is inferior to the parietal lobe.

5. Medial:

- Towards the midline of the body or structure.

- Example: The corpus callosum is medial within the brain.

6. Lateral:

- Away from the midline of the body or structure.

- Example: The temporal lobes are lateral to the thalamus.

7. Proximal:

- Closer to the point of attachment or origin.

- Example: The spinal cord is proximal to the peripheral nerves.

8. Distal:

- Farther from the point of attachment or origin.

- Example: The fingers are distal to the wrist.

Planes of Reference

1. Sagittal Plane:

- Divides the body or brain into right and left parts.
- Midsagittal Plane (Median Plane): Divides the body or brain into equal right and left halves.
- Parasagittal Plane: Divides the body or brain into unequal right and left parts.

2. Coronal Plane (Frontal Plane):

- Divides the body or brain into anterior (front) and posterior (back) parts.
- Example: A coronal section of the brain might show the frontal lobe in front and the occipital lobe at the back.

3. Transverse Plane (Horizontal Plane):

- Divides the body or brain into superior (upper) and inferior (lower) parts.
- Example: A transverse section of the brain might show the separation between the upper part of the cerebrum and the lower part of the brainstem.

Additional Neuroanatomical Terms

1. Ipsilateral:

- On the same side of the body.
- Example: The right hand and right foot are ipsilateral.

2. Contralateral:

- On the opposite side of the body.
- Example: The right hand is contralateral to the left hand.

3. Bilateral:

- Present on both sides of the body.
- Example: Bilateral symmetry in brain hemispheres.

4. Unilateral:

- Present on one side of the body.
- Example: A unilateral stroke affects one hemisphere of the brain.

5. Superficial:


- Closer to the surface of the body.
- Example: The cortex is superficial to the subcortical structures.

6. Deep:

- Further away from the surface of the body.
- Example: The thalamus is deep within the brain.

Summary Diagram

Here is a simplified diagram of these planes and terms as they relate to the brain:



Practical Application

- Medical Imaging: Directional terms and planes are crucial in interpreting MRI, CT scans, and other imaging modalities.
- Neurosurgery: These terms help neurosurgeons navigate the brain accurately during procedures.
- Neuroanatomy Education: Understanding these terms is fundamental for students and professionals studying brain structure and function.

These terms and planes provide a standardized way of describing the location and orientation of brain structures, ensuring clear communication in both clinical and academic settings.