

Assignment no 10


“Communication”

Name: Fatima and Laraib Fatima

Roll no: 392 and 401

Bs Psychology 6 semester

Govt. Graduate College for Women Bagbanpura Lahore



COMMUNICATION

Roll no 392:

Topics: (Factors in Development, Speech abnormalities, Aphasia, Apraxia, Wernicke's Aphasia)

Communication

In behavioral neuroscience, **communication refers to the process by which information is transmitted and received between individuals through various forms of signaling.** This can include verbal language, non-verbal cues (such as body language, facial expressions, and gestures), and other forms of signaling such as pheromones or electrical signals in animals. Communication in this context often involves studying how the brain and nervous system process, interpret, and respond to these signals, as well as understanding the underlying neural mechanisms that facilitate effective communication.

Factors in the Development of Speech:

The development of speech is influenced by a combination of genetic, biological, environmental, and social factors.

Genetic Factors:

- Genetic factors include inherited traits that can influence language development. Specific genes, such as FOXP2, have been linked to speech and language abilities. These genetic components can predispose individuals to varying abilities in language acquisition.

Neurological Factors:

- Neurological factors are crucial since brain development and functioning play a key role. Areas such as Broca's area, which is involved in speech production, and Wernicke's area, which is involved in language comprehension, are essential. Proper neural connectivity and functioning of these areas are vital for effective speech development.

Hearing Ability:

- Hearing ability is fundamental because hearing is essential for learning and producing speech. Children with hearing impairments may face significant challenges in speech development. Early detection and intervention can help mitigate these challenges and support speech development.

Cognitive Development:

- Cognitive development encompasses general cognitive abilities, including memory, attention, and problem-solving skills, which support the learning of language and speech. Cognitive development enables children to understand and produce language, form sentences, and follow conversations.

Environmental Factors:

- Environmental exposure to language-rich environments is critical. This includes hearing spoken language, being read to, and engaging in conversations. Environments that provide diverse and frequent language input can significantly enhance speech development.

Social Interaction:

- Social interaction with family, peers, and others encourages language practice and development. Regular social interactions provide opportunities for children to practice speech, receive feedback, and learn the rules and nuances of language use.

Parental Influence:

- Parental influence plays a key role in speech development. Parents and caregivers model speech, respond to children's attempts at communication, and provide encouragement and correction. The quality and quantity of parental interaction can greatly impact a child's speech development.

Health and Nutrition:

- Health and nutrition impact overall physical health and brain development, which consequently affect speech development. Proper nutrition supports brain development, while good health ensures that children have the energy and capability to engage in speech learning activities.

Cultural Factors:

- Cultural factors, including cultural norms and practices regarding communication, influence how and when children learn to speak. Different cultures may have varying expectations and methods of teaching language, which can shape the speech development process.

Educational Opportunities:

- Educational opportunities, such as access to early childhood education and language learning resources, can significantly impact speech development. Programs that focus on language skills and provide structured learning environments can enhance speech development outcomes.

Speech Abnormalities:

Speech abnormalities refer to atypical patterns or disruptions in speech production that deviate from normal speech development or functioning. These abnormalities can affect various aspects of speech, including articulation, fluency, voice quality, and the ability to form sounds, words,

and sentences effectively. Speech abnormalities can result from a range of causes, including neurological disorders, genetic conditions, developmental delays, hearing impairments, and psychological factors. They can manifest as difficulties in pronouncing words correctly (articulation disorders), disruptions in the flow of speech (fluency disorders like stuttering), abnormal pitch, volume, or quality of the voice (voice disorders), and problems with the rhythm and timing of speech (prosodic disorders).

Aphasia:

Aphasia is a communication disorder that results from damage to parts of the brain that are involved in language. This condition affects a person's ability to speak, understand speech, read, or write. Aphasia can occur suddenly after a stroke or head injury, or it can develop gradually from a slow-growing brain tumor or a disease that causes progressive, permanent damage (degenerative).

Types of Aphasia, each characterized by specific language deficits:

1. Broca's Aphasia (Non-Fluent Aphasia):

Language Deficits: Difficulty in speech production, writing, and forming complete sentences. Speech is typically halting and effortful, with poor grammar.

Characteristics: Patients understand speech relatively well and are often aware of their difficulties.

2. Wernicke's Aphasia (Fluent Aphasia)

Language Deficits: Problems with understanding spoken and written language. Speech is fluent but often nonsensical, with made-up words and irrelevant word substitutions.

Characteristics: Patients have difficulty grasping the meaning of words and sentences but can speak with normal grammar and syntax.

3. Global Aphasia:

Language Deficits: Severe impairment in both speech production and comprehension. Patients have very limited ability to speak or understand language.

Characteristics: This type of aphasia results from extensive damage to the brain's language areas and is often associated with other cognitive and physical deficits.

4. Anomic Aphasia:

Language Deficits: Difficulty in finding the right words, especially names of objects. Speech is fluent and grammatically correct but interrupted by frequent pauses and circumlocutions.

Characteristics: Patients understand speech well and can read adequately but struggle with naming and word retrieval.

Causes of Aphasia:

- The primary causes of aphasia include **stroke**, which is the most common cause, accounting for about 25-40% of all aphasia cases. Other causes include severe **head injuries, brain tumors, infections, or progressive neurological disorders such as Alzheimer's disease**.

Symptoms:

- Symptoms of aphasia vary depending on the part of the brain affected and the extent of the damage. Common symptoms include **difficulty in speaking, trouble finding the right words, speaking in short or incomplete sentences, using words in the wrong order, difficulty understanding conversations, problems with reading or writing, and difficulty with numbers or arithmetic**. Aphasia can be categorized into different types based on the symptoms and the location of the brain damage, such as Broca's aphasia, Wernicke's aphasia, and global aphasia.

Diagnosis:

- Diagnosis of aphasia typically involves a comprehensive assessment by a speech-language pathologist. This assessment includes a series of standardized tests that evaluate the individual's ability to speak, understand language, read, and write. **Medical imaging techniques such as MRI or CT scans are also used to identify the location and extent of brain damage**. Additionally, a thorough medical history and neurological examination help in diagnosing aphasia and identifying its underlying cause.

Treatment:

- Treatment for aphasia focuses on improving communication abilities and often involves **speech and language therapy**. This therapy is tailored to the individual's specific needs and may include exercises to improve language skills, alternative communication methods, and strategies to aid in daily communication. The involvement of family members and caregivers is crucial for supporting the person with aphasia in their recovery. In some cases, medications or surgery might be considered to treat the underlying cause of the brain damage. Ongoing support and therapy are essential for managing aphasia and enhancing the quality of life for those affected.

Apraxia:

Introduction:

Apraxia is a neurological disorder characterized by the inability to perform purposeful movements or gestures, despite having the desire and the physical capability to do so. This disorder **affects the brain's ability to plan and coordinate motor tasks**. It can manifest in various forms, including ideomotor apraxia (difficulty with simple tasks like waving goodbye), ideational apraxia (difficulty with complex tasks involving multiple steps), and buccofacial or orofacial apraxia (difficulty with movements of the face and mouth).

Two common types of Apraxia:

1. Apraxia of Speech (AOS):

Characteristics: Difficulty planning and coordinating speech movements, leading to inconsistent speech errors (e.g., sound distortions, substitutions).

Causes: Damage to the brain's motor planning areas, often due to stroke or brain injury.

2. Limb Apraxia:

Characteristics: Difficulty performing purposeful movements with arms and legs, such as gesturing or using tools correctly.

Causes: Damage to the parietal or frontal lobes, commonly due to stroke or brain injury.

Causes of Apraxia:

- The primary causes of apraxia include **brain damage** resulting from a stroke, head injury, brain tumor, or neurodegenerative diseases such as Alzheimer's disease or Parkinson's disease. It can also be associated with conditions like dementia or may occur alongside other motor disorders.

Symptoms:

- Symptoms of apraxia vary **depending on the type and severity of the disorder**. Common symptoms include difficulty in performing everyday tasks that require coordination, trouble with speech (speech apraxia), problems with fine motor skills like buttoning a shirt or using utensils, and difficulty with complex movements or sequences.

of movements. In children, developmental apraxia of speech can affect their ability to form words and sounds correctly.

Diagnosis:

- Diagnosis of apraxia typically involves a comprehensive **evaluation by a neurologist or a speech-language pathologist**. The evaluation includes a detailed medical history, physical and neurological examinations, and various tests to assess motor skills and the ability to perform specific tasks. Imaging studies such as MRI or CT scans may be used to identify areas of brain damage or abnormalities.

Treatment:

- Treatment for apraxia **focuses on improving motor function and communication abilities**. This often involves physical therapy, occupational therapy, and speech therapy, tailored to the individual's specific needs. Therapy may include exercises to improve coordination, strategies to compensate for motor deficits, and practice with specific tasks or movements. For those with speech apraxia, speech therapy aims to enhance articulation and communication skills. The involvement of family members and caregivers is crucial in providing support and encouragement during the rehabilitation process. In some cases, medications or other treatments may be used to address underlying conditions contributing to apraxia. Ongoing therapy and support are essential for managing apraxia and improving the individual's quality of life.

Wernicke's Aphasia:

Introduction:

Wernicke's aphasia, also known as **receptive aphasia**, is a type of aphasia where individuals have difficulty understanding spoken and written language, while their ability to produce speech remains relatively fluent but nonsensical. This condition results from damage to Wernicke's area, a region in the left hemisphere of the brain responsible for language comprehension.

Causes:

- The **primary cause** of Wernicke's aphasia is a stroke, particularly one that affects the left posterior temporal lobe. Other causes can include head injuries, brain tumors, infections, or neurodegenerative diseases such as Alzheimer's disease. Any condition that results in damage to Wernicke's area can potentially cause this type of aphasia.

Symptoms:

- Symptoms of Wernicke's aphasia include **fluent but nonsensical speech**, the use of made-up or inappropriate words, difficulty understanding spoken and written language, inability to recognize language errors, and problems with reading and writing. Individuals

with Wernicke's aphasia may speak in long, complete sentences that lack meaning or contain incorrect words or phrases, making their speech difficult to understand. They may also have trouble following conversations or understanding what others are saying.

Diagnosis:

- Diagnosis of Wernicke's aphasia involves a **thorough evaluation** by a speech-language pathologist, which includes a **series of standardized tests to assess language comprehension, production, and other cognitive functions**. Medical imaging techniques such as MRI or CT scans are used to identify the location and extent of brain damage. A detailed medical history and neurological examination also help in diagnosing the condition and determining its cause.

Treatment:

- Treatment for Wernicke's aphasia primarily involves **speech and language therapy** aimed at improving language comprehension and communication skills. Therapy may include exercises to enhance understanding of spoken and written language, **strategies to improve expressive language abilities, and alternative communication methods if needed**. Family involvement is essential in providing support and helping to practice communication strategies. Depending on the underlying cause, additional treatments such as medications or surgery may be considered to address the brain damage. Continuous therapy and support are crucial for managing Wernicke's aphasia and enhancing the quality of life for those affected.

Topics (Lateralization in speech, Brain areas in speech, Broca's Area, Wernicke's Area, Angular Gyrus)

Lateralization In Speech

Lateralization in speech refers to the distribution of language processing across the two hemispheres of the brain. In most people, language processing is lateralized to the left hemisphere, which means that the left hemisphere is dominant for language processing.

• **Left Hemisphere Dominance:**

- **Structure:** The left hemisphere has a larger cortical surface area dedicated to language processing, particularly in areas such as Broca's area and Wernicke's area.
- **Function:** The left hemisphere is specialized for language processing, including speech production, comprehension, and semantic processing.
- **Evolution:** Lateralization of language to the left hemisphere may have evolved to optimize language processing and reduce interference between language and other cognitive processes.
- The left hemisphere is responsible for processing most aspects of language, including:
 - **Speech production:** articulation, phonology, syntax
 - **Language comprehension:** semantic and syntactic processing
- The left hemisphere is dominant for language processing in approximately **95-98%** of the population

• **Right Hemisphere Involvement:**

- The right hemisphere is involved in processing certain aspects of language, including:
 - **Emotional processing:** emotional prosody, emotional expression
 - **Speech perception:** phonological processing, semantic processing
 - **Attentional processes:** focused attention, divided attention

- The right hemisphere plays a supporting role in language processing and is often involved in processing more nuanced or subtle aspects of language

- **Bilateral Processing:**

- Both hemispheres are involved in processing certain aspects of language, including:

- **Speech perception:** phonetic and semantic aspects

- **Language comprehension:** pragmatic and contextual aspects

- Bilateral processing allows for more efficient and effective language processing, as both hemispheres work together to facilitate communication

Understanding lateralization in speech is essential for addressing communication disorders like aphasia, which can result from damage to specific areas of the brain. By recognizing how language processing is distributed between the hemispheres, researchers and clinicians can develop more effective treatment strategies to support language recovery.

Brain Areas In Speech

The brain areas that involved in speech processing:

1. **Broca's area (BA 44 & 45):** Located in the frontal lobe, responsible for speech production, articulation, and phonology.

2. **Wernicke's area (BA 22):** Located in the temporal lobe, involved in speech comprehension, semantics, and syntactic processing.

3. **Primary auditory cortex (BA 41):** Processes basic auditory information.

4. **Angular gyrus (BA 39):** Involved in language processing, semantic processing, and attention.

5. **Prefrontal cortex (BA 46 & 47):** Supports working memory, attention, and executive functions essential for language processing.

6. **Primary motor cortex (BA 4):** Controls articulation and speech motor planning.

7. **Supplementary motor area (BA 6):** Involved in speech planning, articulation, and coordination.

These brain areas work together to enable us to understand and produce speech. Damage to any of these areas can result in specific language processing deficits, like aphasias.

Broca's Area

Broca's area, located in the frontal lobe of the left hemisphere (**BA 44 and 45**), is a critical region for speech processing and language production. It is named after French neurologist **Paul Broca**, who first described its role in speech in the **19th century**.

Functions:

- **Speech production:** Broca's area is responsible for articulation, phonology, and syntax. It coordinates the movement of the lips, tongue, and jaw to produce speech sounds.
- **Language processing:** Involved in lexical-semantic processing, grammatical processing, and sentence construction.
- **Working memory:** Plays a role in holding and manipulating linguistic information in working memory.
- **Executive functions:** Supports cognitive processes like attention, planning, and error monitoring.

Damage To Broca's Area Can Result In:

- **Broca's Aphasia:** Characterized by difficulty speaking in grammatically correct sentences, struggling to find the right words, and distorted speech articulation.
- **Agrammatism:** Difficulty with sentence construction and grammatical errors.
- **Anomia:** Difficulty naming objects or people.

In summary, Broca's area is essential for transforming thoughts into spoken language, enabling us to communicate effectively.

Wernicke's Area

Wernicke's area is a critical region in the left hemisphere of the brain, located in the **superior temporal gyrus (BA 22)**, that plays a central role in language processing, particularly in comprehension. It is named after German neurologist **Carl Wernicke**, who first described its function in the **late 19th century**.

Functions:

- **Language comprehension:** Wernicke's area is responsible for understanding spoken language, including semantic processing (meaning), syntactic processing (grammar), and lexical processing (vocabulary).
- **Speech perception:** Involved in processing basic auditory information and phonological processing (sound structure).
- **Semantic processing:** Plays a role in accessing word meanings, object recognition, and conceptual processing.

- **Working memory:** Supports holding and manipulating linguistic information in working memory.

Damage To Wernicke's Area Can Result In:

- **Wernicke's aphasia:** Characterized by difficulty understanding spoken language, speaking in complete but nonsensical sentences, and using made-up words.

- **Receptive aphasia:** Difficulty comprehending language, including spoken and written language.

- **Anomia:** Difficulty naming objects or people.

Wernicke's area works closely with Broca's area to enable us to understand and produce language. While Broca's area is responsible for language production, Wernicke's area is essential for language comprehension.

Angular Gyrus (AG)

The Angular Gyrus (AG) is a region in the brain located in the parietal lobe, specifically in the inferior parietal lobe (IPL). It plays a crucial role in various cognitive processes, including:

1. Language processing:

- The AG is involved in language comprehension, particularly in semantic processing, syntactic processing, and lexical-semantic processing.
- It helps to comprehend written and spoken language, accessing word meanings and integrating context.
- The AG is also involved in language production, including writing and speaking.

2. Number processing:

- The AG is essential for numerical cognition, including number comprehension, calculation, and mathematical reasoning.

- It helps to understand quantity, magnitude, and numerical relationships.
- The AG is involved in mental arithmetic and mathematical problem-solving.

3. **Spatial Attention:**

- The AG is involved in directing attention to specific locations in space, tracking objects, and understanding spatial relationships.
- It plays a key role in spatial reasoning, mental rotation, and navigation.
- The AG helps to focus attention on relevant sensory information and filter out irrelevant information.

4. **Memory:**

- The AG is involved in directing attention to specific locations in space, tracking objects, and understanding spatial relationships.
- It plays a key role in spatial reasoning, mental rotation, and navigation.
- The AG helps to focus attention on relevant sensory information and filter out irrelevant information.

5. **Multimodal integration:**

- The AG integrates information from different sensory modalities, such as vision, hearing, and touch.
- It helps to combine sensory inputs to understand the world around us.
- The AG is involved in cross-modal processing, such as reading lips and understanding spoken language.


Damage to the Angular Gyrus can result in:

- **Agnosia:** Difficulty recognizing objects or understanding written language
- **Acalculia:** Difficulty with mathematical calculations
- **Spatial neglect:** Difficulty paying attention to one side of space
- **Memory impairments:** Difficulty with working memory and learning new information

The Angular Gyrus is a critical region for integrating information from multiple cognitive systems, and its damage can lead to significant cognitive impairments.



Speech Areas



Damage In Speech Area

