

Differences in Toxic Chemical Release Pathways: A Comparison of Air and Water Releases

Kira Lu

2026-05-02

Contents

1	Introduction	1
2	Methods	2
3	Results	2
4	Conclusion	4
5	Limitation	5
6	GitHub Repository	5
7	AI Use Disclosure	5
8	References	5

1 Introduction

1.1 Research Question

This study investigates whether the magnitude of toxic chemical releases differs between air and water release pathways.

1.2 Scientific Background

Environmental exposure to toxic chemicals varies by release pathway, with air and water pathways presenting different risks to human health and ecosystems. Previous studies have shown that toxic exposure may be associated with adverse health outcomes, particularly in vulnerable populations.^{1,2} The U.S. Environmental Protection Agency's Toxics Release Inventory (TRI) provides a comprehensive dataset to examine these patterns across industrial facilities.³

2 Methods

2.1 Data

The data used in this study come from the U.S. Environmental Protection Agency (EPA) Toxics Release Inventory (TRI) 2024 dataset. Due to GitHub file size limitations, a subset of the dataset (`tri_small.csv`) was used.

The TRI dataset includes facility-level records reported by industrial facilities to the EPA. The analytic dataset included 5,000 facility records with non-missing values for both air and water releases.

The key variables include:

- **air_releases:** sum of fugitive and stack air emissions
- **water:** total water releases

A long-format dataset was created to compare the two release pathways. A log transformation ($\log(1 + \text{releases})$) was applied to reduce skewness and improve comparability between distributions.

2.2 Statistical Analysis

To compare release amounts between pathways, a Wilcoxon rank-sum test was used because the release data are highly skewed and do not satisfy the assumptions of normality required for a two-sample t-test.

The hypotheses were:


- **H0:** There is no difference in toxic release amounts between air and water pathways.
- **H1:** There is a difference in toxic release amounts between air and water pathways.


3 Results

3.1 Visualization

The following histogram and boxplot were used to visualize the distribution of log-transformed toxic releases by pathway.

Distribution of Log-Transformed Toxic Releases by Pathway





3.2 Statistical Test

A Wilcoxon rank-sum test was conducted to assess differences in log-transformed toxic release amounts between air and water pathways. The test revealed a statistically significant difference between the two pathways ($W = 19635229$, $p < 2.2e-16$).

3.3 Interpretation

The distributions of log-transformed toxic releases were visualized using histograms and boxplots. The results show that water releases are highly concentrated near zero with several extreme outliers, while air releases exhibit a broader distribution.

Descriptive patterns suggest that air releases have a higher median and greater variability compared to water releases.

The Wilcoxon rank-sum test indicated a statistically significant difference between pathways ($W = 19635229$, $p < 2.2e-16$), suggesting that release distributions differ between air and water pathways.

4 Conclusion

This analysis demonstrates that toxic release amounts differ significantly between air and water pathways in the TRI dataset. Air releases show a broader distribution, while water releases are concentrated near zero with extreme outliers. These findings suggest that release pathway is strongly associated with differences in both the magnitude and variability of toxic releases.

5 Limitation

One limitation of this analysis is that it relies on a subset of the TRI dataset, which may not fully represent all facilities or geographic variation. In addition, the analysis does not account for differences in facility size or industry type, which may influence the magnitude of reported releases.

6 GitHub Repository

The GitHub repository containing the data, scripts, and related project materials is available here:

<https://github.com/puffconw/r-project-test>

7 AI Use Disclosure

This document was developed with assistance from ChatGPT for code debugging, formatting, and preparation of the R Markdown structure. All analytical decisions, statistical interpretation, and final writing were reviewed and completed by the author.

8 References

1. Agarwal N, Banerghansa C, Bui LTM. Toxic exposure in America: estimating fetal and infant health outcomes from 14 years of TRI reporting. *J Health Econ.* 2010;29(4):557-574.
2. Wilson SM, Fraser-Rahim H, Williams E, et al. Assessment of the distribution of toxic release inventory facilities in metropolitan Charleston: an environmental justice case study. *Am J Public Health.* 2012;102(10):1974-1980.
3. U.S. Environmental Protection Agency. Toxics Release Inventory (TRI) Program. Accessed April 10, 2026. <https://www.epa.gov/toxics-release-inventory-tri-program>