

# Assignment 16

Pulkit Saxena

## 1 PROBLEM HOFFMAN PG 242 Q7

Find the minimal polynomials and the rational forms of the following real matrices

$$1) \begin{pmatrix} 0 & -1 & -1 \\ 1 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}$$

$$2) \begin{pmatrix} c & 0 & -1 \\ 0 & c & 1 \\ 1 & 1 & c \end{pmatrix}$$

$$3) \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$$

## 2 THEOREMS

Theorem 1	<p>A Rational canonical form is a matrix <math>\mathbf{R}</math> that is Direct sum of companion matrix.</p> <p><math>\mathbf{R} = \mathbf{C}(\mathbf{p}_1) \oplus \cdots \oplus \mathbf{C}(\mathbf{p}_r)</math></p> $\mathbf{R} = \begin{pmatrix} \mathbf{C}(\mathbf{p}_1) & 0 & 0 & \cdots & 0 \\ 0 & \mathbf{C}(\mathbf{p}_2) & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & 0 & \cdots & \mathbf{C}(\mathbf{p}_r) \end{pmatrix} \quad (2.0.1)$ <p>where <math>\mathbf{C}(\mathbf{p}_i)</math> is the <math>k_i \times k_i</math> companion matrix of <math>p_i</math> where polynomial <math>p_1, p_2 \dots p_r</math> are called invariant factors for Given Matrix .Where <math>k_i</math> denotes the degree of annihilator of <math>p_i</math>.</p> <p>This representation is called rational form.</p>
Theorem 2	<p>If <math>p_i(x) = x + a_0</math> then its companion matrix <math>\mathbf{C}(\mathbf{p})</math> is 1 x 1 matrix as <math>(-a_0)</math>.</p> <p>If <math>k_i \geq 2</math> then <math>p(x) = x^k + a_{k-1}x^{k-1} + \cdots + a_1x + a_0</math> then its companion matrix is</p> $\mathbf{C}(\mathbf{p}_i) = \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 & -a_0 \\ 1 & 0 & 0 & \cdots & 0 & -a_1 \\ 0 & 1 & 0 & \cdots & 0 & -a_2 \\ 0 & 0 & 1 & \cdots & 0 & -a_3 \\ \vdots & \vdots & \vdots & \cdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & -a_{k-1} \end{pmatrix} \quad (2.0.2)$

TABLE 1: Illustration of theorem.

## 3 SOLUTION

Given part 1	$\mathbf{A} = \begin{pmatrix} 0 & -1 & -1 \\ 1 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}$
Characteristics and Minimal Polynomial	<p>Characteristics polynomial of the matrix is <math>\det(x\mathbf{I} - \mathbf{A})</math></p> $\det(x\mathbf{I} - \mathbf{A}) = \begin{vmatrix} (x) & 1 & 1 \\ -1 & (x) & 0 \\ 1 & 0 & (x) \end{vmatrix} = x(x^2) - 1(-x) - x = x^3$ <p>Characteristic Polynomial = <math>x^3</math></p> <p>Minimal Polynomial can be <math>x, x^2</math> or <math>x^3</math> of lowest degree satisfying <math>p(\mathbf{A}) = 0</math></p> <p>Let take <math>p(x) = x \implies p(\mathbf{A}) = \mathbf{A} = \begin{pmatrix} 0 &amp; -1 &amp; -1 \\ 1 &amp; 0 &amp; 0 \\ -1 &amp; 0 &amp; 0 \end{pmatrix} \neq 0</math></p> <p>Let take <math>p(x) = x^2 \implies p(\mathbf{A}) = \mathbf{A}^2 = \begin{pmatrix} 0 &amp; -1 &amp; -1 \\ 1 &amp; 0 &amp; 0 \\ -1 &amp; 0 &amp; 0 \end{pmatrix} \neq 0</math></p> <p>Take <math>p(x) = x^3 \implies p(\mathbf{A}) = \mathbf{A}^3 = \begin{pmatrix} 0 &amp; 0 &amp; 0 \\ 0 &amp; 0 &amp; 0 \\ 0 &amp; 0 &amp; 0 \end{pmatrix} = \mathbf{0}</math></p> <p>Thus minimal polynomial <math>p(x) = x^3</math>.</p>
Companion Matrix and Rational Form	<p>Since</p> <p>Characteristics polynomial=Minimal polynomial=Invariant factors</p> $p(x) = x^3 + 0x^2 + 0x + 0$ <p>So Companion Matrix is of dimension 3x3 and from theorem 2</p> $\mathbf{C}(\mathbf{p}) = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$ <p>Since there is only one minimal polynomial of degree 3 which is equal to characteristics equation therefore</p> <p>Rational matrix=companion matrix</p>

	$\mathbf{R} = \mathbf{C}(\mathbf{p}) = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$ <p>which is in rational form.</p>
Given Part 2	$\mathbf{A} = \begin{pmatrix} c & 0 & -1 \\ 0 & c & 1 \\ 1 & 1 & c \end{pmatrix}$
Characteristics and Minimal Polynomial	<p>Characteristics polynomial of the matrix is <math>\det(x\mathbf{I} - \mathbf{A})</math></p> $\det(x\mathbf{I} - \mathbf{A}) = \begin{vmatrix} (x-c) & 0 & 1 \\ 0 & (x-c) & -1 \\ -1 & -1 & (x-c) \end{vmatrix}$ $= (x-c)((x-c)^2 + 1) - (x-c) = (x-c)^3$ <p>Characteristic Polynomial = <math>(x-c)^3</math></p> <p>Minimal Polynomial can be <math>(x-c)</math>, <math>(x-c)^2</math> or <math>(x-c)^3</math></p> <p>of lowest degree satisfying <math>p(\mathbf{A}) = 0</math></p> <p>Let take <math>p(x) = (x-c) \implies p(\mathbf{A}) = \mathbf{A} - c\mathbf{I} = \begin{pmatrix} 0 &amp; 0 &amp; -1 \\ 0 &amp; 0 &amp; 1 \\ 1 &amp; 1 &amp; 0 \end{pmatrix} \neq 0</math></p> <p>Let take <math>p(x) = (x-c)^2 \implies p(\mathbf{A}) = (\mathbf{A} - c\mathbf{I})^2 = \begin{pmatrix} -1 &amp; -1 &amp; 0 \\ 1 &amp; 1 &amp; 0 \\ 0 &amp; 0 &amp; 0 \end{pmatrix} \neq 0</math></p> <p>Take <math>p(x) = (x-c)^3 \implies p(\mathbf{A}) = (\mathbf{A} - c\mathbf{I})^3 = \begin{pmatrix} 0 &amp; 0 &amp; 0 \\ 0 &amp; 0 &amp; 0 \\ 0 &amp; 0 &amp; 0 \end{pmatrix} = \mathbf{0}</math></p> <p>Thus minimal polynomial <math>p(x) = (x-c)^3</math>.</p>
Companion Matrix and Rational Form	<p>Since</p> <p>Characteristics polynomial=Minimal polynomial=Invariant factors</p> $p(x) = x^3 - 3cx^2 + 3c^2x - c^3$ <p>So Companion Matrix is of dimension 3x3 and from theorem 2</p>

	$\mathbf{C}(\mathbf{p}) = \begin{pmatrix} 0 & 0 & c^3 \\ 1 & 0 & -3c^2 \\ 0 & 1 & 3c \end{pmatrix}$ <p>Since there is only one minimal polynomial of degree 3 which is equal to characteristics equation therefore Rational matrix=companion matrix</p> $\mathbf{R} = \mathbf{C}(\mathbf{p}) = \begin{pmatrix} 0 & 0 & c^3 \\ 1 & 0 & -3c^2 \\ 0 & 1 & 3c \end{pmatrix}$ <p>which is in rational form.</p>
Given part 3	$\mathbf{A} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$
Characteristics and Minimal Polynomial	<p>Characteristics polynomial of the matrix is <math>\det(x\mathbf{I} - \mathbf{A})</math></p> $\det(x\mathbf{I} - \mathbf{A}) = \begin{vmatrix} (x - \cos \theta) & -\sin \theta \\ \sin \theta & (x - \cos \theta) \end{vmatrix} = x^2 - 2 \cos \theta x + 1$ <p>Characteristic Polynomial = Minimal Polynomial = <math>x^2 - 2 \cos \theta x + 1</math></p>
Companion Matrix and Rational Form	<p>Since</p> <p>Characteristics polynomial=Minimal polynomial=Invariant factors</p> $p(x) = x^2 - 2 \cos \theta x + 1$ <p>So Companion Matrix is of dimension 2x2 and from theorem 2</p> $\mathbf{C}(\mathbf{p}) = \begin{pmatrix} 0 & -1 \\ 1 & 2 \cos \theta \end{pmatrix}$ <p>Since there is only one minimal polynomial of degree 3 which is equal to characteristics equation therefore Rational matrix=companion matrix</p> $\mathbf{R} = \mathbf{C}(\mathbf{p}) = \mathbf{C}(\mathbf{p}) = \begin{pmatrix} 0 & -1 \\ 1 & 2 \cos \theta \end{pmatrix}$ <p>which is in rational form.</p>

TABLE 2: Solution Table