Bloch-McConnell Simulation Study – Comparison of Various CEST Simulation Frameworks

Markus Huemer[1,2,3], Patrick Schuenke[4,5], Viktoria Buchegger[1], Martin Juschitz[1], Nick Scholand[1], Felix F. Zimmermann[4], Christoph Kolbitsch[4], Moritz Simon Fabian[6], Nikita Vladimirov[7], Or Perlman[7,8], Swee Qi Pan[9,10], Yee Kai Tee[9], Ouri Cohen[11], Huabin Zhang[12], Jianpan Huang[12], Jian Hou[13], Qing Zeng[14,15], Nirbhay Yadav[14,15], Jiadi Xu[14,15], Kexin Wang[14,15,16], Beomgu Kang[14,15], Munendra Singh[14,15], Hye-Young Heo[14,15], Jingpu Wu[15,17], Jinyuan Zhou[15], Dario Livio Longo[18], Feriel Ramdhane[18], Zhongliang Zu[19], Malvika Viswanathan[19], Pedram Parnianpour[20], Erik P. Pioro[20], Mara Quach[21,22], Rudolf Stollberger[1], Martin Uecker[1], Moritz Zaiss[6]

[1]Institute of Biomedical Imaging, Graz University of Technology, Graz, Austria
[2]Athinoula A. Martinos Center for Biomedical Imaging, Department of Radiology, Massachusetts General Hospital, Charlestown, USA
[3]Department of Radiology, Harvard Medical School, Boston, USA
[4]Physikalisch-Technische Bundesanstalt (PTB), Braunschweig and Berlin, Germany
[5]NVision Quantum Technologies, Ulm, Germany
[6]Institute of Neuroradiology, Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU), University Hospital Erlangen, Erlangen, Germany
[7]School of Biomedical Engineering, Tel Aviv University, Tel Aviv, Israel
[8]Sagol School of Neuroscience, Tel Aviv University, Tel Aviv, Israel
[9]Lee Kong Chian Faculty of Engineering and Science, Universiti Tunku Abdul Rahman, Kajang, Malaysia
[10]Philips Singapore, Singapore, Singapore
[11]Department of Medical Physics, Memorial Sloan Kettering Cancer Center, New York, USA
[12]Department of Diagnostic Radiology, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong
[13]Department of Imaging and Interventional Radiology, The Chinese University of Hong Kong, Hong Kong
[14]F.M. Kirby Research Center for Functional Brain Imaging, Kennedy Krieger Institute, Baltimore, USA
[15]Department of Radiology and Radiological Science, Johns Hopkins University School of Medicine, Baltimore, USA
[16]Biomedical Imaging Research Institute, Cedars-Sinai Medical Center, Los Angeles, California, USA
[17]Department of Electrical and Computer Engineering, Johns Hopkins University, Baltimore, USA
[18]Institute of Biostructures and Bioimaging, National Research Council of Italy, Torino, Italy
[19]Institute of Imaging Science, Vanderbilt University Medical Center, Nashville, USA
[20]Djavad Mowafaghian Centre for Brain Health, University of British Columbia, Vancouver, Canada
[21]Department of Biomedical Engineering and Graeme Clark Institute, The University of Melbourne, Parkville, Australia
[22]Melbourne Brain Centre Imaging Unit, The University of Melbourne, Parkville, Australia

October 30, 2026

Abstract

Purpose: Quantitative chemical exchange saturation transfer (CEST) MRI depends on Bloch-McConnell simulations, yet it is unclear how consistent implementations are across research groups. This study quantified inter-framework variability and identified dominant sources of disagreement.

Methods: Eight benchmark cases were defined and distributed online in two rounds: four continuous-wave preparations on a two-pool creatine and a five-pool white-matter model

(cases 1–4), followed by Gaussian and pulsed WASABI preparations (cases 5–8). Nineteen groups submitted Z-spectra for comparison.

Results: For simple two-pool CW saturation (cases 1–2), most submissions agreed within $\sim 10^{-4}$ in Z-spectra difference, but residual spread exceeded numerical precision. By contrast, marked differences were observed in the simulations for the five-pool model (cases 3–4) and time-dependent saturation pulses (cases 5–7). The different implementation of the magnetization transfer (MT) pool was identified as a major contributor to these differences.

Good agreement was achieved for a single Gaussian pulse (case 5); pulse trains (cases 6–7) displayed larger differences, related to the interpulse delays. The two-pulse WASABI scheme (case 8) showed similar problems.

The final observed differences in the simulations could lead to biases of tens of percent in the fitted pool fractions or exchange rates.

Conclusion: Although the presented study helped to find common ground, Bloch-McConnell simulators from different groups are not yet interchangeable, even for fully specified protocols. Harmonized MT modeling (full M_{xyz} versus z-only MT), shaped-pulse handling, and clear RF power definitions are prerequisites for reproducible quantitative CEST. This open benchmark supports ongoing comparison and community standardization.

Keywords: CEST, MRI, Bloch-McConnell, magnetization transfer, quantitative MRI, inter-site comparison

Correspondence: Markus Huemer, Athinoula A. Martinos Center for Biomedical Imaging, 149 Thirteenth Street, Charlestown, USA. Email: `mhuemer@mgh.harvard.edu`

Wordcount: Abstract: 250, Main text: 5241

1 Introduction

Chemical Exchange Saturation Transfer (CEST) MRI is a powerful imaging technique that provides contrast sensitive to exchange processes between solute and water protons. This sensitivity enables novel contrasts that are increasingly explored in clinical MRI applications. Among the most commonly studied CEST contrasts is amide proton transfer weighted (APTw) imaging, which has been used in various organs and has been shown to be particularly useful in the assessment of cancer [1], neurodegenerative diseases [2], and other pathologies like ischemic stroke [3]. In the context of cancer, CEST MRI has been used to differentiate tumor progression from radiation necrosis [4], assess treatment response [5], and detect genetic mutations such as IDH in gliomas [6]. Recently, there has been growing interest in quantitative CEST approaches (qCEST), which aim to provide quantitative insights into exchange rates and solute concentrations. Such approaches include Bloch-McConnell fitting, ratiometric methods, and CEST MR fingerprinting. Both, Bloch-McConnell fitting and CEST MRF are based on the Bloch-McConnell (BMC) equations, which extend the classical Bloch equations to systems of coupled proton pools with chemical exchange [7]. For a system of N pools, the magnetization vector $\mathbf{M}$ contains the transverse and longitudinal components of each pool. During RF irradiation at offset $\Delta\omega$, the time evolution is described by

$$\dot{\mathbf{M}} = \mathbf{A}(\Delta\omega, B_1, \ldots)\mathbf{M} + \mathbf{c}, \tag{1}$$

where $\mathbf{A}$ encodes relaxation, chemical exchange, off-resonance effects, and RF saturation, and $\mathbf{c}$ accounts for equilibrium magnetization. By simulating this evolution for a defined saturation scheme and a range of offsets, one obtains model Z-spectra and derived contrasts such as magnetization transfer ratio asymmetry (MTR_{asym}) [8].

A Bloch-McConnell simulation evaluates Eq. 1 for a chosen pool model and saturation scheme. In practice this requires assembling the system matrix $\mathbf{A}$ on each time interval, representing the RF waveform (at some frequency offset, B_1 scaling, and sampling or interpolation of the waveform), and propagating $\mathbf{M}$ with a numerical solver. Common choices include a matrix exponential of $\mathbf{A}$ (via Padé approximation or eigenvalue decomposition), ODE integrators, and

operator-splitting methods. These approaches can agree when applied carefully, but they differ in time-step or tolerance policy, how piecewise-constant RF intervals are formed, and in convergence. Independently developed frameworks may therefore disagree even for an identical protocol, if the RF is discretized differently, the offset is not reset during interpulse delays, or the MT pool is formulated as z-only MT rather than full M_{xyz}.

Because Bloch-McConnell model fitting and fingerprinting pipelines depend directly on these forward simulations, differences between implementations can translate into biases in estimated pool sizes and exchange rates. Numerous simulation frameworks have been developed to perform Bloch-McConnell simulations, ranging from tools used exclusively within individual research groups to open-source platforms such as Pulseq-CEST [9], BMCTool [10], and CEST GUI [11]. Despite the availability of these tools, no systematic comparison between different simulation frameworks has been conducted. Without a shared reference or ground truth, it remains difficult to judge whether observed variability in qCEST metrics derived from BMC simulations reflects biology, acquisition, or simulation implementation, which makes cross-study comparisons challenging.

The objective of this work is to address this gap by providing a baseline to compare Bloch-McConnell simulations from various CEST research groups. To achieve this, we defined two rounds of Bloch-McConnell simulation scenarios, each comprising four cases with well-defined saturation schemes and pool models. These cases were designed to capture a range of commonly encountered CEST saturation settings, including simple CW preparations, shaped-pulse trains as used on clinical MRI scanners, and pool configurations ranging from simple phantoms to brain tissue. Using this benchmark, we compared simulation results from up to 20 different groups, offering new insights into the consistency and reliability of Bloch-McConnell simulations across the field.

2 Methods

All necessary simulation settings and information were provided on the public GitHub repository at `https://github.com/pulseq-cest/BMsim_challenge`. The dataset used for the figures and analyses in this manuscript is archived on Zenodo at `https://doi.org/10.5281/zenodo.21265597`.

2.1 General Simulation Settings

The following settings applied to all benchmark cases unless stated otherwise.

- Fully relaxed initial longitudinal magnetization ($Z_i = 1$) before simulating each offset. Equivalently, the recovery interval between offsets must satisfy $T_{\text{rec}} > 10 \cdot T_1$ of the slowest pool (e.g. water with $T_1 = 3$ s in the 2-pool model requires $T_{\text{rec}} > 30$ s). All pool-model YAML files specify `reset_init_mag: True` for this purpose. The shorter recovery delays $T_{\text{rec}} = 3.5$ s (cases 1–3, 5–7) or 3 s (case 4) in the reference Pulseq files are *not* part of the simulation specification; they reflect typical scanner timing between acquisitions and must not be used as a substitute for resetting $Z_i = 1$.
- Relative concentration of water is set to one ($f_{water} = 1$)
- After each saturation a delay of 6.5 ms was simulated to account for the spoiler gradient, which would normally be applied in measurements (post-preparation delay $t_{\text{delay}} = 6.5$ ms).
- Gyromagnetic ratio: 42.5764 MHz/T, which leads to a Larmor frequency (for 3 T) of 127.7292 MHz
- Normalization scan at −300 ppm using the same preparation

2.2 Pool Models

Two different pool models were defined and used in the eight simulation cases. The first pool model is a 2-pool creatine model consisting of water and one CEST pool, mimicking a creatine phantom [12]. The second pool model is a 5-pool model consisting of a water pool, a semisolid magnetization transfer (ssMT) pool, an amide-like CEST pool, a guanidine-like CEST pool and a rNOE pool, in order to mimic more complex in vivo like environments, using a simplified parameterization inspired by [13]. The pool model configuration files were defined in human- and machine readable yaml file format and can be found in the GitHub repository. Numerical values are listed in Table 1.

Table 1: Pool-model parameters used in the benchmark. f is the relative pool size, T_1 and T_2 are the longitudinal and transverse relaxation times, k is the exchange rate to water, and δ_ω is the chemical shift relative to water.

Model	Pool	f	T_1 (s)	T_2 (s)	k (Hz)	δ_ω (ppm)
2-pool	water	1	3	2	–	0
	creatine	5×10^{-4}	1.05	0.1	50	1.9
5-pool	water	1	1	0.04	–	0
	ssMT	0.1351	1	4×10^{-5}	30	−3
	amide	9.009×10^{-4}	1	0.1	50	3.5
	guanidine	9.009×10^{-4}	1	0.1	1000	2
	NOE	4.5×10^{-3}	1.3	0.005	20	−3

2.3 Simulation Cases

In total, eight cases of increasing complexity were compared in two rounds on the pool models described in Section 2.2. An overview with the parameters for all cases is listed in Table 2, the saturation pulses are shown in Figure 1 (left column) and example Z-spectra for all cases are shown in Figure 1 (middle and right columns). These spectra were generated using the MATLAB implementation of Pulseq-CEST [9] and are equivalent to submission S01.

2.3.1 Round 1

In the first round of the challenge, four cases with continuous-wave (CW) saturation and different saturation times were tested. The aim was to cover cases that should first be comparatively straightforward to reproduce and then add complexity from shorter saturation times, additional exchanging pools, and MT modeling afterwards.

- **Case 1:** A 15 s CW saturation on the 2-pool creatine model with $B_1 = 2\,\mu T$. This case tests the steady-state for a simple creatine phantom and was intended as a baseline test for exchange, relaxation, offset handling, and sign conventions.
- **Case 2:** A 2 s CW saturation on the same 2-pool model with $B_1 = 2\,\mu T$. Compared with Case 1, this case emphasizes transient saturation behavior while keeping the pool model unchanged, thereby testing whether implementations agree away from the full steady-state.
- **Case 3:** The same 2 s CW saturation as in Case 2, but applied to the 5-pool white-matter model. This case tests the handling of multiple exchanging pools and, in particular, the implementation of the semisolid MT pool.
- **Case 4:** A short block-pulse preparation as used in the WASABI method [14], consisting of a 5 ms pulse with $B_1 = 3.7\,\mu T$ applied to the 5-pool white-matter model. This case probes short-pulse and off-resonance behavior in a multi-pool system, including the influence of MT and phase evolution during a brief RF event.

Table 2: Overview over the different simulation cases. CW block pulses (cases 1–4) use constant B_1 amplitude; WASABI block pulses (cases 4 and 8) use $B_{1,\mathrm{peak}}$; Gaussian pulses (cases 5–7) use $B_{1,\mathrm{rms}}$.

Case #	**Pool model**	**# Pulses**	**Pulse shape**	**Pulse duration**	**Interpulse delay**	B_1	**Offsets**
1	2-pool	1	block	15 s	-	2 µT	-15:0.1:15 ppm
2	2-pool	1	block	2 s	-	2 µT	-15:0.1:15 ppm
3	5-pool	1	block	2 s	-	2 µT	-15:0.1:15 ppm
4	5-pool	1	block	5 ms	-	3.7 µT	-2:0.05:2 ppm
5	2-pool	1	Gaussian	50 ms	-	1.996 µT (rms)	-2:0.02:2 ppm
6	2-pool	36	Gaussian	50 ms	5 ms	1.996 µT (rms)	-15:0.1:15 ppm
7	5-pool	36	Gaussian	50 ms	5 ms	1.996 µT (rms)	-15:0.1:15 ppm
8	5-pool	2	block	5 ms	100 µs	3.7 µT	-2:0.05:2 ppm

2.3.2 Round 2

In the second round, pulsed saturation using Gaussian and block-pulse preparations was compared on the same models. These cases were designed to test whether groups implemented shaped RF pulses, interpulse delays, and pulse trains consistently when the saturation preparation is provided in a machine-readable format.

- **Case 5:** A single Gaussian pulse with $t_p = 50$ ms and $B_{1,\mathrm{rms}} = 1.9962\,\mu T$ on the 2-pool model. The pulse shape was provided as a CSV file with time and amplitude values. This isolated pulse tests interpolation, pulse handling and RF amplitude scaling, without additional complications from a pulse train.

- **Case 6:** The Gaussian pulse from Case 5 repeated 36 times with an interpulse delay $t_d = 5$ ms, for a combined T_{sat} of 1.975 s, again on the 2-pool model. This case extends Case 5 to a clinically relevant pulse train and tests accumulation of saturation and relaxation over repeated RF events.

- **Case 7:** The same Gaussian pulse train as in Case 6, but applied to the 5-pool white-matter model. This case combines shaped-pulse train implementation with the multi-pool and MT-modeling complexity introduced in Case 3.

- **Case 8:** The WASABI pulse from Case 4 ($B_1 = 3.7\,\mu T$) played out twice with an interpulse delay $t_d = 100\,\mu s$ on the 5-pool white-matter model. This short two-pulse preparation tests whether implementations handle phase accumulation and relaxation during very short delays consistently.

2.4 Entries and participating groups

In total, 20 groups or solver teams participated in the study. Some groups submitted multiple solver variants or additional z-only MT simulations. Additionally, some groups did not submit results for all cases, leading to the following number of submitted entries per case:

- **Case 1:** 22 entries

- **Case 2:** 23 entries

- **Case 3:** 24 entries
- **Case 4:** 23 entries
- **Case 5:** 18 entries
- **Case 6:** 19 entries
- **Case 7:** 19 entries
- **Case 8:** 19 entries

Fewer entries were received for Cases 5–8 than for the CW cases. There was a time gap between the two challenge rounds, and shaped-pulse simulations may be of limited interest for groups that work mostly in the pre-clinical setting with continuous-wave saturation, which likely explains why some participants did not submit Round 2 results. Further, some additional settings for solvers were tested for the earlier cases and not submitted for the later cases. Additional information on the participating groups, solver variants, and implementation details is provided in Table 3 and Supporting Information Section 1.

The main comparison figures exclude entries labeled as z-only MT simulations, except for the dedicated M_{xyz} versus z-only MT comparison in Section 3.2.

Table 3: Overview of participating groups, solver variants, and implementation details. Submission numbers are used as labels in the comparison figures and were assigned in the order of submission.

Subm.	Corresponding Participant	Prog. Language	Number of Pools	MT	Solver	Paper & Code
S01	**Moritz Zaiss - FAU**	Matlab 2017b / C++	dynamic	x,y,z	expm with Pade (6 iterations)	pulseq-cest matlab
S02	**Patrick Schuenke - PTB**	Python	dynamic	x,y,z	expm with eig	BMCTool - `github.com/schuenke/BMCTool`
S03	**Markus Huemer - IBI TUG**	Matlab	5 fixed	x,y,z	expm with Operator Splitting	`gitlab.tugraz.at/ibi/mrirecon/miscellaneous/bmc-simulation-matlab`[15]
S04	**Markus Huemer - IBI TUG**	Matlab	7 fixed	x,y,z	expm with eig	
S05	**Qing Zeng - KKI/JHU**	Python	dynamic	x,y,z	expm	
S06	**Nikita Vladimirov - TAU**	Python/C++	dynamic	x,y,z	Pade	open-py-cest-mrf [16] `github.com/momentum-laboratory/molecular-mrf`
S07	**Zhongliang Zu - VUMC**	Matlab	dynamic	z-only	expm with eig	
S08	**Kexin Wang - KKI/JHU**	Matlab	fixed	x,y,z	expm	`github.com/Kexin-Wang/BMsim`[17]
S09	**Dario Longo - CNR**	Matlab 2021b	fixed	x,y,z	expm	extension of `github.com/cest-sources/BM_sim_fit`[18]
S10	**Moritz Fabian - FAU**	Matlab 2019b/2022b	dynamic	x,y,z	RKDP	`gitlab.com/mrilab/sim/bmsim.git`
S11	**Beomgu Kang - KKI/JHU**	Matlab / Python	dynamic	x,y,z	expm	`github.com/Heo-Group`[19]
S12	**Huabin Zhang - HKU**	Matlab 2023a	dynamic	x,y,z	expm	`github.com/JianpanHuang/BMsim_challenge`
S13	**Jian Hou - CUHK**	Matlab 2023a	dynamic	z-only	expm	
S14	**Viktoria Buchegger - IBI TUG**	C	fixed	x,y,z	RKDP	BART [20] `codeberg.org/mrirecon/bart`
S15	**Mara Quach - UniMelb**	Julia v1.11	dynamic	x,y,z	Rosenbrock	`github.com/maraquach/CESTSim.jl`
S16	**Felix F. Zimmermann - PTB**	Python (torch)	dynamic	x,y,z	expm	`github.com/fzimmermann89/mr2`
S17	**Ouri Cohen - MSK**	pytorch	dynamic	x,y,z	expm	
S18	**Jingpu Wu - JHU**	Python	dynamic	x,y,z	expm	`github.com/JimmyWu990110/BMsim_challenge_Jingpu`
S19	**Pedram Parnianpour - UBC**	Python	dynamic	x,y,z	expm	`github.com/Pedram-Parnianpour/bmsim-challenge-solver.git`
S20	**Swee Qi Pan - UTAR**	Matlab	dynamic	x,y,z	expm	`github.com/sweeqipan/BMsim_SQPan_YKTee`

2.5 Magnetization transfer pool formulation

In Bloch–McConnell simulations of magnetization transfer (MT), the semisolid pool is often modeled using only longitudinal (Z) magnetization, assuming that transverse components decay instantaneously due to extremely short T_2 values. This yields a simplified scalar description in which RF irradiation affects the pool through saturation governed by an effective absorption lineshape and is also known as the Henkelmann model [21]. We refer to this formulation as z-only MT. In contrast, a full M_{xyz} formulation explicitly includes transverse components, enabling the modeling of coherent effects such as off-resonance excitation and phase evolution. The z-only MT approach is computationally efficient and frequently used in simulations for MT and also CEST experiments. In the original formulation of this study, the way to simulate the MT pool was not explicitly stated, but results of the first round showed significant differences between M_{xyz} and z-only MT models for the same parameters. An example of this is shown in Section 3.2. These differences prompted a switch in the study to explicitly require M_{xyz} simulations. The z-only MT simulations are kept in the data source, but are not shown for all results. All submitted z-only MT entries reported using a Lorentzian lineshape for the MT pool.

2.6 Data analysis

Magnetization transfer ratio asymmetry (MTR_{asym}) was computed from each Z-spectrum as $Z(-\Delta\omega) - Z(+\Delta\omega)$ for positive offsets $\Delta\omega$. Inter-submission spread was quantified point-wise at each offset as the absolute difference between the maximum and minimum value across included submissions, separately for the spectra and MTR_{asym}; for each case we report the maximum and median spread across offsets. The normalization measurement at −300 ppm was excluded from Z-spectrum spread calculations. To place the final spreads in context, the same point-wise Z metric was also evaluated on archived revisions of the shared results spreadsheet, restricted to labeled submissions and excluding half-filled spectra and z-only MT columns. For each case we report the largest maximum Z spread observed during the study, the median spread at that stage, and the corresponding values of the latest submission. In the main comparison figures, spectra and asymmetry curves are also shown as differences to submission S02 (BMCTool), which was used as a documented Pulseq-based reference implementation. It generally lies close to the main cluster of submissions in terms of the point-wise Z-spectrum spread, although it is not the median submission for every case.

The main cluster was identified automatically for each case by ranking submissions by their root-mean-square Z difference to S02 and splitting at the largest relative gap in this ranking. If the remaining group still exceeded a maximum point-wise Z spread of 0.01, the split was repeated. Any submission within 10^{-3} of S02 at all offsets was always retained, as to avoid excluding submissions that were very close to the reference for cases with small overall spread.

2.7 Numerical implementation checks

Two additional analyses were performed to illustrate how numerical settings can affect CEST simulations. For the BART submission (S14), single- and double-precision variants were compared with the reference S02 in Case 1 and Case 2. For the operator-splitting MATLAB submission S03, Case 7 Z-spectra were calculated using multiple fixed timesteps from 500 to 5 μs (Supporting Information Section 4).

3 Results

The results of the study were collected online at: `https://docs.google.com/spreadsheets/d/1JN7VN-f1ktDrJgokb0FlUFwkHOMWYlPA_jSfnQoFOVc/` and are archived here: `https://doi.org/10.5281/zenodo.21265597`. A simple evaluation script is available at `https://colab.research.google.com/drive/1csiIjK-fiftdb7OwvJ84gWuv8lLADgv7`.

3.1 Round 1 results

All results for Cases 1 and 2 are shown in Figure 2. Case 1 showed the closest agreement among the Round 1 scenarios. After excluding the normalization offset, the maximum spread across the submitted Z-spectra was 1.4×10^{-3}, while the median point-wise Z spread was 4.0×10^{-5}. The corresponding maximum and median MTR_{asym} spreads were 9.3×10^{-5} and 1.3×10^{-5}, respectively.

For Case 2, the shorter saturation time increased the sensitivity to implementation details. Most submissions followed the same overall Z-spectrum shape, but one or more traces deviated visibly over parts of the offset range, resulting in a maximum point-wise Z spread of 0.44 and a median Z spread of 3.9×10^{-3}. The maximum and median MTR_{asym} spreads were 0.0014 and 1.0×10^{-4}, respectively.

All results for Cases 3 and 4 are shown in Figure 3. Introducing the 5-pool white-matter model in Case 3 led to broader, but more consistent deviations around the offsets affected by the CEST and MT pools. The maximum and median Z spreads were 0.0026 and 0.0023, and the

maximum and median MTR_{asym} spreads were 3.2×10^{-4} and 1.3×10^{-4}, respectively. Case 4, which used a short WASABI-like block pulse on the same pool model, showed visible differences in the Z-spectra, with maximum and median Z spreads of 1.1×10^{-3} and 6.8×10^{-4}, and maximum and median MTR_{asym} spreads of 0.0015 and 7.3×10^{-4}, respectively.

3.2 Magnetization transfer pool formulation results

Selected entries are shown for Case 3 to highlight the difference between full M_{xyz} MT simulations and z-only MT simulations in Figure 4. The z-only MT simulations formed a distinct response compared with the full M_{xyz} MT implementations, especially in spectral regions where the semisolid pool contributes strongly. Previous analysis [22] showed that the difference between z-only MT and M_{xyz} MT simulations can lead to significant differences in fitted pool fractions and exchange rates, which can be as high as 30% for the semisolid pool fraction or 50% for the exchange rate.

Because this difference reflects a modeling choice rather than only a numerical solver difference, z-only MT entries were excluded from the main comparison figures.

3.3 Round 2 results

All results for Cases 5 and 6 are shown in Figure 5. Case 5, the single Gaussian pulse, agreed closely, with a maximum Z spread of 3.8×10^{-3} and a median Z spread of 6.7×10^{-5}. The MTR_{asym} spread was much smaller, with maximum and median values of 1.0×10^{-4} and 5.7×10^{-5}, respectively, because the largest spectrum deviations were largely symmetric. A comparison of nearest-neighbor, linear, and cubic interpolation of the sampled Gaussian in S03 produced Z differences below 3×10^{-4} relative to Pulseq-CEST, smaller than this inter-submission spread (Supporting Information Section 3).

In Case 6, where the same Gaussian pulse was repeated as a pulse train, the largest absolute Z-spectrum spread was observed. The maximum and median Z spreads were 0.69 and 3.5×10^{-5}, and the maximum and median MTR_{asym} spreads were 0.0073 and 1.7×10^{-5}, respectively. The repeated pulses produced additional sideband features in the Z-spectra, which were reproduced with varying accuracy across submissions, hinting at differences in pulse train implementation, interpulse delay and frequency handling, and phase evolution during the delays. A possible implementation error is that ω_{RF} is not reset during t_{delay} (Supporting Information Section 2).

All results for Cases 7 and 8 are shown in Figure 6. Case 7 combined the Gaussian pulse train with the 5-pool model and showed maximum and median Z spreads of 0.012 and 2.9×10^{-3}, and maximum and median MTR_{asym} spreads of 3.5×10^{-4} and 1.5×10^{-4}, respectively. Compared with Case 6, the multi-pool model shows much closer agreement.

Case 8 showed a different pattern. The short two-pulse WASABI-like preparation produced visible Z-spectrum differences, with maximum and median Z spreads of 0.079 and 0.033. The MTR_{asym} curves showed maximum and median spreads of 6.4×10^{-4} and 1.7×10^{-4}, respectively.

3.4 Numerical precision and time-step convergence

Supporting Information Figure 4 summarizes two numerical implementation checks. The BART comparison shows that single- and double-precision arithmetic can produce visibly different spectra, even when the underlying solver and pool model are identical. The clearest example was Case 2 (2 s CW saturation on the 2-pool model), where single-precision BART showed high-frequency oscillatory noise in the difference plots. The maximum point-wise deviation from S02 was approximately 1.1×10^{-3} in Z for single precision and 2.6×10^{-4} for double precision, corresponding to an approximately fourfold reduction in peak numerical error. The mean point-wise deviation was reduced from 5.6×10^{-4} to 3.2×10^{-5}. Case 1 (15 s CW saturation) showed the same pattern at smaller amplitude.

The fixed-timestep analysis for submission S03 in Case 7 shows a related effect. A timestep of 500 μs produced a clearly different Z-spectrum, reducing the timestep to 100 μs produces a normal looking spectrum, which still deviates from the finest timestep of 5 μs by over 10%. Halving the timestep to 50 μs reduces the deviation to less than 5% and further reducing the timestep to 10 μs produces a spectrum that is indistinguishable (at the shown scale) from the finest timestep.

3.5 Largest observed versus final inter-submission spread

Archived spreadsheet revisions show that the final spreads are much smaller than the largest values observed during the study for most cases (Figure 7). Cases 3 and 5 are shown as examples: at the stage of the largest observed Case 5 spread the maximum and median point-wise Z spreads were 1.45 and 6.3×10^{-3}, compared with the final 3.8×10^{-3} and 6.7×10^{-5}.

4 Discussion

In this study we showed that a main cluster of agreeing simulations could be found for each case, consisting of 16 entries in the CW Case 2, and 9 entries in Cases 6 and 8, which tentatively allows us to formulate a consensus on simulations. However, this study also shows that Bloch-McConnell simulations are not automatically interchangeable across independent implementations, even when the nominal pool model, RF preparation pulse trains, offsets, and initialization conditions are specified. The degree of agreement depended strongly on the simulation case. The long CW 2-pool case showed close agreement, whereas shorter saturation, shaped pulses, pulse trains, and the 5-pool white-matter model all increased the observed variability. These findings indicate that benchmark cases need to include both simple reference scenarios and more clinically realistic preparations; agreement in a steady-state-like 2-pool case alone is not sufficient to validate a simulation framework for quantitative CEST applications.

A major limitation of this study is the absence of an independent ground truth. The benchmark compares independently developed implementations on the same specified Bloch-McConnell problem. Agreement of a main cluster therefore indicates consistency among frameworks, but does not prove correctness of the underlying solution. Analytical closed-form solutions exist only for highly restricted cases or with several simplifying assumptions, and experimental spectra cannot serve as a unique reference because they include acquisition imperfections and possible model mismatch. This work therefore does not validate Bloch-McConnell simulations directly, but provides a community comparison and a practical baseline against which new implementations can be checked.

The first round separated several sources of variability. Case 1 was intended as the simplest baseline and showed the smallest spread, suggesting that most implementations handle the basic Bloch-McConnell evolution, offset convention, and relaxation/exchange terms similarly under near steady-state conditions. The larger spread in Case 2 suggests that transient behavior is more sensitive to numerical and implementation choices. Since the pool model and RF amplitude were unchanged between Cases 1 and 2, these differences point to details such as matrix exponentiation, ODE integration tolerances, pulse/discretization conventions for CW irradiation, or the precise handling of delays. The lower spread in Case 1 may reflect that the long saturation approaches a common steady-state solution, whereas the shorter preparation in Case 2 preserves differences in transient evolution.

The 5-pool model introduced a second major source of variability. The comparison of full M_{xyz} and z-only MT implementations showed that these approaches produce systematically different spectra and should not be mixed without explicit labeling. This is expected because a z-only MT model approximates the MT pool using longitudinal saturation and an absorption lineshape, whereas a full M_{xyz} implementation evolves transverse components explicitly. Both

approaches may be useful in specific contexts, but they are not equivalent for the benchmark cases used here. For this reason, z-only MT results were retained in the data source but excluded from the main figures. The lineshape of the ssMT effect is an open question in the field, with Lorentzian, super Lorentzian, Gaussian, and other lineshapes being used in different studies [23]. Depending on the investigated tissue and the specific application, different lineshapes may be more appropriate. The present benchmark does not address this question, but it highlights that the choice of MT modeling can have a substantial impact on the simulated spectra and should be clearly specified in future studies. Furthermore, these lineshapes are often implemented using numerical approximations, such as discretization of the lineshape or interpolation around pole locations. This can introduce additional variability, and possibly errors if not implemented carefully.

The second round showed that shaped RF preparations are another important source of disagreement. The Gaussian pulse train (Case 6) produced the largest absolute Z-spectrum spread, even though the pulse shape was provided in a machine-readable file. This suggests that providing the shape alone does not fully eliminate ambiguity: interpolation of the RF waveform, time discretization, conversion between peak and RMS B_1, treatment of phase, and the specific pulse integration can all affect the result. Among standard interpolation methods the effect is small: nearest-neighbor, linear, and cubic resampling of the Case 5 Gaussian in one solver differed from Pulseq-CEST by less than 3×10^{-4} in Z (see Supporting Information Section 3). The pulse-train cases further amplify these choices because small differences are repeated many times and multiple pulses necessitate accurate handling of phase evolution and relaxation during interpulse delays. In particular, leaving a static RF frequency offset in the system matrix during t_{delay} is a possible implementation error that can separate submissions into distinct clusters (see Supporting Information Section 2 for more details). Ambiguities in the challenge definition of the pulse shape were also reported by several participants, including the exact B_1 scaling, the time discretization of the pulse, and the handling of phase in the pulse train case although we hope that the updated challenge definition in the second round clarified these points.

The multi-pool model in Case 7 shows much closer agreement, indicating that the differences are not caused by the multi-pool model itself, but by the shaped pulse implementation and the long T_1 and T_2 values of water for the 2-pool model. The long relaxation times of water in the 2-pool model lead to a slow decay of the magnetization, which necessitates accurate simulation throughout the pulse train, as any small difference in the magnetization evolution can accumulate and lead to significant differences in the final spectrum.

4.1 Solver choices and differences

In general, the results show that the same solution can be reached with different numerical strategies, provided that the implementations are sufficiently accurate for the RF preparation and pool model. Closely agreeing submissions used different approaches, including matrix exponentiation, Padé approximations, eigenvalue decompositions, ODE solvers, operator splitting, and state-transition matrices that combine aspects of matrix-exponential and ODE propagation. Thus, the solver class itself does not appear to determine the result uniquely. Instead, deviations are more likely caused by the details of how the solver is applied, such as time-step selection, waveform discretization, convergence criteria, and the handling of very short relaxation constants.

One important distinction is whether the simulation uses fixed or adaptive time steps. A fixed time-step implementation can be efficient and reproducible, but the chosen step size must be short enough to resolve the fastest dynamics in the system. This becomes particularly relevant for shaped pulses and for full M_{xyz} MT implementations, where very short T_2 values lead to rapidly decaying transverse components. If these dynamics are under-resolved, the resulting spectrum can deviate even when the pool parameters are nominally identical. Conversely, very small fixed time steps may increase memory use and computation time, especially for long pulse

trains and many offsets.

Adaptive ODE solvers and matrix-exponential approaches avoid some of these limitations, but introduce their own implementation choices. ODE solvers depend on tolerances, step sizes, and event discretization. Matrix-exponential approaches depend on how the RF waveform is divided into constant-amplitude intervals and how the exponential is evaluated, for example by Padé approximation or eigenvalue decomposition. For well-conditioned problems and carefully implemented solver choices these approaches should converge to the same solution, but differences can appear for long transient solutions with slow decays, when the RF pulse is coarsely sampled, when the ODE becomes stiff due to very short decay times, or approximations are used without convergence checks.

Supporting Information Figure 4 illustrates the impact of floating point precision for the BART implementation. In Case 2 (CW transient saturation), single-precision arithmetic introduced visible high-frequency noise in the difference spectra, whereas the double-precision variant reduced the peak numerical error by approximately a factor of four and the mean point-wise error by more than an order of magnitude. This indicates that floating-point precision can affect even CW CEST simulations when transient evolution is resolved over many integration steps. While only shown here for BART, this is expected to affect all solver classes. Nevertheless, the single-precision result still were precise enough for practical applications in all examples.

The same figure shows a related effect in Case 7 for fixed-step operator-splitting timescale in S03: an overly coarse timestep of 500 μs produced a clearly incorrect spectrum, while progressively smaller timesteps converged toward the same solution. These findings support the view that solver accuracy must be validated for the specific RF preparation and exact numerical settings are important to achieve reproducible results.

4.2 Implications for qMRI and future Bloch-McConnell simulations

The relationship between Z-spectrum spread and MTR_{asym} spread was case dependent. In some cases, large Z-spectrum differences produced comparatively small MTR_{asym} differences, consistent with approximately symmetric deviations. In other cases, especially when the 5-pool model was combined with pulsed saturation, differences became strongly asymmetric and led to larger MTR_{asym} variability. This is important for qCEST because many downstream analyses use asymmetry or fitted pool parameters rather than the raw Z-spectrum. A deviation that appears modest in Z can therefore still be relevant if it affects the spectral asymmetry near the fitted pool resonance.

To place this variability in a clinical context, we compare it with APTw imaging, which reports MTR_{asym} at 3.5 ppm after pulsed saturation with $B_{1,\mathrm{rms}} = 2\,\mu\mathrm{T}$ and $T_{\mathrm{sat}} = 2$ s [4], which is very close to the saturation used in Case 7. In Case 7 the MTR_{asym} peak of the main cluster is 0.29% (2.9×10^{-3}), while the inter-submission MTR_{asym} spread is 0.035% (3.5×10^{-4}; median 0.015%). The Case 7 pool model represents healthy white matter rather than tumor, so the absolute APTw amplitude is not expected to match the glioblastoma values. In IDH-wildtype glioblastoma, the median APTw signal in contrast-enhancing lesions was 2.23% in progressive disease versus 1.76% in treatment-related effects, a difference of 0.47 percentage points; within six months after radiotherapy the medians were 2.7% versus 1.26% (1.4 percentage points). The maximum Case 7 inter-submission MTR_{asym} spread observed during the study was 0.09% (9.2×10^{-4}; median 0.067%), about a fifth of the 0.47 percentage-point APTw contrast and about a third of the simulated peak. The current spread of 0.035%, after debugging simulations and revised submissions, is about an eighth of that peak and about an order of magnitude below the 0.47 percentage-point contrast.

The current Case 7 spread is therefore smaller than those clinical contrasts, whereas the maximum spread during the study was a substantial fraction of the smaller APTw contrast. Other implementation choices can still be large enough to affect quantitative interpretation when the expected contrast is small. Studies that rely on Bloch-McConnell simulations for quantitative

CEST analysis should therefore be aware of the potential for implementation-dependent variability and the effect that this can have on the interpretation of results. For example, if a study uses a Bloch-McConnell simulation to fit pool fraction or exchange rate, the choice of z-only versus full M_{xyz} MT implementation can lead to systematic differences in the fitted parameters in the order of 30% as shown by Schuenke et al. [22].

The present comparison does not yet assign all deviations to a single cause. Several factors can interact, including RF definitions, MT-pool formulation, solver choice, implementation and accuracy, and waveform discretization. In addition, some differences arose from ambiguity in the original challenge description rather than from an error in a particular simulation package. The benchmark therefore serves two purposes: it identifies cases where implementations diverge, and it tested which parts of the simulation specification need to be stated for reproducible quantitative CEST.

As a reference simulation we suggest the BMCTool results (S02) [10], which showed no significant deviation from the main cluster of results. The tool is easy to use with its Python interface and support for pulseq files. Therefore, comparing against BMCTool provides a quick and easy check for reasonable results, with small deviations being possible due to the above-mentioned implementation details.

4.3 Take-home recommendations and reporting guidelines

The challenge shows that some implementation choices are critical, whereas others are less consequential if they are applied carefully. Critical choices include the MT formulation (full M_{xyz} versus z-only MT), the definition and discretization of the RF waveform (peak versus RMS B_1, interpolation, and phase), the handling of interpulse delays, and whether magnetization is reset between offsets. The solver class itself appears less consequential: matrix exponentials, ODE integrators, and operator splitting all reached very close results in the main cluster when time steps or tolerances were adequate. Floating-point precision can introduce visible noise in long transient simulations, but the single-precision examples remained close enough for practical purposes.

When developing a new Bloch-McConnell simulation implementation, we recommend the following steps before using it for quantitative CEST analysis: First, compare Case 1 against the main cluster of results to test offset convention, sign, relaxation, and exchange under near steady-state CW conditions. Second, Case 2 tests transient CW evolution, where differences due to time-step, tolerance, and floating-point choices may become visible. Third, if an MT pool is used, Case 3 should be compared, if the implementation supports full M_{xyz} MT. If only z-only MT is implemented, the results should be clearly labeled as such and compared with other z-only MT implementations. Fourth, if the intended application uses shaped pulses or pulse trains, Cases 5–7 should be run to check for the correct implementation of the RF waveform, interpulse delays, and phase evolution.

The main clusters of agreeing submissions (Figures 2–6) contained 18 of 19, 16 of 20, 18 of 19, 18 of 19, 13 of 16, 9 of 17, 14 of 17, and 9 of 17 entries for Cases 1–8, with maximum pointwise deviations within the cluster ($\Delta Z_{\mathrm{cluster}}$) of 9.2×10^{-4}, 2.7×10^{-4}, 5.9×10^{-4}, 9.5×10^{-4}, 2.0×10^{-4}, 9.8×10^{-3}, 5.8×10^{-3}, and 2.0×10^{-4}, respectively. These cluster ranges can be used as a benchmark: a new implementation that remains within the corresponding cluster range for the relevant cases, i.e. if comparing against S02, is consistent with the community result reported here. They indicate the level of agreement achieved by the community and should not be read as an acceptable-error threshold, a validated error bound, or proof of correctness.

To make comparisons and reproducibility easier, future Bloch-McConnell simulation studies should report at least the following:

- Pool model: relative pool sizes f, T_1, T_2, exchange rates k, and chemical shifts, ideally as

a machine-readable file.

- MT pool: full M_{xyz} or z-only MT, and the absorption lineshape if z-only MT is used.
- RF and sequence: B_1 definition (peak or RMS), waveform sampling or interpolation, pulse phase, interpulse delays, whether the RF frequency offset is reset during delays, spoiler delay, and gyromagnetic ratio.
- Initialization: whether $Z_i = 1$ is reset before each offset, or a recovery delay is simulated.
- Numerical solver: solver class, time-step strategy or tolerances, floating-point precision, and any convergence checks.
- Furthermore authors could reference this work and note if they compared against the main-cluster ranges reported here.

5 Conclusion

This benchmark shows that independent Bloch-McConnell simulation frameworks can agree closely for simple reference cases, but may diverge substantially for transient saturation, shaped RF pulses, pulse trains, and multi-pool models with MT. Agreement improved strongly over the course of the study: for most cases the final maximum and median Z spreads are substantially smaller than the largest values observed earlier, by up to two to three orders of magnitude (e.g. Case 5 from a maximum of 1.45 to 3.8×10^{-3}; Figure 7). Clear specification of MT modeling, RF power definitions, pulse discretization, and initialization conditions is therefore essential for reproducible quantitative CEST. For newly implemented simulations we suggest comparing against the BMCTool results (S02) and against the cluster ranges and reporting guidelines in Section 4.3. The provided open challenge data and analysis scripts provide a basis for continued community comparison and for developing more standardized CEST simulation practice.

Author contributions

M.Z. and P.S. designed the study. M.Z., and P.S. defined the simulation cases and specifications. M.Z., P.S. and M.H coordinated the study and collected results. M.H. performed the analysis and wrote the initial paper draft. All named authors submitted their results and contributed to writing, editing and reviewing of the manuscript.

Acknowledgements

The authors thank all participating groups for contributing simulation results to the benchmark. AI-assisted coding tools were used to develop and refine the Python analysis scripts for parsing submitted results, computing summary metrics, and generating manuscript figures. AI-assisted writing tools were also used to support drafting and editing of the manuscript text. All authors reviewed, edited, and approved the final content and take responsibility for the published work. This research was funded in whole or in part by the Austrian Science Fund (FWF) 10.55776/F100800 and 10.55776/I4870. This work was partially supported by Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) grants 372486779 (SFB1340) and ZA 814/15-1.

References

[1] Zhou Jinyuan, Zaiss Moritz, Knutsson Linda, et al. Review and consensus recommendations on clinical APT-weighted imaging approaches at 3T: Application to brain tumors. *Magnetic Resonance in Medicine.* 2022;88:546-574.

[2] Chen Lin, Wei Zhiliang, Chan Kannie W.Y., et al. Protein aggregation linked to Alzheimer's disease revealed by saturation transfer MRI. *NeuroImage.* 2019;188:380-390.

[3] Tietze Anna, Blicher Jakob, Mikkelsen Irene Klærke, et al. Assessment of ischemic penumbra in patients with hyperacute stroke using amide proton transfer (APT) chemical exchange saturation transfer (CEST) MRI. *NMR in Biomedicine.* 2014;27:163-174.

[4] Zeyen Thomas, Krause Inga, Decker Andreas, et al. Amide proton transfer-weighted (APTw) CEST MRI in clinical routine for single time point diagnosis of pseudoprogression in IDH-wildtype glioblastoma. *Neuro-oncology.* 2026;28(3):790.

[5] Kroh Florian, Knebel Doeberitz Nikolaus, Breitling Johannes, et al. Semi-solid MT and APTw CEST-MRI predict clinical outcome of patients with glioma early after radiotherapy. *Magnetic Resonance in Medicine.* 2023;90:1569-1581.

[6] Paech Daniel, Windschuh Johannes, Oberhollenzer Johanna, et al. Assessing the predictability of IDH mutation and MGMT methylation status in glioma patients using relaxation-compensated multipool CEST MRI at 7.0 T. *Neuro-Oncology.* 2018;20:1661-1671.

[7] McConnell Harden M.. Reaction Rates by Nuclear Magnetic Resonance. *The Journal of Chemical Physics.* 1958;28:430-431.

[8] Zhou Jinyuan, Payen Jean Francois, Wilson David A., Traystman Richard J., Zijl Peter C.M. Van. Using the amide proton signals of intracellular proteins and peptides to detect pH effects in MRI. *Nature Medicine 2003 9:8.* 2003;9:1085-1090.

[9] Herz Kai, Mueller Sebastian, Perlman Or, et al. Pulseq-CEST: Towards multi-site multi-vendor compatibility and reproducibility of CEST experiments using an open-source sequence standard. *Magnetic Resonance in Medicine.* 2021;86:1845-1858.

[10] Schuenke Patrick. *BMCTool: Python Bloch-McConnell (BMC) simulation tool.* 2026.

[11] Zhang Huabin, Qiu Bensheng, Xu Jiadi, Chan Kannie W.Y., Huang Jianpan. CESTsimu: An open-source GUI for spectral and spatial CEST simulation. *Magnetic Resonance in Medicine.* 2025;93:2595-2609.

[12] Goerke Steffen, Zaiss Moritz, Bachert Peter. Characterization of creatine guanidinium proton exchange by water-exchange (WEX) spectroscopy for absolute-pH CEST imaging in vitro. *NMR in Biomedicine.* 2014;27(5):507–518.

[13] Zijl Peter C. M., Lam Wilfred W., Xu Jiadi, Knutsson Linda, Stanisz Greg J.. Magnetization Transfer Contrast and Chemical Exchange Saturation Transfer MRI. Features and analysis of the field-dependent saturation spectrum. *NeuroImage.* 2018;168:222–241.

[14] Schuenke Patrick, Windschuh Johannes, Roeloffs Volkert, Ladd Mark E., Bachert Peter, Zaiss Moritz. Simultaneous mapping of water shift and B1(WASABI)—Application to field-Inhomogeneity correction of CEST MRI data. *Magnetic Resonance in Medicine.* 2017;77:571-580.

[15] Graf Christina, Rund Armin, Aigner Christoph Stefan, Stollberger Rudolf. Accuracy and performance analysis for Bloch and Bloch-McConnell simulation methods. *Journal of Magnetic Resonance.* 2021;329:107011.

[16] Vladimirov Nikita, Cohen Ouri, Heo Hye Young, Zaiss Moritz, Farrar Christian T., Perlman Or. Quantitative molecular imaging using deep magnetic resonance fingerprinting. *Nature Protocols 2025 20:10.* 2025;20:3024-3054.

[17] Wang Kexin, Huang Jianpan, Ju Licheng, et al. Creatine mapping of the brain at 3T by CEST MRI. *Magnetic Resonance in Medicine.* 2024;91:51-60.

[18] Zaiss Moritz, Angelovski Goran, Demetriou Eleni, McMahon Michael T., Golay Xavier, Scheffler Klaus. QUESP and QUEST revisited – fast and accurate quantitative CEST experiments. *Magnetic Resonance in Medicine.* 2018;79:1708-1721.

[19] Singh Munendra, Jiang Shanshan, Li Yuguo, Zijl Peter, Zhou Jinyuan, Heo Hye Young. Bloch simulator–driven deep recurrent neural network for magnetization transfer contrast MR fingerprinting and CEST imaging. *Magnetic Resonance in Medicine.* 2023;90:1518-1536.

[20] Scholand Nick, Wang Xiaoqing, Roeloffs Volkert, Rosenzweig Sebastian, Uecker Martin. Quantitative MRI by nonlinear inversion of the Bloch equations. *Magnetic Resonance in Medicine.* 2023;90:520-538.

[21] Graham S. J., Henkelman R. Mark. Understanding pulsed magnetization transfer. *Journal of Magnetic Resonance Imaging.* 1997;7:903-912.

[22] Schuenke Patrick, Herz Kai, Zu Zhongliang, et al. Validate your CEST simulation!. In: 0907; 2023.

[23] Assländer Jakob, Gultekin Cem, Flassbeck Sebastian, Glaser Steffen J., Sodickson Daniel K.. Generalized Bloch model: A theory for pulsed magnetization transfer. *Magnetic Resonance in Medicine.* 2022;87:2003-2017.




Figure 1: Overview of all eight cases, one row per case. Left: saturation scheme. Cases 1 to 4 use CW saturation with different saturation times. Cases 5, 6 and 7 use one or multiple Gaussian pulses and Case 8 repeats the CW saturation of Case 4 with a short interpulse delay. The gray area marks the 6.5 ms spoiler delay after saturation, which is visible for short saturation cases. Middle and right: exemplary Z-spectra and MTR_{asym}, simulated with the MATLAB Pulseq-CEST implementation (equivalent to submission S01).




Figure 2: Cases 1 and 2: Z- and MTR_{asym} spectra with differences to S02. The main cluster is marked in the Z-spectra plots and contained 18 (of 19) entries in Case 1 and 16 (of 20) entries in Case 2, with maximum deviations of $9.24 \cdot 10^{-04}$ and $2.71 \cdot 10^{-04}$, respectively within those clusters. S17 shows the largest deviation in Case 1. S14b represents a solver variant of S14, using double precision instead of single precision. In Case 2, S14, S15, S15b, and S17 fall outside the main cluster; S15b is an alternative solver variant of S15 and shows the largest deviation. More information on these solver variants can be found in Supporting Information Section 1.




Figure 3: Cases 3 and 4: simulated Z- and MTR_{asym} spectra with differences to S02. The main cluster is marked in the Z-spectra plots and contained 18 (of 19) entries in Case 3 and 18 (of 19) entries in Case 4, with maximum deviations of $5.90 \cdot 10^{-04}$ and $9.55 \cdot 10^{-04}$, respectively within those clusters. S17 shows the largest deviation in Case 3. In Case 4, S18 falls outside the main cluster and also differs noticeably in MTR_{asym}.




Figure 4: Selected entries for Case 3 showing full M_{xyz} MT simulations (S01, S06, S10 and S19) in comparison with z-only MT simulations (S01, S10 and S13; labeled zMT in the figure). The M_{xyz} entries closely match with the chosen reference (S01, labeled xyzMT). The z-only MT entries show a significant difference to the reference, with close agreement between each other.




Figure 5: Cases 5 and 6: simulated Z- and MTR_{asym} spectra with differences to S02. The main cluster is marked in the Z-spectra plots and contained 13 (of 16) entries in Case 5 and 9 (of 17) entries in Case 6, with maximum deviations of $2.02 \cdot 10^{-04}$ and $9.85 \cdot 10^{-03}$, respectively within those clusters. S08, S17, and S18 fall outside the Case 5 main cluster.




Figure 6: Cases 7 and 8: simulated Z- and MTR_{asym} spectra with differences to S02. The main cluster is marked in the Z-spectra plots and contained 14 (of 17) entries in Case 7 and 9 (of 17) entries in Case 8, with maximum deviations of $5.85 \cdot 10^{-03}$ and $1.95 \cdot 10^{-04}$, respectively within those clusters. In Case 7, S08, S17, and S18 deviate most strongly; Case 8 shows broader spread with several separated traces.




Figure 7: Top: Case 3 Z and MTR$_{\text{asym}}$ (3 to 0 ppm) beside Case 5 Z, with largest observed deviation (top row, red traces; Case 3 includes z-only MT, dotted) versus current (bottom row, green traces; z-only MT omitted). Bottom: for each case, the largest maximum point-wise Z spread observed during the study and the median spread at that same stage, versus the final maximum and median, shown on a log scale.