
Introduction to Computational Chemistry

Part 7: Chemical concepts

Eno Paenurk & Patrick Finkelstein & Felix Pultar



07.06.2023

Webinar overview



- Part 1: Takeoff
- Part 2: Reaction pathways from scratch
- Part 3: Preparing structures
- Part 4: Density Functional Theory
- Part 5: Organometallics
- Part 6: Spectroscopy
- Part 7: Chemical concepts

Goals accomplished so far



- Obtaining reaction profiles with autodE
- Obtaining good starting structures
 - from SMILES
 - from a guess
 - refining with xtb
- Running DFT jobs with ORCA
 - Minima intermediates
 - Saddle-points transition states
 - Transition metals
 - Spectroscopy: IR, NMR, UV/Vis

- Reading relevant data from the output
 - Energies
 - Thermochemistry
- Visualizing the results
 - 3D structures/trajectories
 - Vibrations

Goals today



- Extract information about the HOMO/LUMO
- Visualize molecular orbitals
- Calculate and visualize orbital interactions
- Calculate partial charges
- Visualize electrostatic potential maps

Orbitals



Remember that orbitals are not observables, but:

"Whatever makes you inspired to do new experiments, it's fine."

-Frank Neese

Orbital occupations and energies are listed after the SCF calculation:

```
ORBITAL ENERGIES
```

UNDITAL ENERGIES

. . .

NO	OCC	E(Eh)	E(eV)	
 6 7	2.0000	-0.324924 -0.077083	-8.8416 -2.0975	




To visualize the orbitals, you can either:

- generate .cube files of orbitals with orca_plot (one .cube file per orbital)
- request extra printout to the output:
 - ! Printbasis PrintMOs (recommended only for single points)




The .cube or the output file can be opened by Chemcraft ...





... and Avogadro.





How big is the orbital?

You need to choose the contour value (or isovalue) when visualizing 3D functions like orbitals. Usually values between 0.02 and 0.06 (e/Bohr 3) give reasonable results.

There is more than one flavor

We show how to plot "canonical" molecular orbitals. There are various transformation schemes that can produce "more intuitive" orbitals.

Orbital interactions



If we want more insight into the orbital interactions between different parts of the molecule, we can perform energy decomposition analysis techniques.


In ORCA, we can use the ETS-NOCV* workflow:

- Run SP calculations on the two interacting fragments
- 2 Combine the wavefunctions of the fragments
- Run SP calculation of the combined fragments starting from the combined wavefunction
- Read the orbital interaction energies from the output
- Visualize the respective orbitals and the electron density flow (so-called deformation density)

Orbital interactions



The NOCV .cube files can be opened by Chemcraft to calculate the *deformation density*.



2nd strongest orbital interaction (30% of total interaction energy)

Red – electron density depletion Blue – electron density accumulation

Partial charges



There are a lot of methods for calculating partial charges

Partial charges in ORCA:

Method	Approach	How to invoke	Comment
Mulliken and Löwdin (LOEWDIN) population analyses	Dissect the wavefunction	printed by default	basis set dependent, not very reliable
ChEIPG (Charges from the Electrostatic Potential on a Grid)	Fit charges to electrostatic potential	! CHELPG	stable, but not al- ways great for large molecules
Hirshfeld population analysis	Dissect the electron density	! Hirshfeld	stable, but typically too low

Partial charges



Showcase: CO₂ (showing C)

	B3LYP/def2-		BP86/def2-	
Method	SVP	TZVP	SVP	TZVP
Mulliken	0.26	0.37	0.21	0.31
Löwdin	0.13	-0.41	0.09	-0.45
ChEIPG	0.74	0.71	0.69	0.66
Hirshfeld	0.36	0.35	0.33	0.32

- ChEIPG and Hirshfeld preferred over Mulliken and Löwdin
- Look into different options if good partial charges are crucial for your study

Note


Partial charges are not oxidation states. If you want to calculate oxidation states, talk to your local computationalist.



Electrostatic potential map



The electostatic potential (calculated from Mulliken charges) can be mapped onto the electron density or van der Waals surface in Avogadro.



Summary



There are a lot of ways to chemically interpret and analyze your calculations – we barely scratched the surface.

We covered:

- Orbital analyses
 - Visualization
 - Orbital interaction calculations
- Electrostatics
 - Partial charges
 - Electrostatic potential maps from partial charges

You may further find interesting:

- Formal oxidation states
- Spin density
- Energy decomposition analyses
- Localized orbitals
- Electrostatic potential maps from electron density

Live-Demo





Exercises





17 / 17