ORTHOPEDIC & JOINT RECONSTRUCTION HOSPITAL

POST-OPERATIVE DISCHARGE SUMMARY

PATIENT INFORMATION:

Patient Name: Nirmala Devi | Age: 66 | Gender: Female
UHID: OJR-ORTHO-2026-3391 | Surgery Date: 2026-10-01 | Discharge Date: 2026-10-06
Operative Surgeon: Dr. Vikramaditya Rao, MS (Ortho)

PROCEDURE PERFORMED:

Right Total Knee Arthroplasty (Cruciate-Retaining Tricompartmental Prosthesis with High-Flexion Polyethylene Insert) under Subarachno

PRE-OPERATIVE DIAGNOSIS:
Severe End-Stage Osteoarthritis of Right Knee (Kellgren-Lawrence Grade IV) with 15-degree varus deformity and severe functional mobi

POST-OPERATIVE HOSPITAL COURSE:
Surgical wound clean, intact with subcuticular Monocryl closure and negative pressure waterproof dressing. Drain removed on post-op Da

POST-OP INVESTIGATIONS:
- Post-op Hemoglobin: 10.2 g/dL (Stable, no transfusion required)
- Serum Electrolytes: Na 138, K 4.1 (Normal)
- Bilateral Lower Extremity Venous Doppler: Negative for Deep Vein Thrombosis (DVT)
- Post-op Check X-Ray Right Knee: Prosthesis in anatomical alignment, cement mantle intact.

DISCHARGE PRESCRIPTION:

1. Tab. Rivaroxaban 10 mg - 1 tablet orally once daily with food for 14 days (Deep Vein Thrombosis DVT Thromboprophylaxis).
2. Tab. Tramadol 37.5 mg + Paracetamol 325 mg (Ultracet) - 1 tablet orally twice daily after meals as needed for moderate pain.
3. Tab. Paracetamol 650 mg - 1 tablet orally every 8 hours as baseline pain relief.
4. Tab. Pantoprazole 40 mg - 1 tablet orally once daily morning before meals for 14 days.
5. Tab. Calcium Carbonate + Cholecalciferol - 1 tablet daily with lunch for 3 months.
6. Syp. Lactulose 15 mL - at bedtime as needed for opioid-induced bowel constipation.

CRITICAL SAFETY & MEDICATION ALERTS:

- DVT PROPHYLAXIS COMPLIANCE: Take Rivaroxaban daily without missing doses. Joint replacement carries elevated risk of post-op venous t
- Avoid additional NSAIDs (Diclofenac/Ibuprofen) due to bleeding risk when combined with Rivaroxaban.
- Do NOT wet surgical dressing. Keep dry until suture line review.

PHYSICAL THERAPY & HOME RECOVERY:

- Perform ankle pumps 20 times every hour, quadriceps sets, and straight leg raises 3 times daily.
- Use four-legged walker for support while walking. Avoid squatting or sitting cross-legged on the floor.
- Apply ice packs over anterior knee for 15 minutes after physiotherapy sessions.
- Red Flags: Sudden unilateral calf swelling/pain (DVT), chest pain/breathlessness (pulmonary embolism), or active redness/foul drainage from in

FOLLOW-UP:
Suture line inspection and dressing change in Orthopedic OPD on 2026-10-18.