

Are declines concentrated in particular regions, habitats or ecological groups?


Regions

Declines are widespread, a little steeper on the coast than inland

Median species change in share of squares where recorded, 1977–1990 → 2006–2020 (fair comparison)







– Typical species change: coast -12%, tablelands -10%, inland -1%.

– 41% of map cells are down 10% or more, scattered across the state rather than clustered.

– Continuing decliners fall hardest: coast 11, inland 9, tablelands 7 species.


Atlas surveys 1977–2020; same effort per grid square in each period (432 squares). Change is 1977–1990 to 2006–2020. Region names are not yet confirmed, so zones are used.

Are declines concentrated in particular regions, habitats or ecological groups?

Habitats and ecological groups

Declines are concentrated in small, insect-eating woodland birds

Median change vs typical species by group, 1977–1990 → 2006–2020, with 95% range (n = species)



– Clearly below typical: small birds (-18%), dome nesters (-17%), insect-eaters (-16%), summer migrants (-15%), woodland birds (-15%).

– Clearly above typical: fruit-eaters (+45%), omnivores (+16%), large birds (+11%).

– Groups differ by more than chance for body size, main diet, nest type, movement; habitat differences are suggestive only.

Atlas surveys 1977–2020; same effort per grid square in each period (432 squares). Change is 1977–1990 to 2006–2020. Traits are provisional (species_traits_provisional.csv) and should be checked against the team's trait table.