0.0.1 $\langle \Delta H_{SD} \rangle$ Derivation

The equation $H = -q(1-q)$ defines a constant of motion for the SD part of the game, where q is the fraction of players in the 4th strategy. Using the transition probabilities of the different process we can derive an expression for the expected change in H within the simplex.

Where $i, j, k, N-i-j-k$ are the players playing R, P, S, and the 4th strategy respectively.

$\Delta H = H(t+1) - H(t)$,
$\Delta H = -x_{t+1}(1-x_{t+1}) - (-x_t(1-x_t))$
$\Delta H = -x_{t+1}(1-x_{t+1}) + x_t(1-x_t)$
$\Delta H = x_t(1-x_t) - x_{t+1}(1-x_{t+1})$

$\langle \Delta H \rangle = \sum_{i,j,k} \Big(H_s - H_{s'} \Big) T^{s \to s'}$, s is a particular state in the simplex.

$$\begin{aligned}
\langle \Delta H_{SD} \rangle = \frac{6}{N^5} \sum_{i=1}^{N} \sum_{j=1}^{N-i} \sum_{k=1}^{N-i-j} \Big[& (N-i-j-k)(1-N+i+j+k)(T^{R+} + T^{P+} + T^{S+} + T^{+R} + T^{+P} + T^{+S}) \\
& - (N-i-j-k+1)(-N+i+j+k)T^{R+} \\
& - (N-i-j-k+1)(-N+i+j+k)T^{P+} \\
& - (N-i-j-k-1)(2-N+i+j+k)T^{+R} \\
& - (N-i-j-k-1)(2-N+i+j+k)T^{+P} \\
& - (N-i-j-k-1)(2-N+i+j+k)T^{+S} \Big]
\end{aligned} \tag{1}$$

The terms with transitions within the RPS simplex can be ignored as q would not change between these states, therefore the term $H_s - H_{s'} = 0$

$p = N-i-j-k$, the number of players playing the 4th strategy.

$$\begin{aligned}
\langle \Delta H_{SD} \rangle = \frac{6}{N^5} \sum_{i=1}^{N} \sum_{j=1}^{N-i} \sum_{k=1}^{N-i-j} \Big[& p(1-p)(T^{R+} + T^{P+} + T^{S+} + T^{+R} + T^{+P} + T^{+S}) \\
& - (p+1)(-p)(T^{R+} + T^{P+} + T^{S+}) \\
& - (p-1)(2-p)(T^{+R} + T^{+P} + T^{+S}) \Big]
\end{aligned}$$

$$\begin{aligned}
\langle \Delta H_{SD} \rangle &= \frac{6}{N^5} \sum_{i=1}^{N} \sum_{j=1}^{N-i} \sum_{k=1}^{N-i-j} \Big[T^{R+} \left(p(1-p) + p(p+1) \right) + T^{+R} \left(p(1-p) - (p-1)(2-p) \right) \\
&\quad + T^{P+} \left(p(1-p) + p(p+1) \right) + T^{+P} \left(p(1-p) - (p-1)(2-p) \right) \\
&\quad + T^{S+} \left(p(1-p) + p(p+1) \right) + T^{+S} \left(p(1-p) - (p-1)(2-p) \right) \Big] \\
&= \frac{6}{N^5} \sum_{i=1}^{N} \sum_{j=1}^{N-i} \sum_{k=1}^{N-i-j} \Big[2p \left(T^{R+} + T^{P+} + T^{S+} \right) + (p - p^2 - (3p - p^2 - 2))(T^{+R} + T^{+P} + T^{+S}) \Big] \\
&= \frac{6}{N^5} \sum_{i=1}^{N} \sum_{j=1}^{N-i} \sum_{k=1}^{N-i-j} \Big[2p \left(T^{R+} + T^{P+} + T^{S+} \right) + (2-2p)(T^{+R} + T^{+P} + T^{+S}) \Big]
\end{aligned}$$

$$\langle \Delta H_{SD} \rangle = \frac{12}{N^5} \sum_{i=1}^{N} \sum_{j=1}^{N-i} \sum_{k=1}^{N-i-j} \Big[p \left(T^{R+} + T^{P+} + T^{S+} \right) + (1-p)(T^{+R} + T^{+P} + T^{+S}) \Big]$$

$$\langle \Delta H_{SD} \rangle = \frac{12}{N^5} \sum_{i=1}^{N} \sum_{j=1}^{N-i} \sum_{k=1}^{N-i-j} p \Big[(T^{R+} - T^{+R}) + (T^{P+} - T^{+P}) + (T^{S+} - T^{+S}) \Big] + T^{+R} + T^{+P} + T^{+S} \tag{2}$$

The continuous limit, where $x = i/N$, $y = j/N$, $z = k/N$, and $q = p/N$, $p = Nq$ and $q = 1 - x - y - z$ leads to:

$$\begin{aligned} \langle \Delta H_{SD} \rangle = \frac{12}{N^2} \int_0^1 dx \int_0^{1-x} dy \int_0^{1-x-y} dz \Big[& Nq \left[(T^{R+} - T^{+R}) + (T^{P+} - T^{+P}) + (T^{S+} - T^{+S}) \right] \\ & + (T^{+R} + T^{+P} + T^{+S}) \Big] \end{aligned}$$

Finally,

$$\begin{aligned} \langle \Delta H_{SD} \rangle = & \frac{12}{N} \int_0^1 dx \int_0^{1-x} dy \int_0^{1-x-y} dz \Big[q \left[(T^{R+} - T^{+R}) + (T^{P+} - T^{+P}) + (T^{S+} - T^{+S}) \right] \Big] \\ & + \frac{12}{N^2} \int_0^1 dx \int_0^{1-x} dy \int_0^{1-x-y} dz \left[T^{+R} + T^{+P} + T^{+S} \right] \end{aligned} \tag{3}$$

This can then be solved numerically and the critical population values can be found where $\langle \Delta H_{SD} \rangle = 0$.

Critical N found matches nicely with the simulated versions. Numerical integration in python code ./augRps.py, shows change of sign as expected. Matches nicely with the approximated values for the Moran process. The specific expression for Moran process is very long. Computed numerically and solved with scipy.integrate (reference scipy) Maybe can plot the simulated critical population sizes against the analyical on the same graph.

0.0.2 Analytical solution for the Local update process

Considering the local update process, an exact analytical solution can be found for $\langle \Delta H_{SD} \rangle$.

$$\begin{bmatrix} 0 & c & 1 & \gamma \\ 1 & 0 & c & \gamma \\ c & 1 & 0 & \gamma \\ \beta & \beta & \beta & 0 \end{bmatrix} \tag{4}$$

To simplify these derivations, I have fixed the values of $a = 0, b = 1$, allowing still to adjust the dynamics of the RPS game via c and the SD game via γ, β as used in all the results.

Evaluating the terms of the type $T^{R+} - T^{+R}$ for the local update process:

$$\begin{aligned} T^{R+} - T^{+R} &= \frac{i}{N} \frac{N-i-j-k}{N} \left(\frac{1}{2} + \frac{w}{2} \frac{\pi_+ - \pi_R}{\Delta \pi_{\max}} \right) - \frac{i}{N} \frac{N-i-j-k}{N} \left(\frac{1}{2} + \frac{w}{2} \frac{\pi_R - \pi_+}{\Delta \pi_{\max}} \right) \\ &= \frac{1}{2} \frac{i}{N} \frac{N-i-j-k}{N} w \left(\frac{\pi_+ - \pi_R}{\Delta \pi_{\max}} - \frac{\pi_R - \pi_+}{\Delta \pi_{\max}} \right) \\ &= w \frac{i}{N} \frac{N-i-j-k}{N} \left(\frac{\pi_+ - \pi_R}{\Delta \pi_{\max}} \right) \end{aligned}$$

substituting the population frequencies:

$$T^{R+} - T^{+R} = wx(1-x-y-z) \frac{\pi_+ - \pi_R}{\Delta \pi_{\max}} \tag{5}$$

the difference for the other two strategies can simply be derived by swapping the corresponding payoffs in the above expression.

Now considering the second integral with terms $T^{+R} + T^{+P} + T^{+S}$:

$$T^{+R} + T^{+P} + T^{+S} = \frac{i}{N}\frac{N-i-j-k}{N}\left(\frac{1}{2} + \frac{w}{2}\frac{\pi_R - \pi_+}{\Delta\pi_{\max}}\right) \tag{6}$$

$$+ \frac{j}{N}\frac{N-i-j-k}{N}\left(\frac{1}{2} + \frac{w}{2}\frac{\pi_P - \pi_+}{\Delta\pi_{\max}}\right) \tag{7}$$

$$+ \frac{k}{N}\frac{N-i-j-k}{N}\left(\frac{1}{2} + \frac{w}{2}\frac{\pi_S - \pi_+}{\Delta\pi_{\max}}\right) \tag{8}$$

Again replacing with the frequencies:

$$T^{+R} + T^{+P} + T^{+S} = (1-x-y-z)\Bigg[x\left(\frac{1}{2} + \frac{w}{2}\frac{\pi_R - \pi_+}{\Delta\pi_{\max}}\right) \tag{9}$$

$$+ y\left(\frac{1}{2} + \frac{w}{2}\frac{\pi_P - \pi_+}{\Delta\pi_{\max}}\right) + z\left(\frac{1}{2} + \frac{w}{2}\frac{\pi_S - \pi_+}{\Delta\pi_{\max}}\right)\Bigg] \tag{10}$$

$$= \frac{1}{2}(1-x-y-z)\left[x + y + z + \frac{w}{\Delta\pi_{\max}}\Big[x(\pi_R - \pi_+) + y(\pi_P - \pi_+) + z(\pi_S - \pi_+)\Big]\right] \tag{11}$$

Substituting in the expanded expressions for payoff differences e.g. $\pi_R - \pi_+$:

$$T^{+R} + T^{+P} + T^{+S} = \frac{1}{2}(1-x-y-z)\Bigg[x + y + z + \frac{w}{\Delta\pi_{\max}} \tag{12}$$

$$\Big[x(z + cy + \gamma(1-x-yz) - \beta(x+y+z)) \tag{13}$$

$$+ y(x + cz + \gamma(1-x-yz) - \beta(x+y+z)) \tag{14}$$

$$+ z(y + cx + \gamma(1-x-yz) - \beta(x+y+z))\Big]\Bigg] \tag{15}$$

Finally substituting everything back into $\langle \Delta H_{SD} \rangle$:

$$\langle \Delta H_{SD} \rangle = \frac{12}{N} \int_0^1 dx \int_0^{1-x} dy \int_0^{1-x-y} dz (1-x-y-z) \Bigg[\tag{16}$$

$$\frac{w}{\Delta\pi_{\max}} x(1-x-y-z)\left(\beta(x+y+z) - \gamma(1-x-y-z) - z - cy\right) \tag{17}$$

$$\frac{w}{\Delta\pi_{\max}} y(1-x-y-z)\left(\beta(x+y+z) - \gamma(1-x-y-z) - x - cz\right) \tag{18}$$

$$\frac{w}{\Delta\pi_{\max}} z(1-x-y-z)\left(\beta(x+y+z) - \gamma(1-x-y-z) - y - cx\right) \Bigg] \tag{19}$$

$$+ \frac{6}{N^2} \int_0^1 dx \int_0^{1-x} dy \int_0^{1-x-y} dz (1-x-y-z) \Bigg[\tag{20}$$

$$x + y + z + \frac{w}{\Delta\pi_{\max}} \Bigg(x(z + cy + \gamma(1-x-yz) - \beta(x+y+z)) \tag{21}$$

$$+ y(x + cz + \gamma(1-x-yz) - \beta(x+y+z)) \tag{22}$$

$$+ z(y + cx + \gamma(1-x-yz) - \beta(x+y+z)) \Bigg) \Bigg] \tag{23}$$

Solving the integrals leads to the exact analytical expression for local update process $\langle \Delta H_{SD} \rangle$:

$$\langle \Delta H_{SD} \rangle_{LU} = \frac{42\Delta\pi_{\max} + 2w\gamma(7-6N) + (-1+4\beta-c)(-7+4N)w}{280N^2\Delta\pi_{\max}} \tag{24}$$

which is nicely expressed by the actual values in the payoff matrix for the game, here I have fixed the values of $a = 0, b = 1$ so that the expected drift can be written as an expression of c, β, γ, N and w. $\Delta\pi_{\max}$ is the maximum payoff difference which will depend on the choice of parameters, however it is a constant and usually defined as $b - c$ or $1 - c$ when b is fixed to 1 in this case as mostly we consider $0 < \gamma, \beta < 1$ so the maximal difference is between the winning and losing case of the rock paper scissors sub game.

Since the critical population size is defined as a change in sign in $\langle \Delta H_{SD} \rangle$, the critical population size, N_c can also be derived by setting the above equation to 0 and solvign for N.

This leads to:

$$N_c = \frac{-42\Delta\pi_{\max} + w(28\beta - 7c - 14\gamma - 7)}{w(16\beta - 4c - 12\gamma - 4)} \tag{25}$$

$$= \frac{-42\Delta\pi_{\max}}{w(16\beta - 4c - 12\gamma - 4)} + \frac{28\beta - 7c - 14\gamma - 7}{16\beta - 4c - 12\gamma - 4} \tag{26}$$

I have compared this value to the numerically solved solutions of the triple integral and the results match perfectly. For both the values of the integral and also the critical population size which was previously calculated using brentq root finding method. Due to increasing complexity in the Moran process, I have used numerical methods to compute the value of the integral and critical population size instead of a closed form solution.

See if its possible to also derive closed from solution for RPS and 4 strategy observation variables but this was a very long process for SD case so may be worth solving with numerical method. Definitely not for the Moran process.



Figure 1: Comparison of simulated critical population sizes against the derived expected change in H for Moran process. Small deviations due to stochasticity in the simulations. $\gamma = 0.2, \beta = 0.1$ standard RPS. Small deviations due to random nature of simulations, but close match to analytical result. Drift reversal is present in the agent based simulations and exactly matching analytical prediction for these paremeters.

0.0.3 $\langle \Delta H_{RPS} \rangle$ Derivation

$\langle \Delta H \rangle$ within the RPS plane $H = -xyz$.

$$\begin{aligned}\langle \Delta H_{RPS} \rangle = \frac{2}{N^6} \sum_{i=1}^{N} \sum_{j=1}^{N-i} \sum_{k=1}^{N-i-j} \Big[& ijk(T^{RP} + T^{RS} + T^{R+} + T^{PR} + T^{PS} + T^{P+} \\ & + T^{SR} + T^{SP} + T^{S+} + T^{+R} + T^{+P} + T^{+S}) \\ & - k(i-1)(j+1)T^{RP} - (i-1)j(k+1)T^{RS} - jk(i-1)T^{R+} \\ & - k(i+1)(j-1)T^{PR} - i(j-1)(k+1)T^{PS} - ik(j-1)T^{P+} \\ & - (i+1)j(k-1)T^{SR} - i(j+1)(k-1)T^{SP} - ij(k-1)T^{S+} \\ & - jk(i+1)T^{+R} - ik(j+1)T^{+P} - ij(k+1)T^{+S} \Big]\end{aligned}$$

$$\begin{aligned}\langle \Delta H_{RPS} \rangle = \frac{2}{N^6} \sum_{i=1}^{N} \sum_{j=1}^{N-i} \sum_{k=1}^{N-i-j} \Big[& k(j-i)(T^{RP} - T^{PR}) + j(k-i)(T^{RS} - T^{SR}) + i(j-k)(T^{SP} - T^{PS}) \\ & + k(T^{RP} + T^{PR}) + j(T^{RS} + T^{SR}) + i(T^{SP} + T^{PS}) \\ & + jk(T^{R+} - T^{+R}) + ik(T^{P+} - T^{+P}) + ij(T^{S+} - T^{+S}) \Big]\end{aligned} \tag{27}$$

Compared to the derivation of simply the RPS case in[**?**], this has the additional terms including the difference in tranition probabilities in and out of the 4th strategy.

$$
\begin{aligned}
\langle \Delta H_{RPS} \rangle = \frac{2}{N^3} \int_0^1 dx \int_0^{1-x} dy \int_0^{1-x-y} dz \Big[& N^2 z(y-x)(T^{RP} - T^{PR}) + N^2 y(z-x)(T^{RS} - T^{SR}) \\
& + N^2 x(y-z)(T^{SP} - T^{PS}) \\
& + Nz(T^{RP} + T^{PR}) + Ny(T^{RS} + T^{SR}) + Nx(T^{SP} + T^{PS}) \\
& + N^2 xz(T^{P+} - T^{+P}) + N^2 yz(T^{R+} - T^{+R}) + N^2 xy(T^{S+} - T^{+S}) \Big]
\end{aligned}
$$

$$
\begin{aligned}
\langle \Delta H_{RPS} \rangle = \frac{2}{N} \int_0^1 dx \int_0^{1-x} dy \int_0^{1-x-y} dz \Big[& z(y-x)(T^{RP} - T^{PR}) + y(z-x)(T^{RS} - T^{SR}) \\
& + x(y-z)(T^{SP} - T^{PS}) + xz(T^{P+} - T^{+P}) + yz(T^{R+} - T^{+R}) + xy(T^{S+} - T^{+S}) \Big] \\
& + \frac{2}{N^2} \int_0^1 dx \int_0^{1-x} dy \int_0^{1-x-y} dz \Big[z(T^{RP} + T^{PR}) + y(T^{RS} + T^{SR}) + x(T^{SP} + T^{PS}) \Big]
\end{aligned} \tag{28}
$$

0.0.4 Analytical solution for the Local update process $\langle \Delta H_{RPS} \rangle$

An analytical solution for $\langle \Delta H_{RPS} \rangle$ can also be derived for for the local update process, this will require the simplified terms of the forms $T^{RP} - T^{PR}$ and $T^{RP} + T^{PR}$ within the RPS game, and the terms of form $T^{R+} - T^{+R}$ have already been derived in the solution for $\langle \Delta H_{SD} \rangle$.

$$
\begin{aligned}
T^{RP} - T^{PR} &= \left(\frac{1}{2} + \frac{w}{2} \left(\frac{\pi_P - \pi_R}{\Delta \pi_{max}} \right) \right) \frac{i}{N} \frac{j}{N} - \left(\frac{1}{2} + \frac{w}{2} \left(\frac{\pi_R - \pi_P}{\Delta \pi_{max}} \right) \right) \frac{i}{N} \frac{j}{N} \\
&= \frac{i}{N} \frac{j}{N} \left[\frac{w}{2\Delta \pi_{max}} \left(2(\pi_P - \pi_R) \right) \right] \\
&= \frac{w}{\Delta \pi_{max}} xy(\pi_P - \pi_R)
\end{aligned} \tag{29}
$$

where population counts e.g. $i/N, j/N$ are replaced with frequencies x, y respectively. Similarly the other 2 terms within the RPS game can be derived leading to:

$$
\begin{aligned}
T^{RS} - T^{SR} &= \\
T^{SP} - T^{PS} &=
\end{aligned} \tag{30}
$$

Substituting in the expanded payoff differences from the payoff matrix:
Rough figures of rps and SD delta H values.



Figure 2: Blue - SD



Figure 3: Double reversal case. Blue - SD



Figure 4: Comparison of the 3 drift measures $\langle \Delta H_{SD} \rangle, \langle \Delta H_{RPS} \rangle$, and$\langle \Delta H_{+} \rangle$ for the local update process. $\beta = 0.3, \gamma = 0.14, c = -0.2$, strong central attracting RPS case ($c > -1$) and $N_{RPS} < N_{SD}$ so SD game component reverses first, at a higher population size. This results in the vertical column point cloud where RPS fixates at the central fixed point where $x \approx y \approx z$ and q drifts away from the fixed point ($N_{RPS} < N < N_{SD}$)

Testing point cloud at N = 200 $\frac{N}{N_{SD}} = 0.14$, $\frac{N}{N_{RPS}} = 2.26$, rod case
For below at N = 300 $\frac{N}{N_{SD}} = 4.25$, $\frac{N}{N_{RPS}} = 0.54$
For final case at N = 2000 $\frac{N}{N_{SD}} = 4.76$, $\frac{N}{N_{RPS}} = 3.57$
Blob case N = 1000 $\frac{N}{N_{SD}} = 2.38$, $\frac{N}{N_{RPS}} = 1.79$ Final case at N = 30000, $\frac{N}{N_{SD}} = 71.43$, $\frac{N}{N_{RPS}} = 53.57$

N	N/N_{SD}	N/N_{RPS}
200	0.14	2.26
200	2.83	0.36
2000	4.76	3.57

Table 1: Table of N values and corresponding ratios, depending on the values of N_{SD}, N_{RPS} and N when simulating trajectories, the drift reversal cases will be different, large N used to show the cases above the critical population sizes where system converges to interior nash equilibrium.



Figure 5: Comparison of drift measures for local update process, where $\beta = 0.4, \gamma = 0.2, c = -0.8$. Still an interior attracting RPS in large population case, however this time RPS drifts at a higher population size marked by N'_{RPS} so drifts to RPS boundaries whilst SD does not reverse and stays at the fixed point for SD part of the game. Results in disk shape around the central fixed point as RPS goes to boundary while SD is stable in the interior.

For either of these cases, when N is below the lower of the two critical population sizes, this results in a double reversal so the trajectories end up being absorbed by the four corners of the simplex.



Figure 6: $\beta = 0.1, \gamma = 0.1, c = -0.8, N_{SD} = 420, N_{RPS} = 560$



Figure 7: SD drift case N = 200



Figure 8: RPS drift case N = 300



0.0.5 $\langle \Delta H_4 \rangle$ Derivation

All 4 strategies, $H = -xyz(1-x-y-z)$, Therefore $\langle \Delta H_4 \rangle$ for the population counts can be defined as:

$$\begin{aligned}
\langle \Delta H_4 \rangle = \frac{6}{N^7} \sum_{i=1}^{N} \sum_{j=1}^{N-i} \sum_{k=1}^{N-i-j} \Big[& ijk(N-i-j-k)(T^{RP} + T^{RS} + T^{R+} + T^{PR} + T^{PS} + T^{P+} \\
& + T^{SR} + T^{SP} + T^{S+} + T^{+R} + T^{+P} + T^{+S}) - (i-1)(j+1)k(N-i-j-k)T^{RP} \\
& - (i-1)j(k+1)(N-i-j-k)T^{RS} - (i-1)jk(N-i-j-k+1)T^{R+} \\
& - (i+1)(j-1)k(N-i-j-k)T^{PR} - i(j-1)(k+1)(N-i-j-k)T^{PS} \\
& - i(j-1)k(N-i-j-k+1)T^{P+} - (i+1)j(k-1)(N-i-j-k)T^{SR} \\
& - i(j+1)(k-1)(N-i-j-k)T^{SP} - ij(k-1)(N-i-j-k+1)T^{S+} \\
& - (i+1)jk(N-1-i-j-k)T^{+R} - i(j+1)k(N-1-i-j-k)T^{+P} - ij(k+1)(N-1-i-j-k)T^{+S} \Big]
\end{aligned} \tag{31}$$

Normalization $\frac{6}{N^7}$, as its over the whole simplex (pyramid volume 1/6) and triple summation, (N^3), then 4 populations N^4.

Let p = N - i - j - k (4th strategy count):

$$\begin{aligned}\langle \Delta H_4 \rangle = \frac{6}{N^7} \sum_{i=1}^{N} \sum_{j=1}^{N-i} \sum_{k=1}^{N-i-j} \Big[& ijkp(T^{RP} + T^{RS} + T^{R+} + T^{PR} + T^{PS} + T^{P+} \\ & + T^{SR} + T^{SP} + T^{S+} + T^{+R} + T^{+P} + T^{+S}) - (i-1)(j+1)kpT^{RP} \\ & - (i-1)j(k+1)pT^{RS} - (i-1)jk(p+1)T^{R+} \\ & - (i+1)(j-1)kpT^{PR} - i(j-1)(k+1)pT^{PS} \\ & - i(j-1)k(p+1)T^{P+} - (i+1)j(k-1)pT^{SR} \\ & - i(j+1)(k-1)pT^{SP} - ij(k-1)(p+1)T^{S+} \\ & - (i+1)jk(p-1)T^{+R} - i(j+1)k(p-1)T^{+P} - ij(k+1)(p-1)T^{+S} \Big]\end{aligned} \tag{32}$$

Looking at one pair of transitions in and out of the same state. ($R \to P, P \to R$) this can be rewritten as before for $\langle \Delta H_{RPS} \rangle$ as differences of transition probabilities in and out of a state.

$$\begin{aligned}\langle \Delta H_4 \rangle = \frac{6}{N^7} \sum_{i=1}^{N} \sum_{j=1}^{N-i} \sum_{k=1}^{N-i-j} \Big[& kp(j-i)(T^{RP} - T^{PR}) + jp(k-i)(T^{RS} - T^{SR}) \\ & + ip(j-k)(T^{SP} - T^{PS}) + kp(T^{RP} + T^{PR}) + jp(T^{RS} + T^{SR}) \\ & + ip(T^{SP} + T^{PS}) + jk(p-i)(T^{R+} - T^{+R}) + ik(p-j)(T^{P+} - T^{+P}) \\ & + ij(p-k)(T^{S+} - T^{+S}) + jk(T^{R+} + T^{+R}) + ik(T^{P+} + T^{+P}) + ij(T^{S+} + T^{+S}) \Big]\end{aligned} \tag{33}$$

As before, converting this to the integral form where $x = i/N, y = j/N, z = k/N$ and $q = p/N, p = Nq$ where $q = 1 - x - y - z$ leads to:

$$\begin{aligned}\langle \Delta H_4 \rangle = \frac{6}{N^4} \int_0^1 dx \int_0^{1-x} dy \int_0^{1-x-y} dz \Big[& N^3 zq(y-x)(T^{RP} - T^{PR}) + N^3 yq(z-x)(T^{RS} - T^{SR}) \\ & + N^3 xq(y-z)(T^{SP} - T^{PS}) + N^2 zq(T^{RP} + T^{PR}) + N^2 yq(T^{RS} + T^{SR}) + N^2 xq(T^{SP} + T^{PS}) \\ & + N^3 yz(q-x)(T^{R+} - T^{+R}) + N^3 xz(q-y)(T^{P+} - T^{+P}) + N^3 xy(q-z)(T^{S+} - T^{+S}) \\ & + N^2 yz(T^{R+} + T^{+R}) + N^2 xz(T^{P+} + T^{+P}) + N^2 xy(T^{S+} + T^{+S}) \Big]\end{aligned} \tag{34}$$

$$\begin{aligned}\langle \Delta H_4 \rangle = & \frac{6}{N} \int_0^1 dx \int_0^{1-x} dy \int_0^{1-x-y} dz \Big[zq(y-x)(T^{RP} - T^{PR}) + yq(z-x)(T^{RS} - T^{SR}) + xq(y-z)(T^{SP} - T^{PS}) \\ & + yz(q-x)(T^{R+} - T^{+R}) + xz(q-y)(T^{P+} - T^{+P}) + xy(q-z)(T^{S+} - T^{+S}) \Big] \\ & + \frac{6}{N^2} \int_0^1 dx \int_0^{1-x} dy \int_0^{1-x-y} dz \Big[zq(T^{RP} + T^{PR}) + yq(T^{RS} + T^{SR}) + xq(T^{SP} + T^{PS}) \\ & + yz(T^{R+} + T^{+R}) + xz(T^{P+} + T^{+P}) + xy(T^{S+} + T^{+S}) \Big]\end{aligned} \tag{35}$$

Final derivation of $\langle \Delta H_4 \rangle$. There are clear similarities to the other two expected values $\langle \Delta H_{SD} \rangle, \langle \Delta_{RPS} \rangle$ with some terms missing due to the different observation variable H used but they can all be rewritten in terms of differences of transition probabilities into and out of state (e.g. $(T^{RP} - T^{PR})$) in an integral scaled by $1/N$ plus positive terms in a second integral scaled by $1/N^2$.

Perhaps also derive analytical form of this equation too for LOCAl update

0.0.6 Graphical comparison of H

Discuss how the H4 is a composition of the drift and diffusion terms of the other two expected delta H.

Point cloud - look at how H-RPS shows the attraction or repulsion of the fixed rps center, SD - attraction to the top point or the bottom RPS pure plane,

Then H4 - combination where it sign change would indicate whether or not the system is attracted to the central internal fixed point in the pyramid - where a sign change would suggest it pushed to one of the faces e.g. the rps plane where q = 0, step towards the boundary means larger H4 - so delta H addition of larger value so sign flip...

Change in sign in SD - up or down Change in sign in RPS - into out out of center or RPS plane Change in sign in H4 - away from internal fixed point in any direction

Compare the 3 together on same plot for some of the interesting cases.

$\langle \Delta H_4 \rangle$ seems to have a sign change when either the RPS or SD game does or both, double check the code copy of the formula, and see if theres a pattern as one does not seem obvious at the moment.