

A Proof of the Erdős–Gallai Theorem


Stephen G. Hartke

Department of Mathematics
University of Nebraska–Lincoln
www.math.unl.edu/~shartke2
hartke@math.unl.edu

Joint work with Tyler Seacrest

Degree Sequences

Def. The **degree sequence** π of a graph G is a list of its degrees (with multiplicity), usually listed in decreasing order.



$$\pi = (3, 3, 3, 2, 1, 0)$$


Graphic Sequences

Given a list π , when is it the **degree sequence** of some graph?

Graphic Sequences

Given a list π , when is it the **degree sequence** of some graph?


We only consider **simple** graphs:
no **loops** or **multiple edges**.



Graphic Sequences

Given a list π , when is it the **degree sequence** of some graph?

We only consider **simple** graphs:
no **loops** or **multiple edges**.




Def. A list π that is the **degree seq** of some **simple** graph is said to be a **graphic sequence**.

Graphic Sequences

Given a list π , when is it the **degree sequence** of some graph?

We only consider **simple** graphs:
no **loops** or **multiple edges**.



Def. A list π that is the **degree seq** of some **simple** graph is said to be a **graphic sequence**.


What lists are **graphic**?

Graphic Sequences

Ex. Is $\pi = (2, 2, 2, 2, 1, 1)$ graphic?

Graphic Sequences

Ex. Is $\pi = (2, 2, 2, 2, 1, 1)$ graphic? **Yes**



Graphic Sequences

Ex. Is $\pi = (2, 2, 2, 2, 1, 1)$ graphic? Yes

Ex. Is $\pi = (4, 3, 2, 2, 2, 1, 1)$ graphic?

Graphic Sequences

Ex. Is $\pi = (2, 2, 2, 2, 1, 1)$ graphic? **Yes**

Ex. Is $\pi = (4, 3, 2, 2, 2, 1, 1)$ graphic? **No**

For a **simple** graph, $\sum_{v \in V(G)} \deg(v) = 2|E(G)|$.

Hence, the sum of π must be **even**.

Graphic Sequences

Ex. Is $\pi = (2, 2, 2, 2, 1, 1)$ graphic? **Yes**

Ex. Is $\pi = (4, 3, 2, 2, 2, 1, 1)$ graphic? **No**

For a **simple** graph, $\sum_{v \in V(G)} \deg(v) = 2|E(G)|$.

Hence, the sum of π must be **even**.

Ex. Is $\pi = (6, 6, 5, 4, 4, 2, 1)$ graphic?

Graphic Sequences

Ex. Is $\pi = (2, 2, 2, 2, 1, 1)$ graphic? **Yes**

Ex. Is $\pi = (4, 3, 2, 2, 2, 1, 1)$ graphic? **No**

For a **simple** graph, $\sum_{v \in V(G)} \deg(v) = 2|E(G)|$.

Hence, the sum of π must be **even**.

Ex. Is $\pi = (6, 6, 5, 4, 4, 2, 1)$ graphic? **No** ... but **why**?

The Erdős–Gallai Theorem

Thm. [Erdős–Gallai 1960]

Let $\pi = (d_1 \geq d_2 \geq \dots \geq d_n)$ be a nonneg int list with **even sum**.

Then π is **graphic** if and only if

$$\text{for all } 1 \leq k \leq n, \quad \sum_{i=1}^k d_i \leq k(k-1) + \sum_{i=k+1}^n \min(d_i, k).$$

The Erdős–Gallai Theorem


Thm. [Erdős–Gallai 1960]

Let $\pi = (d_1 \geq d_2 \geq \dots \geq d_n)$ be a nonneg int list with **even sum**.

Then π is **graphic** if and only if

$$\text{for all } 1 \leq k \leq n, \quad \sum_{i=1}^k d_i \leq k(k-1) + \sum_{i=k+1}^n \min(d_i, k).$$

Why is this condition **necessary**?



The Erdős–Gallai Theorem


Thm. [Erdős–Gallai 1960]

Let $\pi = (d_1 \geq d_2 \geq \dots \geq d_n)$ be a nonneg int list with **even sum**.

Then π is **graphic** if and only if

$$\text{for all } 1 \leq k \leq n, \quad \sum_{i=1}^k d_i \leq k(k-1) + \sum_{i=k+1}^n \min(d_i, k).$$

Why is this condition **necessary**?



The **left side** counts **degree** among the highest k vertices.

The Erdős–Gallai Theorem


Thm. [Erdős–Gallai 1960]

Let $\pi = (d_1 \geq d_2 \geq \dots \geq d_n)$ be a nonneg int list with **even sum**.

Then π is **graphic** if and only if

$$\text{for all } 1 \leq k \leq n, \quad \sum_{i=1}^k d_i \leq k(k-1) + \sum_{i=k+1}^n \min(d_i, k).$$

Why is this condition **necessary**?



At most $k(k-1)$ can be from edges among the highest k vertices.

The Erdős–Gallai Theorem


Thm. [Erdős–Gallai 1960]

Let $\pi = (d_1 \geq d_2 \geq \dots \geq d_n)$ be a nonneg int list with **even sum**.

Then π is **graphic** if and only if

$$\text{for all } 1 \leq k \leq n, \quad \sum_{i=1}^k d_i \leq k(k-1) + \sum_{i=k+1}^n \min(d_i, k).$$

Why is this condition **necessary**?



Each remaining vertex can **absorb** at most k or its **degree**.

Nongraphic Sequence

Ex. Is $\pi = (6, 6, 5, 4, 4, 2, 1)$ graphic?

Nongraphic Sequence

Ex. Is $\pi = (6, 6, 5, 4, 4, 2, 1)$ graphic?

For $k = 2$, check

$$\sum_{i=1}^k d_i \leq k(k-1) + \sum_{i=k+1}^n \min(d_i, k)$$
$$6 + 6 \leq 2(1) + (2 + 2 + 2 + 2 + 1)$$
$$12 \geq 11$$

Nongraphic Sequence

Ex. Is $\pi = (6, 6, 5, 4, 4, 2, 1)$ graphic?

For $k = 2$, check

$$\sum_{i=1}^k d_i \leq k(k-1) + \sum_{i=k+1}^n \min(d_i, k)$$
$$6 + 6 \leq 2(1) + (2 + 2 + 2 + 2 + 1)$$
$$12 \geq 11$$

So π is **not** graphic!

Sufficiency

Why are the EG inequalities **sufficient**?

Sufficiency

Why are the EG inequalities **sufficient**?

Excursus. Everything is better in **bipartite** land!

Sufficiency

Why are the EG inequalities **sufficient**?

Excursus. Everything is better in **bipartite** land!


Def. A bilist $\tau = (a_1 \geq \dots \geq a_n; b_1 \geq \dots \geq b_m)$ is **bigraphic** if it is the degree sequence of some **simple bipartite** graph.

Sufficiency

Why are the EG inequalities **sufficient**?

Excursus. Everything is better in **bipartite** land!

Def. A bilist $\tau = (a_1 \geq \dots \geq a_n; b_1 \geq \dots \geq b_m)$ is **bigraphic** if it is the degree sequence of some **simple bipartite** graph.



$$\tau = (2, 2, 2, 1; 4, 3, 0)$$

Note that the a_i s are
the degrees in one part,
and the b_j s are the degrees
in the other part.

Bigraphic Sequences

A **necessary** condition:

If a bigraph has degrees $\tau = (a_1 \geq \dots \geq a_n; b_1 \geq \dots \geq b_m)$,

then any realization has $\sum_{i=1}^n a_i = |E(G)|$

Bigraphic Sequences

A **necessary** condition:

If a bigraph has degrees $\tau = (a_1 \geq \dots \geq a_n; b_1 \geq \dots \geq b_m)$,

then any realization has $\sum_{i=1}^n a_i = |E(G)| = \sum_{j=1}^m b_j$.

Bigraphic Sequences

A **necessary** condition:

If a bigraph has degrees $\tau = (a_1 \geq \dots \geq a_n; b_1 \geq \dots \geq b_m)$,

then any realization has $\sum_{i=1}^n a_i = |E(G)| = \sum_{j=1}^m b_j$.

Thm. [Gale 1957, Ryser 1957]

Let $\tau = (a_1 \geq \dots \geq a_n; b_1 \geq \dots \geq b_m)$ be a bilist with $\sum a_i = \sum b_j$.

Then τ is **bigraphic** if and only if

$$\text{for all } 1 \leq k \leq n, \quad \sum_{i=1}^k a_i \leq \sum_{j=1}^m \min(b_j, k).$$

Necessity of Gale–Ryser Theorem

Let $\tau = (a_1 \geq \dots \geq a_n; b_1 \geq \dots \geq b_m)$ be a bilist with $\sum a_i = \sum b_j$.

Then τ bigraphic \Leftrightarrow for all $1 \leq k \leq n$, $\sum_{i=1}^k a_i \leq \sum_{j=1}^m \min(b_j, k)$.

a_1 •

• b_1

a_2 •

• b_2

a_3 •

• b_3

a_4 •

• b_4


a_5 •

• b_5

Necessity of Gale–Ryser Theorem

Let $\tau = (a_1 \geq \dots \geq a_n; b_1 \geq \dots \geq b_m)$ be a bilist with $\sum a_i = \sum b_j$.

Then τ bigraphic \Leftrightarrow for all $1 \leq k \leq n$, $\sum_{i=1}^k a_i \leq \sum_{j=1}^m \min(b_j, k)$.




- b_1
- b_2
- b_3
- b_4
- b_5

Left side counts degree among the highest k vertices in A .

Necessity of Gale–Ryser Theorem

Let $\tau = (a_1 \geq \dots \geq a_n; b_1 \geq \dots \geq b_m)$ be a bilist with $\sum a_i = \sum b_j$.

Then τ bigraphic \Leftrightarrow for all $1 \leq k \leq n$, $\sum_{i=1}^k a_i \leq \sum_{j=1}^m \min(b_j, k)$.



Left side counts degree among the highest k vertices in A .

Each vertex in B can absorb at most k or its degree.

The Gale–Ryser Theorem

Many proofs of sufficiency—my favorite uses network flows.

2 ● ● 4


2 ● ● 3

2 ● ● 0

1 ●


The Gale–Ryser Theorem

Many proofs of sufficiency—my favorite uses network flows.



The Gale–Ryser Theorem

Many proofs of sufficiency—my favorite uses network flows.



The middle edges with flow 1 realize the bipartite graph.

The Gale–Ryser Theorem

Now apply **Max Flow–Min Cut Theorem** to the previous network.

Thm. [Ford–Fulkerson 1956]

The **maximum flow** from the source to the sink is equal to the **minimum size** of a source/sink separating cut.

The Gale–Ryser Theorem

Now apply **Max Flow–Min Cut Theorem** to the previous network.

Thm. [Ford–Fulkerson 1956]

The **maximum flow** from the source to the sink is equal to the **minimum size** of a source/sink separating cut.

The checking of the **Gale–Ryser** conditions is **straightforward**, but involves some cases.

Nonbipartite?

How to relate the **nonbipartite** world to **bipartite** land?

Durfee square

Def. Given a partition $d_1 \geq d_2 \geq \dots \geq d_n$, the Durfee square number m is the largest i such that $d_i \geq i$.

$$d_1 = 4 \quad \bullet \quad \bullet \quad \bullet \quad \bullet$$


$$d_2 = 2 \quad \bullet \quad \bullet$$

$$d_3 = 2 \quad \bullet \quad \bullet$$

$$d_4 = 1 \quad \bullet$$

Durfee square

Def. Given a partition $d_1 \geq d_2 \geq \dots \geq d_n$, the Durfee square number m is the largest i such that $d_i \geq i$.



Ryser Condition

Thm. [Ryser 1957?]

Let $\pi = (d_1 \geq d_2 \geq \dots \geq d_n)$ be a list with Durfee square m .

Let $\tau = (\tilde{d}_1, \dots, \tilde{d}_n; \tilde{d}_1, \dots, \tilde{d}_n)$ where $\tilde{d}_i = \begin{cases} d_i + 1 & \text{if } d_i \leq m, \\ d_i & \text{if } d_i > m. \end{cases}$

Then π is graphic if and only if τ is bigraphic.


Ryser Condition

Thm. [Ryser 1957?]

Let $\pi = (d_1 \geq d_2 \geq \dots \geq d_n)$ be a list with Durfee square m .

Let $\tau = (\tilde{d}_1, \dots, \tilde{d}_n; \tilde{d}_1, \dots, \tilde{d}_n)$ where $\tilde{d}_i = \begin{cases} d_i + 1 & \text{if } d_i \leq m, \\ d_i & \text{if } d_i > m. \end{cases}$

Then π is graphic if and only if τ is bigraphic.




Ryser Condition

Thm. [Ryser 1957?]

Let $\pi = (d_1 \geq d_2 \geq \dots \geq d_n)$ be a list with Durfee square m .

Let $\tau = (\tilde{d}_1, \dots, \tilde{d}_n; \tilde{d}_1, \dots, \tilde{d}_n)$ where $\tilde{d}_i = \begin{cases} d_i + 1 & \text{if } d_i \leq m, \\ d_i & \text{if } d_i > m. \end{cases}$

Then π is **graphic** if and only if τ is **bigraphic**.




Ryser Condition

Thm. [Ryser 1957?]

Let $\pi = (d_1 \geq d_2 \geq \dots \geq d_n)$ be a list with Durfee square m .

Let $\tau = (\tilde{d}_1, \dots, \tilde{d}_n; \tilde{d}_1, \dots, \tilde{d}_n)$ where $\tilde{d}_i = \begin{cases} d_i + 1 & \text{if } d_i \leq m, \\ d_i & \text{if } d_i > m. \end{cases}$

Then π is graphic if and only if τ is bigraphic.




Ryser Condition

Thm. [Ryser 1957?]

Let $\pi = (d_1 \geq d_2 \geq \dots \geq d_n)$ be a list with Durfee square m .

Let $\tau = (\tilde{d}_1, \dots, \tilde{d}_n; \tilde{d}_1, \dots, \tilde{d}_n)$ where $\tilde{d}_i = \begin{cases} d_i + 1 & \text{if } d_i \leq m, \\ d_i & \text{if } d_i > m. \end{cases}$


Then π is graphic if and only if τ is bigraphic.




Ryser Condition

Proof that τ **bigraphic** implies π is **graphic**:


Case 1



Case 2




Case 3



Ryser Condition

Proof that τ bigraphic implies π is graphic:


Case 1



Ryser Condition

Proof that τ bigraphic implies π is graphic:


Case 1



Ryser Condition

Proof that τ bigraphic implies π is graphic:


Case 1



Ryser Condition

Proof that τ bigraphic implies π is graphic:


Case 1



Ryser Condition

Proof that τ bigraphic implies π is graphic:

Case 1



Ryser Condition

Proof that τ bigraphic implies π is graphic:

Case 1


All the instances of Case 1 can be handled similarly.

All instances of Case 1 are handled, and then Cases 2 and 3.

Ryser Condition

Cases 2 and 3: all instances of Case 1 handled.


By parity, there are an even number of **bad** edges.



Ryser Condition

Cases 2 and 3: all instances of Case 1 handled.

By parity, there are an even number of **bad** edges.



Case A: $i \leq m$ and $j > m$


We can find v_k so that

$v_i \leftrightarrow v_k$ but $v_j \nleftrightarrow v_k$.

Ryser Condition

Cases 2 and 3: all instances of Case 1 handled.

By parity, there are an even number of **bad** edges.



Case A: $i \leq m$ and $j > m$


We can find v_k so that

$v_i \leftrightarrow v_k$ but $v_j \nleftrightarrow v_k$.

Ryser Condition

Cases 2 and 3: all instances of Case 1 handled.

By parity, there are an even number of **bad** edges.



Case B: $i, j \leq m$


We can find $k > m$
so that $v_i \leftrightarrow v_k$.

Now proceed with Case A.

Ryser Condition

Cases 2 and 3: all instances of Case 1 handled.

By parity, there are an even number of **bad** edges.



Case B: $i, j \leq m$


We can find $k > m$
so that $v_i \leftrightarrow v_k$.

Now proceed with Case A.

Ryser Condition

Cases 2 and 3: all instances of Case 1 handled.

By parity, there are an even number of **bad** edges.



Case C: $i, j > m$


We can find $k < m$
so that $v_i \not\leftrightarrow v_k$.

Now proceed with Case A.

Ryser Condition

Cases 2 and 3: all instances of Case 1 handled.

By parity, there are an even number of **bad** edges.



Case C: $i, j > m$

We can find $k < m$
so that $v_i \not\leftrightarrow v_k$.

Now proceed with Case A.

Strengthened Erdős–Gallai

Thm. [Zverovich–Zverovich 1992]

It is sufficient to check the first m EG inequalities,
where m is the Durfee square number.

That is,

Let $\pi = (d_1 \geq d_2 \geq \dots \geq d_n)$ be a nonneg int list with even sum.

Then π is graphic if and only if

$$\text{for all } 1 \leq k \leq m, \quad \sum_{i=1}^k d_i \leq k(k-1) + \sum_{i=k+1}^n \min(d_i, k).$$

Key point: for $k > m$, we have $\min(d_i, k) = d_i$.

Equivalent Gale–Ryser

Lem. Let $\tau = (\tilde{d}_1, \dots, \tilde{d}_n; \tilde{d}_1, \dots, \tilde{d}_n)$. Then

$$\sum_{i=1}^k \tilde{d}_i \leq \sum_{i=1}^n \min(\tilde{d}_i, k) \text{ iff } \sum_{i=1}^k \max(k, \tilde{d}_i) \leq k^2 + \sum_{i=k+1}^n \min(\tilde{d}_i, k).$$

Equivalent Gale–Ryser

Lem. Let $\tau = (\tilde{d}_1, \dots, \tilde{d}_n; \tilde{d}_1, \dots, \tilde{d}_n)$. Then

$$\sum_{i=1}^k \tilde{d}_i \leq \sum_{i=1}^n \min(\tilde{d}_i, k) \text{ iff } \sum_{i=1}^k \max(k, \tilde{d}_i) \leq k^2 + \sum_{i=k+1}^n \min(\tilde{d}_i, k).$$

Proof.

Subtract first k terms from right side:

$$\sum_{i=1}^k \tilde{d}_i - \min(\tilde{d}_i, k) \leq \sum_{i=k+1}^n \min(\tilde{d}_i, k)$$

Equivalent Gale–Ryser

Lem. Let $\tau = (\tilde{d}_1, \dots, \tilde{d}_n; \tilde{d}_1, \dots, \tilde{d}_n)$. Then

$$\sum_{i=1}^k \tilde{d}_i \leq \sum_{i=1}^n \min(\tilde{d}_i, k) \text{ iff } \sum_{i=1}^k \max(k, \tilde{d}_i) \leq k^2 + \sum_{i=k+1}^n \min(\tilde{d}_i, k).$$

Proof.

Subtract first k terms from right side:

$$\begin{aligned} \sum_{i=1}^k \tilde{d}_i - \min(\tilde{d}_i, k) &\leq \sum_{i=k+1}^n \min(\tilde{d}_i, k) \\ \sum_{i=1}^k \max(\tilde{d}_i - k, 0) &\leq \sum_{i=k+1}^n \min(\tilde{d}_i, k) \end{aligned}$$

Equivalent Gale–Ryser

Lem. Let $\tau = (\tilde{d}_1, \dots, \tilde{d}_n; \tilde{d}_1, \dots, \tilde{d}_n)$. Then

$$\sum_{i=1}^k \tilde{d}_i \leq \sum_{i=1}^n \min(\tilde{d}_i, k) \text{ iff } \sum_{i=1}^k \max(k, \tilde{d}_i) \leq k^2 + \sum_{i=k+1}^n \min(\tilde{d}_i, k).$$

Proof.

Subtract first k terms from right side:

$$\sum_{i=1}^k \tilde{d}_i - \min(\tilde{d}_i, k) \leq \sum_{i=k+1}^n \min(\tilde{d}_i, k)$$

$$\sum_{i=1}^k \max(\tilde{d}_i - k, 0) \leq \sum_{i=k+1}^n \min(\tilde{d}_i, k)$$

$$\sum_{i=1}^k \max(\tilde{d}_i, k) \leq k^2 + \sum_{i=k+1}^n \min(\tilde{d}_i, k)$$



Strengthened Gale–Ryser Inequalities

Lem.

It is sufficient to check the first m GR inequalities for τ ,
where m is the Durfee square number for π .

That is,

Let $\tau = (\tilde{d}_1, \dots, \tilde{d}_n; \tilde{d}_1, \dots, \tilde{d}_n)$.

Then τ is bigraphic if and only if

$$\text{for all } 1 \leq k \leq m, \quad \sum_{i=1}^k \tilde{d}_i \leq \sum_{i=k+1}^n \min(\tilde{d}_i, k).$$

This is more complicated than for the ER inequalities.

Strengthened Gale–Ryser

Lem. Let $\tau = (\tilde{d}_1, \dots, \tilde{d}_n; \tilde{d}_1, \dots, \tilde{d}_n)$.

Then τ bigraphic iff $\forall 1 \leq k \leq m, \sum_{i=1}^k \tilde{d}_i \leq \sum_{i=k+1}^n \min(\tilde{d}_i, k)$.

Proof. (\Leftarrow) We induct on k to show the remaining GR ineq.

Strengthened Gale–Ryser

Lem. Let $\tau = (\tilde{d}_1, \dots, \tilde{d}_n; \tilde{d}_1, \dots, \tilde{d}_n)$.

Then τ bigraphic iff $\forall 1 \leq k \leq m, \sum_{i=1}^k \tilde{d}_i \leq \sum_{i=k+1}^n \min(\tilde{d}_i, k)$.

Proof. (\Leftarrow) We induct on k to show the remaining GR ineq.

Let $k > m$ (so $d_k < k$). Let ℓ be largest so that $d_\ell \geq k$ (note $\ell < k$).

Strengthened Gale–Ryser

Lem. Let $\tau = (\tilde{d}_1, \dots, \tilde{d}_n; \tilde{d}_1, \dots, \tilde{d}_n)$.

Then τ bigraphic iff $\forall 1 \leq k \leq m, \sum_{i=1}^k \tilde{d}_i \leq \sum_{i=k+1}^n \min(\tilde{d}_i, k)$.

Proof. (\Leftarrow) We induct on k to show the remaining GR ineq.
Let $k > m$ (so $d_k < k$). Let ℓ be largest so that $d_\ell \geq k$ (note $\ell < k$).

$$\sum_{i=1}^k \max(\tilde{d}_i, k) = \sum_{i=1}^{\ell} \tilde{d}_i + k(k - \ell)$$

Strengthened Gale–Ryser

Lem. Let $\tau = (\tilde{d}_1, \dots, \tilde{d}_n; \tilde{d}_1, \dots, \tilde{d}_n)$.

Then τ bigraphic iff $\forall 1 \leq k \leq m, \sum_{i=1}^k \tilde{d}_i \leq \sum_{i=k+1}^n \min(\tilde{d}_i, k)$.

Proof. (\Leftarrow) We induct on k to show the remaining GR ineq.
Let $k > m$ (so $d_k < k$). Let ℓ be largest so that $d_\ell \geq k$ (note $\ell < k$).

$$\begin{aligned} \sum_{i=1}^k \max(\tilde{d}_i, k) &= \sum_{i=1}^{\ell} \tilde{d}_i + k(k - \ell) \\ &\leq \sum_{i=1}^n \min(\tilde{d}_i, \ell) + k(k - \ell) \quad \text{by induction} \end{aligned}$$

Strengthened Gale–Ryser

Lem. Let $\tau = (\tilde{d}_1, \dots, \tilde{d}_n; \tilde{d}_1, \dots, \tilde{d}_n)$.

Then τ bigraphic iff $\forall 1 \leq k \leq m, \sum_{i=1}^k \tilde{d}_i \leq \sum_{i=k+1}^n \min(\tilde{d}_i, k)$.

Proof. (\Leftarrow) We induct on k to show the remaining GR ineq.
Let $k > m$ (so $d_k < k$). Let ℓ be largest so that $d_\ell \geq k$ (note $\ell < k$).

$$\begin{aligned} \sum_{i=1}^k \max(\tilde{d}_i, k) &= \sum_{i=1}^{\ell} \tilde{d}_i + k(k - \ell) \\ &\leq \sum_{i=1}^n \min(\tilde{d}_i, \ell) + k(k - \ell) \quad \text{by induction} \\ &= k^2 - k\ell + \sum_{i=1}^k \min(\tilde{d}_i, \ell) + \sum_{i=k+1}^n \min(\tilde{d}_i, \ell) \end{aligned}$$

Strengthened Gale–Ryser

Lem. Let $\tau = (\tilde{d}_1, \dots, \tilde{d}_n; \tilde{d}_1, \dots, \tilde{d}_n)$.

Then τ bigraphic iff $\forall 1 \leq k \leq m, \sum_{i=1}^k \tilde{d}_i \leq \sum_{i=k+1}^n \min(\tilde{d}_i, k)$.

Proof. (\Leftarrow) We induct on k to show the remaining GR ineq.
Let $k > m$ (so $d_k < k$). Let ℓ be largest so that $d_\ell \geq k$ (note $\ell < k$).

$$\begin{aligned} \sum_{i=1}^k \max(\tilde{d}_i, k) &= \sum_{i=1}^{\ell} \tilde{d}_i + k(k - \ell) \\ &\leq \sum_{i=1}^n \min(\tilde{d}_i, \ell) + k(k - \ell) \quad \text{by induction} \\ &= k^2 - k\ell + \sum_{i=1}^k \min(\tilde{d}_i, \ell) + \sum_{i=k+1}^n \min(\tilde{d}_i, \ell) \\ &\leq k^2 + \sum_{i=k+1}^n \min(\tilde{d}_i, \ell) \leq k^2 + \sum_{i=k+1}^n \min(\tilde{d}_i, k) \quad \blacksquare \end{aligned}$$

Proving Erdős–Gallai

Sufficiency: We need to verify the first m GR ineq for τ .

Proving Erdős–Gallai

Sufficiency: We need to verify the first m GR ineq for τ .

Let $1 \leq k \leq m$.

$$\sum_{i=1}^k \tilde{d}_i = \sum_{i=1}^k d_i + k$$

Proving Erdős–Gallai

Sufficiency: We need to verify the first m GR ineq for τ .

Let $1 \leq k \leq m$.

$$\sum_{i=1}^k \tilde{d}_i = \sum_{i=1}^k d_i + k \leq k(k-1) + \sum_{i=k+1}^n \min(d_i, k) + k$$

Proving Erdős–Gallai

Sufficiency: We need to verify the first m GR ineq for τ .

Let $1 \leq k \leq m$.

$$\begin{aligned}\sum_{i=1}^k \tilde{d}_i &= \sum_{i=1}^k d_i + k \leq k(k-1) + \sum_{i=k+1}^n \min(d_i, k) + k \\ &\leq k^2 + \sum_{i=k+1}^n \min(d_i, k)\end{aligned}$$

Proving Erdős–Gallai

Sufficiency: We need to verify the first m GR ineq for τ .

Let $1 \leq k \leq m$.

$$\begin{aligned}\sum_{i=1}^k \tilde{d}_i &= \sum_{i=1}^k d_i + k \leq k(k-1) + \sum_{i=k+1}^n \min(d_i, k) + k \\ &\leq k^2 + \sum_{i=k+1}^n \min(d_i, k) \\ &\leq \sum_{i=1}^k k + \sum_{i=k+1}^n \min(d_i, k)\end{aligned}$$

Proving Erdős–Gallai

Sufficiency: We need to verify the first m GR ineq for τ .

Let $1 \leq k \leq m$.

$$\begin{aligned}\sum_{i=1}^k \tilde{d}_i &= \sum_{i=1}^k d_i + k \leq k(k-1) + \sum_{i=k+1}^n \min(d_i, k) + k \\ &\leq k^2 + \sum_{i=k+1}^n \min(d_i, k) \\ &\leq \sum_{i=1}^k k + \sum_{i=k+1}^n \min(d_i, k) \\ &\leq \sum_{i=1}^n \min(d_i, k)\end{aligned}$$

Proving Erdős–Gallai

Sufficiency: We need to verify the first m GR ineq for τ .

Let $1 \leq k \leq m$.

$$\begin{aligned}\sum_{i=1}^k \tilde{d}_i &= \sum_{i=1}^k d_i + k \leq k(k-1) + \sum_{i=k+1}^n \min(d_i, k) + k \\ &\leq k^2 + \sum_{i=k+1}^n \min(d_i, k) \\ &\leq \sum_{i=1}^k k + \sum_{i=k+1}^n \min(d_i, k) \\ &\leq \sum_{i=1}^n \min(d_i, k) \leq \sum_{i=1}^n \min(\tilde{d}_i, k).\end{aligned}$$

This completes the proof of the Erdős–Gallai Theorem! ■

A Proof of the Erdős–Gallai Theorem

Stephen G. Hartke

Department of Mathematics
University of Nebraska–Lincoln
www.math.unl.edu/~shartke2
hartke@math.unl.edu

Joint work with Tyler Seacrest