

School of Computer Science  
University of Birmingham

# Adelic Geometry via Geometric Logic

(joint work with Steve Vickers)

Ming Ng  
`mxn732@bham.ac.uk`

November 14, 2021

# What this talk is about



I'm going to discuss two basic themes that cross-cut many different areas of mathematics:



1. What kind of info can homotopical data encode?
2. When can we solve a problem by breaking it into smaller pieces?

I'll then discuss how the research project 'Adelic Geometry via Topos Theory' serves as an interesting test problem for illuminating how these two themes interact with each other.



# Homotopical Data = Geometric Data



# Homotopical Data = Geometric Data





# Homotopical Data = Geometric Data





- Question: Are these the only line bundles over  $S^1$  (up to isomorphism)?

# Homotopical Data = Geometric Data





- ▶ Question: Are these the only line bundles over  $S^1$  (up to isomorphism)?
- ▶ Answer: Yes.

# Homotopical Data = Geometric Data



- ▶ Question: Are these the only line bundles over  $S^1$  (up to isomorphism)?
- ▶ Answer: Yes.
- ▶ Why? Exploit the tight relationship between  $[S^{k-1}, GL_n(\mathbb{R})]$  and  $\text{Vect}_n(S^k)$ .

# Homotopical Data = Geometric Data



## Classification Theorem

Suppose that  $X$  is a paracompact space. Let  $\text{Vect}_n(X)$  be the set of isomorphism classes of  $n$ -dimensional vector bundles over  $X$ .

Then the map

$$[X, G_n] \longrightarrow \text{Vect}_n(X)$$

given by  $f \longmapsto f^*(\gamma_n)$  is a bijection, where  $\gamma_n$  is the universal bundle.





A similar attitude occurs in topos theory in regards to geometric logic:

axioms. This gives rise to a geometric *mathematics*, going beyond the merely logical – technically it is the mathematics that can be conducted in the topos-valid internal mathematics of Grothendieck toposes, and is moreover preserved by the inverse image functors of geometric morphisms. To put it another way, the geometric mathematics has an intrinsic continuity (since geometric morphisms are the continuous maps between toposes).

In this paper I shall survey some of the special features of geometric logic, and a body of established results that combine to support a manifesto “*continuity is geometricity*”. In other words, to “do mathematics continuously” is to work within the geometricity constraints.

— Vickers, 'Continuity and Geometric Logic'



## Theories and Models

A *theory* can be viewed as an axiomatic description of mathematical structures (e.g. the theory of groups);



## Theories and Models

A *theory* can be viewed as an axiomatic description of mathematical structures (e.g. the theory of groups); a *model* is an object that 'satisfies' these axioms.



## Theories and Models

A *theory* can be viewed as an axiomatic description of mathematical structures (e.g. the theory of groups); a *model* is an object that 'satisfies' these axioms.

## Geometric Theory

A *geometric theory* is a theory whose (formulae featured in its) axioms are built out of certain logical connectives



## Theories and Models

A *theory* can be viewed as an axiomatic description of mathematical structures (e.g. the theory of groups); a *model* is an object that 'satisfies' these axioms.

## Geometric Theory

A *geometric theory* is a theory whose (formulae featured in its) axioms are built out of certain logical connectives — i.e.  $=$ , finite conjunctions  $\wedge$ , arbitrary (possibly infinite) disjunctions  $\vee$ , and  $\exists$ .

# Example: Theory of Dedekind Reals



As an example, consider the geometric theory of Dedekind reals, which we denote  $\mathbb{R}$ .

# Example: Theory of Dedekind Reals



As an example, consider the geometric theory of Dedekind reals, which we denote  $\mathbb{R}$ . A model  $x$  of  $\mathbb{R}$  is a Dedekind real number, which will be represented by two sets of rationals  $(L, R)$ , whereby:

$$L = \{q \in \mathbb{Q} \mid q < x\}$$

$$R = \{r \in \mathbb{Q} \mid x < r\}$$

Otherwise known as the left and right Dedekind sections of the real number.



## Definition

- ▶ A *geometric morphism*  $f : \mathcal{F} \rightarrow \mathcal{E}$  of toposes is a pair of adjoint functors  $f_* : \mathcal{F} \rightarrow \mathcal{E}$  and  $f^* : \mathcal{E} \rightarrow \mathcal{F}$ , respectively called the *direct image* and the *inverse image* of  $f$ , such that the left adjoint  $f^*$  preserves finite limits.





## Definition

- ▶ A *geometric morphism*  $f : \mathcal{F} \rightarrow \mathcal{E}$  of toposes is a pair of adjoint functors  $f_* : \mathcal{F} \rightarrow \mathcal{E}$  and  $f^* : \mathcal{E} \rightarrow \mathcal{F}$ , respectively called the *direct image* and the *inverse image* of  $f$ , such that the left adjoint  $f^*$  preserves finite limits.



## Definition

- ▶ A *geometric morphism*  $f : \mathcal{F} \rightarrow \mathcal{E}$  of toposes is a pair of adjoint functors  $f_* : \mathcal{F} \rightarrow \mathcal{E}$  and  $f^* : \mathcal{E} \rightarrow \mathcal{F}$ , respectively called the *direct image* and the *inverse image* of  $f$ , such that the left adjoint  $f^*$  preserves finite limits.

## Definition

1. A *global point* of a topos  $\mathcal{E}$  is defined as a geometric morphism  $\mathbf{Set} \rightarrow \mathcal{E}$ .
2. A *generalised point* of a topos  $\mathcal{E}$  is a geometric morphism  $\mathcal{F} \rightarrow \mathcal{E}$ .



## Definition

The classifying topos of a geometric theory  $\mathbb{T}$  is a Grothendieck topos  $\mathbf{Set}[\mathbb{T}]$  that classifies the models of  $\mathbb{T}$  in Grothendieck toposes, i.e. for any Grothendieck topos  $\mathcal{E}$ , we have an equivalence of categories:

$$\mathbf{Geom}(\mathcal{E}, \mathbf{Set}[\mathbb{T}]) \simeq \mathbb{T}\text{-mod}(\mathcal{E})$$



## Definition

The classifying topos of a geometric theory  $\mathbb{T}$  is a Grothendieck topos  $\mathbf{Set}[\mathbb{T}]$  that classifies the models of  $\mathbb{T}$  in Grothendieck toposes, i.e. for any Grothendieck topos  $\mathcal{E}$ , we have an equivalence of categories:

$$\mathbf{Geom}(\mathcal{E}, \mathbf{Set}[\mathbb{T}]) \simeq \mathbb{T}\text{-mod}(\mathcal{E})$$

## Theorem

Every Grothendieck topos is a classifying topos of some geometric theory  $\mathbb{T}$ , and every geometric theory  $\mathbb{T}$  has a classifying topos.

# Topos = Generalised Space



## Definition

The classifying topos of a geometric theory  $\mathbb{T}$  is a Grothendieck topos  $\mathbf{Set}[\mathbb{T}]$  that classifies the models of  $\mathbb{T}$  in Grothendieck toposes, i.e. for any Grothendieck topos  $\mathcal{E}$ , we have an equivalence of categories:

$$\mathbf{Geom}(\mathcal{E}, \mathbf{Set}[\mathbb{T}]) \simeq \mathbb{T}\text{-mod}(\mathcal{E})$$

## Theorem

Every Grothendieck topos is a classifying topos of some geometric theory  $\mathbb{T}$ , and every geometric theory  $\mathbb{T}$  has a classifying topos.

## Slogan

Models = points of a topos. In particular, we can reason in terms of the points of the topos (as a generalised space) as opposed to only reasoning in terms of its objects/sheaves (as a category).



## Point-set Topology

- ▶ Point = Element of a set
- ▶ Space = A set of points, along with a set of opens satisfying some specific axioms.
- ▶ Continuous Maps = A function  $f : X \rightarrow Y$  such that  $f^{-1}(U)$  is open for all opens  $U \subset Y$

## Pointfree Topology

- ▶ Point = Model of a geometric theory
- ▶ Space = The 'World' in which the point lives with other points (i.e. a Grothendieck topos)
- ▶ Continuous Maps = A geometric morphism  $f : \mathcal{E} \rightarrow \mathcal{F}$  such that  $f^* : \mathcal{F} \rightarrow \mathcal{E}$  preserves finite limits and small colimits



## Fact

There exists a *generic model*  $U_{\mathbb{T}}$  living in every classifying topos,



## Fact

There exists a *generic model*  $U_{\mathbb{T}}$  living in every classifying topos, which possesses the universal property that any model  $M$  in a Grothendieck topos  $\mathcal{E}$  can be obtained as  $f^*(U_{\mathbb{T}}) \cong M$  via the inverse image functor of some (unique)  $f : \mathcal{E} \rightarrow \mathbf{Set}[\mathbb{T}]$ .






## Fact

There exists a *generic model*  $U_{\mathbb{T}}$  living in every classifying topos, which possesses the universal property that any model  $M$  in a Grothendieck topos  $\mathcal{E}$  can be obtained as  $f^*(U_{\mathbb{T}}) \cong M$  via the inverse image functor of some (unique)  $f : \mathcal{E} \rightarrow \mathbf{Set}[\mathbb{T}]$ .

An **important** consequence of this is that any geometric sequent that holds for  $U_{\mathbb{T}}$  will hold for all models  $M$  of  $\mathbb{T}$ .



Classifying topoi

# 'Divide-and-Conquer'



$$X^n + Y^n + Z^n = 0 \quad (n > 2)$$



$$X^n + Y^n + Z^n = 0 \quad (n > 2)$$

- Question: What are the rational (equiv. integer) solutions to this polynomial?



$$X^n + Y^n + Z^n = 0 \quad (n > 2)$$

- Question: What are the rational (equiv. integer) solutions to this polynomial? — hard!



$$X^n + Y^n + Z^n = 0 \quad (n > 2)$$

- ▶ Question: What are the rational (equiv. integer) solutions to this polynomial? — hard!
- ▶ Observation #1: Integer solutions imply real and modulo  $p$  solutions (in fact  $p$ -adic solutions).



$$X^n + Y^n + Z^n = 0 \quad (n > 2)$$

- ▶ Question: What are the rational (equiv. integer) solutions to this polynomial? — hard!
- ▶ Observation #1: Integer solutions imply real and modulo  $p$  solutions (in fact  $p$ -adic solutions).
- ▶ Observation #2: Real and  $p$ -adic solutions are easier to deal with than just integer/rational solutions.



$$X^n + Y^n + Z^n = 0 \quad (n > 2)$$

- ▶ Question: What are the rational (equiv. integer) solutions to this polynomial? — hard!
- ▶ Observation #1: Integer solutions imply real and modulo  $p$  solutions (in fact  $p$ -adic solutions).
- ▶ Observation #2: Real and  $p$ -adic solutions are easier to deal with than just integer/rational solutions.
- ▶ New Question: Given a polynomial with  $\mathbb{Q}$ -coefficients, when does knowledge about its  $\mathbb{Q}_p$  and  $\mathbb{R}$ -solutions give us info about its  $\mathbb{Q}$ -solutions?



# Hasse's Local-Global Principle



## Local-Global Principle for $\mathbb{Q}$

Some property  $P$  is true for  $\mathbb{Q}$  iff  $P$  is true for all the completions of  $\mathbb{Q}$ .

# Hasse's Local-Global Principle



## Local-Global Principle for $\mathbb{Q}$

Some property  $P$  is true for  $\mathbb{Q}$  iff  $P$  is true for all the completions of  $\mathbb{Q}$ .

## Definition of adèle ring for $\mathbb{Q}$

The adèle ring  $\mathbb{A}_{\mathbb{Q}}$  is defined to be the restricted product of all the completions of  $\mathbb{Q}$ . Morally, the adèle ring can be viewed as a device that allows us to reason about all the completions of  $\mathbb{Q}$  simultaneously.

# Hasse's Local-Global Principle



## Local-Global Principle for $\mathbb{Q}$

Some property  $P$  is true for  $\mathbb{Q}$  iff  $P$  is true for all the completions of  $\mathbb{Q}$ .

## Definition of adèle ring for $\mathbb{Q}$

The adèle ring  $\mathbb{A}_{\mathbb{Q}}$  is defined to be the restricted product of all the completions of  $\mathbb{Q}$ . Morally, the adèle ring can be viewed as a device that allows us to reason about all the completions of  $\mathbb{Q}$  simultaneously.

## Idea

Instead of asking whether a property simultaneously holds for *all completions* of  $\mathbb{Q}$  (which forces us to use complicated algebraic constructions like the adèle ring  $\mathbb{A}_{\mathbb{Q}}$ ), what if we asked whether a property holds for the *generic completion* of  $\mathbb{Q}$ ?



*“One weakness in the analogy between the collection of  $\{K_s\}_{s \in S}$  for a compact Riemann surface  $S$  and the collection  $\{\mathbb{Q}_p$ , for prime numbers  $p$ , and  $\mathbb{R}\}$  is that [...] no manner of squinting seems to be able to make  $\mathbb{R}$  the least bit mistake-able for any of the  $p$ -adic fields, nor are the  $p$ -adic fields  $\mathbb{Q}_p$  isomorphic for distinct  $p$ .*

***A major theme in the development of Number Theory has been to try to bring  $\mathbb{R}$  somewhat more into line with the  $p$ -adic fields; a major mystery is why  $\mathbb{R}$  resists this attempt so strenuously.”***

— Mazur, ‘Passage from Local to Global in Number Theory’

# Investigation begins...



Starting point:

For simplicity, let us assume that our base field is  $\mathbb{Q}$ . Classically, an absolute value of  $\mathbb{Q}$  is a function  $|\cdot| : \mathbb{Q} \rightarrow \mathbb{R}$  such that for all  $x, y \in \mathbb{Q}$ :

- ▶  $|x| \geq 0$ , and  $|x| = 0$  iff  $x = 0$

# Investigation begins...



Starting point:

For simplicity, let us assume that our base field is  $\mathbb{Q}$ . Classically, an absolute value of  $\mathbb{Q}$  is a function  $|\cdot| : \mathbb{Q} \rightarrow \mathbb{R}$  such that for all  $x, y \in \mathbb{Q}$ :

- ▶  $|x| \geq 0$ , and  $|x| = 0$  iff  $x = 0$
- ▶  $|xy| = |x||y|$

# Investigation begins...



Starting point:

For simplicity, let us assume that our base field is  $\mathbb{Q}$ . Classically, an absolute value of  $\mathbb{Q}$  is a function  $|\cdot| : \mathbb{Q} \rightarrow \mathbb{R}$  such that for all  $x, y \in \mathbb{Q}$ :

- ▶  $|x| \geq 0$ , and  $|x| = 0$  iff  $x = 0$
- ▶  $|xy| = |x||y|$
- ▶  $|x + y| \leq |x| + |y|$

# Investigation begins...



Starting point:

For simplicity, let us assume that our base field is  $\mathbb{Q}$ . Classically, an absolute value of  $\mathbb{Q}$  is a function  $|\cdot| : \mathbb{Q} \rightarrow \mathbb{R}$  such that for all  $x, y \in \mathbb{Q}$ :

- ▶  $|x| \geq 0$ , and  $|x| = 0$  iff  $x = 0$
- ▶  $|xy| = |x||y|$
- ▶  $|x + y| \leq |x| + |y|$

We define a *place* as an equivalence class of absolute values whereby  $|\cdot|_1 \sim |\cdot|_2$  if there exists some  $\alpha \in (0, 1]$  such that  $|\cdot|_1 = |\cdot|_2^\alpha$  or  $|\cdot|_2 = |\cdot|_1^\alpha$ .



# Classifying Topos of Places of $\mathbb{Q}$



- ▶ Intuitively: what does this topos look like?
- ▶ The points of this topos would correspond to equivalence classes of absolute values, such that:
  1.  $|\cdot|^\alpha \sim |\cdot|$for any absolute value  $|\cdot|$ , and  $\alpha \in (0, 1]$

# Classifying Topos of Places of $\mathbb{Q}$



- ▶ Intuitively: what does this topos look like?
- ▶ The points of this topos would correspond to equivalence classes of absolute values, such that:

1.  $|\cdot|^\alpha \sim |\cdot|$

for any absolute value  $|\cdot|$ , and  $\alpha \in (0, 1]$

$$[av] \times (0, 1] \begin{array}{c} \xrightarrow{\pi} \\ \xrightarrow{ex} \end{array} [av]$$



- ▶ Intuitively: what does this topos look like?
- ▶ The points of this topos would correspond to equivalence classes of absolute values, such that:

1.  $|\cdot|^\alpha \sim |\cdot|$

for any absolute value  $|\cdot|$ , and  $\alpha \in (0, 1]$

$$[av] \times (0, 1] \begin{array}{c} \xrightarrow{\pi} \\ \xrightarrow{ex} \end{array} [av]$$

- ▶  $\pi$  is the projection map sending  $(|\cdot|, \alpha) \mapsto |\cdot|$
- ▶  $ex$  is the exponentiation map sending  $(|\cdot|, \alpha) \mapsto |\cdot|^\alpha$

# Classifying Topos of Places of $\mathbb{Q}$



- ▶ Intuitively: what does this topos look like?
- ▶ The points of this topos would correspond to equivalence classes of absolute values, such that:
  1.  $|\cdot|^\alpha \sim |\cdot|$

# Classifying Topos of Places of $\mathbb{Q}$



- ▶ Intuitively: what does this topos look like?
- ▶ The points of this topos would correspond to equivalence classes of absolute values, such that:

1.  $|\cdot|^\alpha \sim |\cdot|$
2.  $|\cdot|^1 = |\cdot|$
3.  $(|\cdot|^\alpha)^\beta = |\cdot|^{\alpha\beta}$

for any absolute value  $|\cdot|$ , and  $\alpha, \beta \in (0, 1]$



- ▶ Intuitively: what does this topos look like?
- ▶ The points of this topos would correspond to equivalence classes of absolute values, such that:

1.  $|\cdot|^\alpha \sim |\cdot|$
2.  $|\cdot|^1 = |\cdot|$
3.  $(|\cdot|^\alpha)^\beta = |\cdot|^{\alpha\beta}$

for any absolute value  $|\cdot|$ , and  $\alpha, \beta \in (0, 1]$

- ▶ In essence, we would like to ‘quotient’ the topos  $[\mathbf{av}]$  by an algebraic action – two questions:



- ▶ Intuitively: what does this topos look like?
- ▶ The points of this topos would correspond to equivalence classes of absolute values, such that:

1.  $|\cdot|^\alpha \sim |\cdot|$
2.  $|\cdot|^1 = |\cdot|$
3.  $(|\cdot|^\alpha)^\beta = |\cdot|^{\alpha\beta}$

for any absolute value  $|\cdot|$ , and  $\alpha, \beta \in (0, 1]$

- ▶ In essence, we would like to ‘quotient’ the topos  $[\mathbf{av}]$  by an algebraic action – two questions:



- ▶ Intuitively: what does this topos look like?
- ▶ The points of this topos would correspond to equivalence classes of absolute values, such that:
  1.  $|\cdot|^\alpha \sim |\cdot|$
  2.  $|\cdot|^1 = |\cdot|$
  3.  $(|\cdot|^\alpha)^\beta = |\cdot|^{\alpha \cdot \beta}$for any absolute value  $|\cdot|$ , and  $\alpha, \beta \in (0, 1]$
- ▶ In essence, we would like to ‘quotient’ the topos  $[\mathbf{av}]$  by an algebraic action – two questions:
  - ▶ Is the notion of (real) exponentiation geometric? Ng-Vickers (2021)





- ▶ Intuitively: what does this topos look like?
- ▶ The points of this topos would correspond to equivalence classes of absolute values, such that:
  1.  $|\cdot|^\alpha \sim |\cdot|$
  2.  $|\cdot|^1 = |\cdot|$
  3.  $(|\cdot|^\alpha)^\beta = |\cdot|^{\alpha\beta}$for any absolute value  $|\cdot|$ , and  $\alpha, \beta \in (0, 1]$
- ▶ In essence, we would like to ‘quotient’ the topos  $[av]$  by an algebraic action – two questions:
  - ▶ Is the notion of (real) exponentiation geometric? Ng-Vickers (2021)
  - ▶ What does it mean to quotient by a monoid action vs. group action?



## Ostrowski's Theorem for $\mathbb{Q}$

Every absolute value of  $\mathbb{Q}$  is equivalent to a (non-Archimedean)  $p$ -adic absolute value  $|\cdot|_p$  (for some prime  $p$ ), or the Archimedean absolute value  $|\cdot|_\infty$ .

# Non-Archimedean Place (for fixed prime $p$ )



$$\begin{array}{ccccc} & \xrightarrow{\pi} & & & \\ [av_{NA}] \times (0, \infty) & \xleftarrow{s} & [av_{NA}] & \dashrightarrow & \mathcal{D} \\ & \xrightarrow{ex} & & & \end{array}$$

# Non-Archimedean Place (for fixed prime $p$ )



$$\begin{array}{ccccc} & \xrightarrow{\pi} & & & \\ [av_{NA}] \times (0, \infty) & \xleftarrow{s} & [av_{NA}] & \dashrightarrow & \mathcal{D} \\ & \xrightarrow{ex} & & & \end{array}$$

# Non-Archimedean Place (for fixed prime $p$ )



$$\begin{array}{ccc} & \xrightarrow{\pi} & \\ [av_{NA}] \times (0, \infty) & \xleftarrow{s} [av_{NA}] & \dashrightarrow \mathcal{D} \\ & \xrightarrow{ex} & \end{array}$$

- For any non-Arch. absolute  $|\cdot|$ , exponentiating  $|\cdot|^\alpha$  still yields a non-Arch. absolute value for any  $\alpha \in (0, \infty)$  (unlike the Archimedean case).

# Non-Archimedean Place (for fixed prime $p$ )



$$\begin{array}{ccc} & \xrightarrow{\pi} & \\ [av_{NA}] \times (0, \infty) & \xleftarrow{s} [av_{NA}] & \dashrightarrow \mathcal{D} \\ & \xrightarrow{ex} & \end{array}$$

- ▶ For any non-Arch. absolute  $|\cdot|$ , exponentiating  $|\cdot|^\alpha$  still yields a non-Arch. absolute value for any  $\alpha \in (0, \infty)$  (unlike the Archimedean case).
- ▶ What is  $\mathcal{D}$ ?



$$\begin{array}{ccc} & \xrightarrow{\pi} & \\ [av_{NA}] \times (0, \infty) & \xleftarrow{s} [av_{NA}] & \dashrightarrow \mathcal{D} \\ & \xrightarrow{ex} & \end{array}$$

- ▶ For any non-Arch. absolute  $|\cdot|$ , exponentiating  $|\cdot|^\alpha$  still yields a non-Arch. absolute value for any  $\alpha \in (0, \infty)$  (unlike the Archimedean case).
- ▶ What is  $\mathcal{D}$ ?

## Theorem

$$\mathcal{D} \simeq \text{Set}$$



$$\begin{array}{ccc} & \xrightarrow{\pi} & \\ [av_A] \times (0, 1] & \xleftarrow{s} [av_A] & \dashrightarrow \mathcal{D}' \\ & \xrightarrow{m} & \end{array}$$





$$\begin{array}{ccc} & \xrightarrow{\pi} & \\ [av_A] \times (0, 1] & \xleftarrow{s} [av_A] & \dashrightarrow \mathcal{D}' \\ & \xrightarrow{m} & \end{array}$$

- Space of Arch. absolute values is acted upon by a **monoid**  $(0, 1]$ -action as opposed to a **group**  $(0, \infty)$ -action.



$$\begin{array}{ccc} & \xrightarrow{\pi} & \\ [av_A] \times (0, 1] & \xleftarrow{s} [av_A] & \dashrightarrow \mathcal{D}' \\ & \xrightarrow{m} & \end{array}$$

- ▶ Space of Arch. absolute values is acted upon by a **monoid**  $(0, 1]$ -action as opposed to a **group**  $(0, \infty)$ -action.
- ▶ Can we play the same game as we did in the Non-Archimedean case?



$$\begin{array}{ccc} & \xrightarrow{\pi} & \\ [av_A] \times (0, 1] & \xleftarrow{s} [av_A] & \dashrightarrow \mathcal{D}' \\ & \xrightarrow{m} & \end{array}$$

- ▶ Space of Arch. absolute values is acted upon by a **monoid**  $(0, 1]$ -action as opposed to a **group**  $(0, \infty)$ -action.
- ▶ Can we play the same game as we did in the Non-Archimedean case? Answer: No!



$$\begin{array}{ccc} & \xrightarrow{\pi} & \\ [av_A] \times (0, 1] & \xleftarrow{s} [av_A] & \dashrightarrow \mathcal{D}' \\ & \xrightarrow{m} & \end{array}$$

- ▶ Space of Arch. absolute values is acted upon by a **monoid**  $(0, 1]$ -action as opposed to a **group**  $(0, \infty)$ -action.
- ▶ Can we play the same game as we did in the Non-Archimedean case? Answer: No! (The topos  $\mathcal{D}'$  has non-trivial forking in its sheaves)



$$\begin{array}{ccc} & \xrightarrow{\pi} & \\ [av_A] \times (0, 1] & \xleftarrow{s} [av_A] & \dashrightarrow \mathcal{D}' \\ & \xrightarrow{m} & \end{array}$$

- ▶ Space of Arch. absolute values is acted upon by a **monoid**  $(0, 1]$ -action as opposed to a **group**  $(0, \infty)$ -action.
- ▶ Can we play the same game as we did in the Non-Archimedean case? Answer: No! (The topos  $\mathcal{D}'$  has non-trivial forking in its sheaves)
- ▶ So what is  $\mathcal{D}'$ ?

# Preliminary Reorientations

Number Theory



23


## Candidate Picture

$$\mathcal{D}' \simeq \overleftarrow{[0, 1]}$$



## Candidate Picture

$\mathcal{D}' \simeq \overleftarrow{[0, 1]}$  (the space of 'upper reals' between 0 and 1)





## Candidate Picture

$\mathcal{D}' \simeq \overleftarrow{[0, 1]}$  (the space of 'upper reals' between 0 and 1)

- The Arakelov compactification of  $\mathrm{Spec}(\mathbb{Z})$  suggests that we add a single point at infinity to  $\mathrm{Spec}(\mathbb{Z})$  corresponding to the 'Archimedean prime' ...





## Candidate Picture

$\mathcal{D}' \simeq \overleftarrow{[0, 1]}$  (the space of 'upper reals' between 0 and 1)

- ▶ The Arakelov compactification of  $\mathrm{Spec}(\mathbb{Z})$  suggests that we add a single point at infinity to  $\mathrm{Spec}(\mathbb{Z})$  corresponding to the 'Archimedean prime' ... our candidate picture suggests that there is some blurring going on at infinity, and that infinity is not just a classical point with no intrinsic structure.



Sullivan's Arithmetic Square (a.k.a. 'The Hasse Square'):

$$\begin{array}{ccc} \mathbb{Z} & \longrightarrow & \prod_p \hat{\mathbb{Z}}_p \\ \downarrow & & \downarrow \\ \mathbb{Q} & \longrightarrow & \mathbb{Q} \otimes_{\mathbb{Z}} \prod_p \hat{\mathbb{Z}}_p = \mathbb{A}_{\mathbb{Q}}^f \end{array}$$

# By way of conclusion...



# By way of conclusion...



- ▶ Theme #1: Viewing toposes as a framework uniting logic and topology
- ▶ Theme #2: Local-Global issues, and its connections to Theme #1 via generic reasoning



- ▶ Theme #1: Viewing toposes as a framework uniting logic and topology
- ▶ Theme #2: Local-Global issues, and its connections to Theme #1 via generic reasoning
- ▶ Pulling away from the set theory reveals key insights into the deep nerve connecting topology and algebra.



- ▶ Theme #1: Viewing toposes as a framework uniting logic and topology
- ▶ Theme #2: Local-Global issues, and its connections to Theme #1 via generic reasoning
- ▶ Pulling away from the set theory reveals key insights into the deep nerve connecting topology and algebra.
- ▶ Some very interesting indications that there is some blurring at infinity in our picture of  $\overline{\text{Spec}(\mathbb{Z})}$  — interesting to explore the precise implications of this.



- [1] Balchin, S. and Greenlees, J.P.C. *Adelic models of tensor-triangulated categories*, Advances in Mathematics, 375:107339, (2020).
- [2] Boileau, A., Joyal, A. *La Logique des Topos*, The Journal of Symbolic Logic, Vol. 46, No. 1, pp. 6-16, (1981).
- [3] Caramello, O., *Theories, Sites, Toposes: Relating and studying mathematical theories through topos-theoretic 'bridges'*, Oxford University Press (2017).
- [4] Connes, A., Consani, C., *Absolute Algebra and Segal's  $\Gamma$ -rings*, Journal of Number Theory Volume 162, pp. 518-551, May 2016.
- [5] Egan, G., *Geometry and Waves [Extra]*,  
<http://www.gregegan.net/ORTHOGONAL/03/WavesExtra.html>



- [6] Hatcher, A.: *Vector Bundles and K-theory*,  
<https://www.math.cornell.edu/hatcher/VBKT/VBpage.html>
- [7] Johnstone, P.T.: *Topos Theory*, Dover Publications Inc., (1977)
- [8] Johnstone, P.T.: *Sketches of an Elephant: A Topos Theory Compendium, Vol. 1*, Clarendon Press, (2002)
- [9] Johnstone, P.T.: *Sketches of an Elephant: A Topos Theory Compendium, Vol. 2*, Clarendon Press, (2002)
- [10] Mazur, B., *On the Passage from Local to Global in Number Theory*, Bulletin of the AMS, 29 No. 1, (1993).
- [11] Moerdijk, I. *The classifying topos of a continuous groupoid, I.*, Transactions of the American Mathematical Society Volume 310, Number 2, pp. 629-668, 1988.





- [12] Nlab, *Vector Bundles*,  
<https://ncatlab.org/nlab/show/vector+bundle>.
- [13] Ng, M., and Vickers, S. *Point-free Construction of Real Exponentiation*, to appear in LMCS.
- [14] Sullivan, D., *Geometric Topology - Localization, Periodicity, and Galois Symmetry (The 1970 MIT notes)*,  
<https://www.maths.ed.ac.uk/~v1ranick/books/gtop.pdf>
- [15] Vickers, S., *Localic Completion of Generalized Metric Spaces*, preprint.
- [16] Vickers, S., *Continuity and Geometric Logic*, J. Applied Logic (12), pp. 14-27, (2014).
- [17] Zakharevich, I., *Attitudes of K-theory*, NOTICES OF THE AMERICAN MATHEMATICAL SOCIETY, Vol. 66, No. 7, pp. 1034-1044.