

Error propagation for TIMS Pb/Pb geochronology

Pieter Vermeesch
p.vermeesch@ucl.ac.uk

August 13, 2023

The Pb-budget of meteorites consists of extraterrestrial and terrestrial components. To separate these two components, we spike the sample with a solution that contains synthetic ^{202}Pb and ^{205}Pb . The spike serves two purposes. First, the ^{205}Pb in it is used as a tracer to compare the sample (which is a mixture of extraterrestrial and terrestrial Pb) with a blank solution (which contains only terrestrial Pb). Second, if the true $^{205}\text{Pb}/^{202}\text{Pb}$ -ratio of the spike is known, then this can be used to correct for mass-dependent fractionation. The extraterrestrial $^{204}\text{Pb}/^{206}\text{Pb}$ and $^{207}\text{Pb}/^{206}\text{Pb}$ -ratios of the sample can then be estimated as

$$\begin{cases} [4/6]_e = \frac{4_s - 4_b}{6_s - 6_b} \\ [7/6]_e = \frac{7_s - 7_b}{6_s - 6_b} \end{cases} \quad (1)$$

where x_s represents the amount (in pmol) of fractionation-corrected ^{20x}Pb in the spike-sample-blank mixture, and x_b is the corresponding amount in the spike-blank mixture:

$$x_* = M_* [x/5]_* F_*^{\beta(x)} \quad (2)$$

where M_* is the amount (in pmol) of ^{205}Pb added to the spike-sample-blank mixture (if $*$ = s) or spike-blank mixture (if $*$ = b); and $[x/5]_*$ is the corresponding $^{20x}\text{Pb}/^{205}\text{Pb}$ -ratio. When $M_s = M_b$, the spike amount cancels out in Equation 1. However, its uncertainty still needs to be included in the error propagation. $F_*^{\beta(x)}$ is the kinetic fractionation factor of Young et al. (2002), with

$$F_* = \frac{[2/5]_t}{[2/5]_*} \text{ and} \quad (3)$$

$$\beta(x) = \frac{\ln(205) - \ln(m_x)}{\ln(205) - \ln(202)} \quad (4)$$

in which $m_x = 20x$ is the molar mass of x , and $[2/5]_t$ is the true atomic $^{202}\text{Pb}/^{205}\text{Pb}$ -ratio of the spike.

Because isotopic ratios are strictly positive quantities whose uncertainties are best expressed in relative terms, it is useful to cast Equation 1 into a logarithmic form:

$$\begin{cases} l_{46} \equiv \ln[4/6]_e = l_4 - l_6 \\ l_{76} \equiv \ln[7/6]_e = l_7 - l_6 \end{cases} \quad (5)$$

where

$$l_4 = \ln(\exp[l_4(s)] - \exp[l_4(b)]) \quad (6)$$

$$l_6 = \ln(\exp[l_6(s)] - \exp[l_6(b)]) \quad (7)$$

$$l_7 = \ln(\exp[l_7(s)] - \exp[l_7(b)]) \quad (8)$$

with

$$l_x(*) = \ln M_* + \ln[x/5]_* + \beta(x) (\ln[2/5]_t - \ln[2/5]_*) \quad (9)$$

In order to solve Equation 5, we must first solve Equation 9 (for $x \in \{4, 6, 7\}$ and $* \in \{b, s\}$), and propagate its uncertainties. Let Σ be the covariance matrix of the blank-corrected $l_*(x)$ -measurements:

$$\Sigma = \begin{bmatrix} s[l_b(4)]^2 & s[l_b(4), l_b(6)] & s[l_b(4), l_b(7)] & s[l_b(4), l_s(4)] & s[l_b(4), l_s(6)] & s[l_b(4), l_s(7)] \\ s[l_b(6), l_b(4)] & s[l_b(6)]^2 & s[l_b(6), l_b(7)] & s[l_b(6), l_s(4)] & s[l_b(6), l_s(6)] & s[l_b(6), l_s(7)] \\ s[l_b(7), l_b(4)] & s[l_b(7), l_b(6)] & s[l_b(7)]^2 & s[l_b(7), l_s(4)] & s[l_b(7), l_s(6)] & s[l_b(7), l_s(7)] \\ s[l_s(4), l_b(4)] & s[l_s(4), l_b(6)] & s[l_s(4), l_b(7)] & s[l_s(4)]^2 & s[l_s(4), l_b(6)] & s[l_s(4), l_s(7)] \\ s[l_s(6), l_b(4)] & s[l_s(6), l_b(6)] & s[l_s(6), l_b(7)] & s[l_s(6), l_s(4)] & s[l_s(6)]^2 & s[l_s(6), l_s(7)] \\ s[l_s(7), l_b(4)] & s[l_s(7), l_b(6)] & s[l_s(7), l_b(7)] & s[l_s(7), l_s(4)] & s[l_s(7), l_s(6)] & s[l_s(7)]^2 \end{bmatrix} \quad (10)$$

where $s[x]^2$, $s[y]^2$ and $s[x, y]$ are the (co) variances of x and y ; then conventional error propagation by first order Taylor approximation dictates that:

$$\Sigma = \begin{bmatrix} I_3 & 0_{3,5} & J_t \\ 0_{5,3} & J_s & J_t \end{bmatrix} \begin{bmatrix} \Sigma_b & 0_{5,3} & 0_{5,1} \\ 0_{3,5} & \Sigma_s & 0_{4,1} \\ 0_{1,5} & 0_{1,4} & s[5/2]_t^2 \end{bmatrix} \begin{bmatrix} I_3 & 0_{3,5} \\ 0_{5,3} & J_s^T \\ J_t^T & J_t^T \end{bmatrix} \quad (11)$$

where I_n is the $n \times n$ identity matrix; $0_{n,m}$ is an $n \times m$ matrix of zeros; J_s and J_t are Jacobian matrices (with J_s^T and J_t^T their transpose):

$$J_s = \begin{bmatrix} \frac{1}{M_s} & \frac{-\beta(4)}{[2/5]_s} & \frac{1}{[4/5]_s} & 0 & 0 \\ \frac{1}{M_s} & \frac{-\beta(6)}{[2/5]_s} & 0 & \frac{1}{[6/5]_s} & 0 \\ \frac{1}{M_s} & \frac{-\beta(7)}{[2/5]_s} & 0 & 0 & \frac{1}{[7/5]_s} \end{bmatrix}; \quad J_t = \begin{bmatrix} \frac{\beta(4)}{[2/5]_t} \\ \frac{\beta(6)}{[2/5]_t} \\ \frac{\beta(7)}{[2/5]_t} \end{bmatrix} \quad (12)$$

and Σ_s and Σ_b are the covariance matrices of the sample and average blank, respectively. Σ_b is estimated from repeat measurements of

$$l'_b(x) = \ln M_b + \ln[x/5]_b - \beta(x) \ln[2/5]_b \quad (13)$$

(for $x \in \{4, 6, 7\}$) whereas Σ_s contains the analytical uncertainties of the raw measurements for the sample:

$$\Sigma_s = \begin{bmatrix} s[M_s]^2 & 0 & 0 & 0 & 0 \\ 0 & s[2/5]_s^2 & s([2/5]_s, [4/5]_s) & s([2/5]_s, [6/5]_s) & s([2/5]_s, [7/5]_s) \\ 0 & s([4/5]_s, [2/5]_s) & s[4/5]_s^2 & s([4/5]_s, [6/5]_s) & s([4/5]_s, [7/5]_s) \\ 0 & s([6/5]_s, [2/5]_s) & s([6/5]_s, [4/5]_s) & s[6/5]_s^2 & s([6/5]_s, [7/5]_s) \\ 0 & s([7/5]_s, [2/5]_s) & s([7/5]_s, [4/5]_s) & s([7/5]_s, [6/5]_s) & s[7/5]_s^2 \end{bmatrix} \quad (14)$$

Although it is possible to solve Equation 5 by plugging the $l_x(*)$ values from Equation 9 into Equations 6–7, this approach breaks down when the blank exceeds the signal. To rule out this possibility, it is better to estimate l_{46} and l_{76} using the method of maximum likelihood. Let Δ be a column vector (and Δ^T a row vector) of misfits:

$$\Delta = \begin{bmatrix} l_4(b) - c_{4b} \\ l_6(b) - c_{6b} \\ l_7(b) - c_{7b} \\ l_4(s) - \ln(\exp[c_{46} + c_6] + \exp[c_{4b}]) \\ l_6(s) - \ln(\exp[c_6] + \exp[c_{6b}]) \\ l_7(s) - \ln(\exp[c_{76} + c_6] + \exp[c_{7b}]) \end{bmatrix} \quad (15)$$

where c_{4b} , c_{6b} , c_{7b} , c_6 , c_{46} , c_{76} are the true (but unknown) values of $l_4(b)$, $l_6(b)$, $l_7(b)$, l_6 , l_{46} and $l_{76}(b)$, respectively. These parameters can be estimated by maximising the likelihood function (\mathcal{L}):

$$\mathcal{L} \propto -\frac{1}{2} \Delta^T \Sigma^{-1} \Delta \quad (16)$$

The uncertainties of c_{4b} , c_{6b} , c_{7b} , c_6 , c_{46} , c_{76} can be obtained by inverting the Hessian matrix of \mathcal{L} with respect to the parameters.

References

Young, E. D., Galy, A., and Nagahara, H. Kinetic and equilibrium mass-dependent isotope fractionation laws in nature and their geochemical and cosmochemical significance. *Geochimica et Cosmochimica Acta*, 66(6):1095–1104, 2002.