

Slides

If the two outcomes are *not* mutually exclusive:

$$P(\{2 \times H\} \cup \{3 \times H\}) = \frac{4}{8} + \frac{1}{8} - \frac{1}{8} = \frac{4}{8} = 0.5$$

$$P(\{2 \times H\} \cap \{1 \times T\}) + P(3 \times H)$$

- the mean of x is 9
- the variance of x is 11
- the mean of y is 7.50
- the variance of y is 4.125
- the correlation coefficient of x and y is 0.816
- the best fit line is given by $y = 3.00 + 0.500x$

urn:	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
10 samples selected with replacement:	5	2	12	19	10	5	3	19	11	4										
10 samples selected with replacement:	5	2	12	19	10	5	3	19	11	4										
10 samples selected without replacement:	15	8	13	5	4	19	7	1	20	10										

$$P(H, H, T) + P(\begin{smallmatrix} \square & \bullet \\ \bullet & \bullet \end{smallmatrix}, \begin{smallmatrix} \bullet & \bullet \\ \bullet & \bullet \end{smallmatrix}) - P(H, H, H \cap \begin{smallmatrix} \square & \bullet \\ \bullet & \bullet \end{smallmatrix}, \begin{smallmatrix} \bullet & \bullet \\ \bullet & \bullet \end{smallmatrix})$$

$$p = 2/3$$

We failed to reject H_0 for $\hat{p} = \frac{2}{5} = 0.4$. How about $\hat{p} = \frac{6}{15} = 0.4$?

$$k \in R$$

Next, how about $\hat{p} = \frac{12}{30} = 0.4$?

Which values of p are compatible with the observation of $k = 2$ successes out of $n = 5$ claims?

$$P(k \leq 2 | p = 2/3, n = 5) = \sum_{i=0}^2 \binom{5}{i} (2/3)^i (1/3)^{n-i} = 0.21 \geq \alpha/2$$

Assuming that H_0 is true, the probability of incurring a type-I error is:

1 experiment: $\alpha = 5\%$

2 experiments: $1 - (1 - \alpha)^2 = 9.75\%$

3 experiments: $1 - (1 - \alpha)^3 = 14\%$

48 experiments: $1 - (1 - \alpha)^{48} = 91.5\%$

N experiments: $1 - (1 - \alpha)^N\%$

Bonferroni correction: α/N

What is λ ?

the binomial distribution

the Poisson distribution

What is so normal about the Gaussian distribution?

What is $s[z]$?

$$s[d] = \sqrt{s[x]^2 + s[y]^2}$$

$$z = x + y$$

$$z = x - y$$

$$s[z]^2 = s[x]^2 + s[y]^2$$

$$z = xy$$

$$\left(\frac{s[z]}{z}\right)^2 = \left(\frac{s[x]}{x}\right)^2 + \left(\frac{s[y]}{y}\right)^2$$

$$s[y]^2 = y^2 \left(2\frac{s[t]}{t}\right)^2$$

$$2 \times 0.0000042 = 0.0000084 < \alpha$$

$$\mathrm{P}(\geq 9.0 \text{ earthquake})?$$

$$\frac{44 \ + \ 51 \ + \ 79 \ + \ 65 \ + \ 27 \ + \ 31 \ + \ 4 \ + \ 355 \ + \ 22 \ + \ 352 \ + \ 287 \ + \ 7 \ + \ 287 \ + \ 339 \ + \ 0}{276 \ + \ 342 \ + \ 355 \ + \ 334 \ + \ 296 \ + \ 7 \ + \ 17 \ + \ 351 \ + \ 349 \ + \ 37 \ + \ 339 \ + \ 40 \ + \ 324 \ + \ 325 \ + \ 334}$$

$$=189.2^{\circ}$$

$$\sin[\theta\pm\phi]=\sin[\theta]\cos[\phi]\pm\cos[\theta]\sin[\phi]$$

$$359^{\circ}-1^{\circ}=2^{\circ}$$

$$\bar{R}=0.88$$

$$s_c=0.51$$

$$s_c=2.04$$

x	0.85	0.84	0.83	0.8	0.77	0.75	0.74	0.73	0.71	0.6
y	0.53	0.54	0.56	0.6	0.64	0.66	0.67	0.68	0.71	0.8
x	-0.87	-0.85	-0.83	-0.8	-0.77	-0.75	-0.74	-0.73	-0.69	-0.64
y	-0.50	-0.53	-0.56	-0.6	-0.64	-0.66	-0.67	-0.68	-0.72	-0.77

Quizzes

Given $k = 9$ successes out of $n=10$ binomial trials, carry out a hypothesis test for

$$H_0 : p = 0.6$$

versus

$$H_a : p > 0.6$$

at a 95% confidence level.

X	0	1	2	3	4	5	6	7	8	9	10
$P(X)$	0.006	0.040	0.12	0.21	0.25	0.20	0.11	0.042	0.011	0.0016	0.00001
$P(x \leq X)$	0.006	0.046	0.17	0.38	0.63	0.83	0.95	0.990	1	1	1
$P(x \geq X)$	1	0.994	0.954	0.833	0.618	0.367	0.166	0.055	0.0123	0.0017	0.0001

Can the null hypothesis be rejected?

Given $k = 9$ successes out of $n=10$ binomial trials, carry out hypothesis test of

$$H_0 : p = 0.6$$

versus

$$H_a : p \neq 0.6$$

at a 95% confidence level.

X	0	1	2	3	4	5	6	7	8	9	10
$P(X)$	0.006	0.040	0.12	0.21	0.25	0.20	0.11	0.042	0.011	0.0016	0.00001
$P(x \leq X)$	0.006	0.046	0.17	0.38	0.63	0.83	0.95	0.990	1	1	1
$P(x \geq X)$	1	0.994	0.954	0.833	0.618	0.367	0.166	0.055	0.0123	0.0017	0.0001

Can the null hypothesis be rejected?

Given $k = 9$ successes out of $n=10$ trials, carry out a hypothesis test for

$$H_o : p = 0.6$$

versus

$$H_a : p < 0.6$$

at a 95% confidence level?

X	0	1	2	3	4	5	6	7	8	9	10
$P(X)$	0.006	0.040	0.12	0.21	0.25	0.20	0.11	0.042	0.011	0.0016	0.00001
$P(x \leq X)$	0.006	0.046	0.17	0.38	0.63	0.83	0.95	0.990	1	1	1
$P(x \geq X)$	1	0.994	0.954	0.833	0.618	0.367	0.166	0.055	0.0123	0.0017	0.0001

Can the null hypothesis be rejected?