

# FAIR fission track analysis with geochron@home

Pieter Vermeesch<sup>1</sup>, Tim Band<sup>1</sup>, Jiangping He<sup>2</sup>, Andrew Carter<sup>3</sup>, and Rex Galbraith<sup>1</sup>

<sup>1</sup>University College London, London WC1E 6BT, United Kingdom

<sup>2</sup>King's College London, London SE1 8WA, United Kingdom

<sup>3</sup>Birkbeck, University of London, London WC1E 7HX, United Kingdom

**Correspondence:** Pieter Vermeesch (p.vermeesch@ucl.ac.uk)

**Abstract.** Fission track geochronology is based on the visual analysis of optical images. This visual process is prone to observer bias. Fission track datasets are currently reported as small data tables. The interpretation of these tables requires a high degree of trust between the fission track analyst and the user of the data. geochron@home is software that removes this requirement of trust. It combines a browser-based ‘virtual microscope’ with an online database to provide FAIR (Findable, Accessible,

5 Interoperable and Reproducible) access to fission track data.

geochron@home serves four different purposes. It can be used (1) to count fission tracks in ‘private mode’, i.e. hidden from other users on the internet; (2) to archive fission track images and counts for inspection by other users; (3) to create tutorials for new students of the fission track method; and (4) to generate crowd-sourced fission track data, by serving randomly selected selections of images to citizen scientists. We illustrate these four applications with examples that demonstrate (1)

10 geochron@home’s ability to compare and combine fission track counts for multiple users within a lab group; (2) the use of the geochron@home archive in the peer review system; (3) the effectiveness of even simple tutorials in teaching novice users how to count fission tracks; and (4) the ‘wisdom of crowds’ in fission track identification.

geochron@home was written in Python and PostgreSQL. Its code is freely available for inspection and modification, allowing users to set up their own geochron@home server. Alternatively, users who would like to upload data to the GaHa, but do not 15 have the facilities to set up their own server, may use the server at University College London free of charge. The GaHa accepts image stacks acquired on any type of digital microscope, and accommodates fission track data (counts and length measurements) from external fission track analysis suites such as Fission Track Studio and TrackFlow.

We anticipate that the introduction of FAIR workflows will make fission track data more accurate and more future proof. Storing fission track data online will benefit future developments in fission track thermochronology. For example, archival 20 datasets of peer reviewed fission track counts can be used to train and improve machine learning algorithms for automated fission track analysis. We hope that other geochronological methods will follow the fission track community’s lead in FAIR data processing, with benefits to all the other Earth Science disciplines that depend on geochronological data.

## 1 Introduction

Science is on an irreversible trajectory towards greater openness. Geochronology is no exception to this trend, as this journal  
25 illustrates with its open access and review policies. Funding agencies increasingly demand that research results and data are shared with the public. Currently, geochronological data are generally provided as flat tables of dates or isotopic ratio estimates. However, in other fields of science such as physics, it is common practice to share the raw unprocessed measurements along with processing instructions (e.g., Abbott et al., 2016). This paper moves geochronology in the same direction. It presents a mechanism to generate and store fully FAIR (Findable, Accessible, Interoperable and Reproducible; Wilkinson et al., 2016)  
30 data in the context of fission track analysis.

Unlike most other geochronometers, which use mass spectrometers to estimate parent-daughter ratios, fission tracks are observed under an optical microscope and counted by a human observer. In recent years, digital microscopy has moved fission track data acquisition from the objective lens of a microscope to the computer screen (Gleadow et al., 2009; Van Ranst et al., 2019; Gleadow, 2019). Ongoing developments in artificial intelligence generate further opportunities to improve the throughput  
35 and accuracy of fission track data (Nachtergael and De Grave, 2020). But despite the richness of the digital datasets produced by these novel tools, fission track data are still reported as small summary tables. These tables require an unnecessary degree of trust between the ‘producer’ and ‘consumer’ of the data.

geochron@home is a free and open software platform that allows geochronologists to share their raw fission track data over the internet for perusal by peer reviewers, colleagues and the general public. geochron@home is a virtual petrographic  
40 microscope connected to a database with digital image stacks of etched fission track samples. The platform can be used to acquire, archive and inspect fission track data in full adherence to the FAIR data principles. In Section 2, we will describe geochron@home’s software architecture in five steps. We will show that this architecture accommodates imagery from any type of digital microscope. It enables flexible workflows that can be adapted to four different applications.

Using image stacks of Mount Dromedary apatite, we will show how geochron@home can be used to count fission tracks  
45 in ‘private mode’ (Section 3); to archive published fission track datasets in ‘public mode’ (Section 4); to build tutorials for training purposes (Section 5); and to crowd-source fission track data on the internet (Section 6). Because geochron@home is free and open, it can be extended and improved by any interested party. We make some suggestions for future improvements in Section 7. We hope that the geochron@home’s example will be followed in other geochronological disciplines, as this will benefit not only geochronology itself, but all the other disciplines that depend on it (Section 8).


## 50 2 Workflow

The geochron@home workflow separates the acquisition of microscope images from their analysis, providing the flexibility to accommodate data from different microscope manufacturers, which typically come with their own instrument control software.  
The workflow can be broken down into five steps.

1. Acquisition of z-stacks of microscope images in reflected and transmitted light for each of the grains in a sample and,  
55     optionally, for the accompanying external detector (Figure 1). At University College London, this first step is currently  
   accomplished by a Python macro within Zeiss' Zen Blue software. However, geochron@home can also accommodate imagery from other platforms, such as Fission Track Studio (Zeiss; Gleadow et al., 2009) and TrackFlow (Nikon;  
   Van Ranst et al., 2019).
2. Prepare the z-stacks for uploading to the geochron@home database. This database requires that the images are organised  
60     as a nested sequence of directories, in which a 'project' consists of 'samples' that comprise a number of 'grains'. Each  
   grain corresponds to a numbered sequence of .jpeg images, one for each layer of the z-stack. Note that the raw  
   microscope images are generally not stored as .jpeg files but in uncompressed .czi (for Zen Blue), .tif (for Fission  
   Track Studio) or .nd2 (for Nikon/TrackFlow) formats. Conversion from these raw images to sequences of .jpeg files  
   is done with a shell script using ImageMagick (Still, 2006).
- 65     In addition to the sequence of .jpeg images, a low level 'grain' folder can also include an optional file called `roi.json`  
   containing the vertices of a default region of interest for spontaneous (and/or induced) track counting. The database struc-  
   ture can also accommodate existing fission track counting results. For example, if a user has already counted their fission  
   tracks in Fission Track Studio, then they can store those results in a `.json` file at the 'sample' level directory. Because  
70     Fission Track Studio stores its results in an `.xml` format, a second conversion script is needed to translate those results  
   into an equivalent `.json` format.
3. Upload the data to the geochron@home platform. geochron@home is a Django web-app with a PostgreSQL database.  
   The database can be accessed via a Python API and a (more limited) web-based GUI interface. Accessing the API  
   requires administrator privileges. Administrators can create projects, samples and grains; download data; and set the  
75     access rights of 'ordinary' users. Projects can be private or public. The source code and installation instructions for  
   geochron@home are freely available over GitHub (see the Data Availability statement at the end of this paper). This  
   allows fission track users to set up their own server. Alternatively, fission track labs can upload their data to the UCL  
   server by contacting the corresponding author.
4. Analyse the images with a browser-based 'virtual fission track microscope'. This virtual instrument has a simple user  
   interface with controls to zoom, pan and focus in or out of the digital image stack. Depending on the permissions granted  
80     to the user by the administrator, the virtual microscope offers a number of different options. Entry level 'ordinary'  
   users are only allowed to count tracks by clicking within a pre-defined 'region of interest'. In contrast, 'superusers'  
   are allowed to define their own regions of interest. Once the user is satisfied that they have counted all the fission tracks in  
   a particular grain, they can submit the results to the server. They are then presented with a new set of images until all  
   grains are counted.

5. Post-processing. The fission track data can either be downloaded as a flat data table of counts and area, or as a `.json` file containing the locations of all the coounted tracks. `geochron@home` does not provide any tools to post-process these files. They are meant to be passed on to other tools such as spreadsheet applications or `IsoplotR` (Vermeesch, 2018).

The five-step workflow can be used for several applications, including (1) conventional fission track analysis; (2) archiving published fission track results; (3) building tutorials; and (4) crowd-sourcing fission track data. The next sections of this paper  
90 will illustrate these four different applications with real world examples.



**Figure 1.** Screenshots of raw fission track data in `geochron@home`. GR: an apatite grain in reflected light; GT: the same grain in transmitted light; MR: the corresponding mica detector in reflected light; MT: the mica detector in transmitted light. White rectangles mark the region of interest (ROI), within which an analyst has counted fission tracks by marking their etch pits (shown in yellow). The raw data for this figure can be viewed on the `geochron@home` archive (<https://github.com/pvermees/GaHa>).

### 3 Counting fission tracks in ‘private mode’

Administrators can define regions of interest (ROI) and count or edit the fission track coordinates of any grain in a `geochron@home` database. They can also assign other users to groups, and give these groups access to a subset of the projects in the database.  
95 Administrators have fine control over the permissions of the groups. For example, they can allow the members of a group to count fission tracks in the predefined ROIs, or to define their own ROIs. Groups provide a mechanism to compare and combine the results of multiple analysts of the same sample. Figure 2 illustrates this with two sets of fission track density estimates (in tracks per  $\mu\text{m}^2$ ) for the same sample of Mount Dromedary apatite (Green, 1985), analysed by two users (PV and AC).

The two analysts defined their own ROIs, resulting in slightly different density estimates (Figure 2a). Figure 2b compares the track density estimates of the two analysts for 25apatite grains on a radial plot, using the assumption that the density estimates are statistically independent from each other. Let  $N_{PV}$  and  $N_{AC}$  be the number of spontaneous tracks counted by PV and AC over areas  $A_{PV}$  and  $A_{AC}$ , respectively. Then Figure 2b assumes that the ratio of the estimated track densities  $\hat{\rho}_{PV} = N_{PV}/A_{PV}$  and  $\hat{\rho}_{AC} = N_{AC}/A_{AC}$  is associated with the following uncertainty:

$$\hat{\sigma} = \sqrt{\text{var}\left(\frac{\hat{\rho}_{PV}}{\hat{\rho}_{AC}}\right)} = \frac{\hat{\rho}_{PV}}{\hat{\rho}_{AC}} \sqrt{\frac{1}{N_{PV}} + \frac{1}{N_{PV}}} \quad (1)$$


In reality, the uncertainties are strongly correlated due to the significant overlap between the counting areas chosen by the two analysts. This overlap causes the density ratios to be underdispersed with respect to the assigned uncertainties (MSWD = 0.34 in Figure 2b). Equation 1 can be modified to remove the error correlation under the assumption that the two analysts counted exactly the same features in the overlapping area ( $A_0$ ):

$$\hat{\sigma}' = \frac{\hat{\rho}_{PV}}{\hat{\rho}_{AC}} \sqrt{\frac{A_{PV} + A_{AC}}{N_{PV} + N_{AC}}} \sqrt{\frac{1}{A_{PV}} + \frac{1}{A_{AC}} - \frac{2A_0}{A_{PV}A_{AC}}} \quad (2)$$

If there is no overlap between the two counting areas (i.e., if  $A_0 = 0$ ), then  $\hat{\sigma}' \approx \hat{\sigma}$ . Conversely, if the counting areas are identical (i.e., if  $A_{PV} = A_{AC} = A_0$ ), and if the two analysts produce perfectly consistent counts, then  $\hat{\sigma}' = 0$ .

Equation 2 produces a radial plot (Figure 2c) that is overdispersed (MSWD = 4.2) with respect to  $\hat{\sigma}'$ . This suggests that the assumption of perfect consistency is also wrong. Detailed inspection of the counted features in the virtual microscope reveals that the two analysts are in perfect agreement for long fission tracks, but that they occasionally disagree about the shorter features. The ratio of the pooled track densities indicates that PV undercounts the Mount Dromedary apatite by 7% relative to AC.

In summary, it could be said that Equation 1 represents an overly ‘pessimistic’ view of analysts’ abilities to reproduce each other’s results, whereas Equation 2 is too ‘optimistic’. The truth lies between the two scenarios. The existence of ‘observer bias’ is well documented (Tamer et al., 2025). It is one of the reasons why fission track analysis is often done relative to age standards: observer bias does not have to be a problem provided that it is consistent between grains, and between samples.



**Figure 2.** Comparison of fission track data for Mount Dromedary apatite by two analysts (PV = Pieter Vermeesch and AC = Andrew Carter). a) the track counts and ROIs of both users for grain 4648, shown in blue (PV) and red (AC); b) radial plot for the ratio of the track densities of both users for all 25 grains analysed by the two analysts, using the ‘pessimistic’ scenario of Equation 1; c) radial plot of the same data, using the ‘optimistic’ error propagation formula of Equation 2.

#### 4 The geochron@home archive (GaHa)

Once a set of fission track images has been analysed and the analyst is confident that the results are accurate, the status of the results can be changed from private to public. This makes the results visible over the internet as a list of URLs, where each grain number and user ID corresponds to a unique address. The geochron@home archive (GaHa) brings fission track geochronology into the era of FAIR science. It allows peer reviewers to inspect the raw data from which thermochronological inferences are made. The archive is open to submissions from any fission track laboratory, free of charge. However, as mentioned in Section 2, it is also possible to establish a new archive elsewhere. At the time of writing, GaHa contains data for three studies:


1. Guo et al. (2025): this is an LA-ICP-MS based fission track study of detrital apatite from the northeastern Tibetan Plateau.  
130 It contains image stacks of semitracks in 1146 apatite grains from 16 different samples.
2. Tamer et al. (2025): this is a round-robin study in which digital image stacks of 44 apatite crystals were circulated among 14 different analysts. These analysts used the FastTrack image analysis software (which is part of the Fission Track Studio suite) to define their own ROIs and count the semitracks and horizontally confined fission tracks in them. GaHa presents the results of the round-robin experiment as a  $44 \times 14$  grid of URLs.
3. This study: All the raw fission track data used in this article are available on GaHa, along with the post-processing software that was used to produce the figures. Together, these resources provide the reader with all the information needed to fully reproduce our results, ‘from cradle to grave’. To our knowledge, this is the first geochronological study to do so.

#### 5 geochron@home tutorials

- 140 Given the right permissions (assigned by an administrator), users can build tutorial pages by annotating features in fission track images. These features can be tracks or other objects such as scratches, inclusions, dislocations or holes. A selection of tutorial pages is presented to new users when they first log into geochron@home. They must complete the tutorial before being allowed to count fission tracks. The tutorial pages can be revisited at any time by visiting the corresponding link on the geochron@home landing page.
- 145 The annotations in the current tutorial pages were made by an experienced fission track analyst (Andrew Carter). They provide a quick and easy mechanism to train novice users in the art of fission track analysis. At the time of writing, the tutorial pages offer only a limited degree of interactivity. Users can click on features to read the annotations. However, future plans

include the addition of fully interactive ‘quizzes’ (see Section 8). Despite the simplicity of the current tutorials, they are already quite effective, as demonstrated by the crowd-sourcing experiment described in the next section of this paper.

## 150 6 Crowd-sourcing fission track data



**Figure 3.** a) box plots of 25 crowd-sourced sets of fission track counts for 70 analysts. b) radial plot comparing the paired track counts for the first two grains in the dataset (4649 and 4673). The estimate of user PV is shown in blue. Analyst AC used a different ROI and is not shown.

In 1906, Sir Francis Galton visited a county fair in which a contest was held to guess the weight of an ox. 787 people participated in the event. Galton discovered that the median of all their estimates was within 0.8% of the true weight of the ox and more accurate than 90% of the individual estimates. Such is the ‘wisdom of crowds’ (Galton, 1907). Similar effects are seen in fission track geochronology.

A 1984 interlaboratory comparison study (Miller et al., 1985) showed that the average of several fission track age estimates is closer to the known age of mineral standards than the age obtained by any individual observer. Unfortunately, routine measurement of fission track samples by multiple analysts is prohibitively expensive in a normal laboratory environment. geochron@home changes this, by bringing fission track dating to a proverbial ‘county fair’ of citizen scientists.

160 Figure 4 presents the results of a crowd-sourcing experiment that was carried out at UCL as part of an undergraduate course in isotope geology. 70 students were asked to create an account on the geochron@home server by selecting a unique username and password. After spending a few minutes to complete the compulsory tutorial (Section 5), the students were asked to count fission tracks in randomly assigned image stacks of Mount Dromedary apatite. Each student was required to analyse at least 15 grains, using pre-defined ROIs. In a matter of hours, the students amassed a dataset of 37,331 fission track counts in 34 grains.


165 This large dataset was parsed into separate data files –one for each student– which they processed during an assessed practical exercise.

The raw counts are shown as a series of box plots in Figure 4a for the same selection of 25 grains that were analysed by AC and PV in Section 3. The box plots exhibit significant variability between the grains and between the students. This inter-grain variability is expected. It reflects the different sizes of the ROIs, and the variable U-content of the grains. The variability of the  
170 track counts for each grain results from observer bias. This phenomenon was previously discussed in Section 3. However, the bias is much stronger for the crowd-sourced dataset than it was for the two ‘expert’ analysts of Section 3.

Figure 4b compares the 42 raw fission track counts of the first two grains on a radial plot. The pairwise ratios of the counts for each analyst are more consistent than the raw counts of Figure 4a. Students who over- or undercount the first grain, tend to over- or undercount the second grain as well. Nevertheless, the results are overdispersed with respect to the counting uncertainties,  
175 even when using the ‘pessimistic’ scenario of Equation 1. The poor consistency of the results reflects the very limited training received by the students. But despite these inconsistencies, the opinion of an ‘expert’ (PV, shown in blue) falls the middle of the estimates obtained by the citizen scientists. This suggests that the collective wisdom of the entire group of students is greater than that of any individual analyst.

As mentioned in Section 2, geochron@home stores the actual track positions marked by the users. This raw data can be  
180 downloaded as a .json file and inspected in detail. Figure 3 is a two-dimensional histogram for the x-y positions of all the track positions generated by the students in the two apatite crystals. Visual comparison of the histogram with the optical image confirms that the students unanimously identified the most obvious semitracks, which contain a clearly visible etch pit and tail. Shorter and fainter tracks received fewer clicks. Datasets like this can be used to replace integer counts of fission tracks with probabilities, reflecting the ambiguity of some fission track datasets.

185 Figure 3 also shows that some students counted the tails of the fission tracks rather than their etch pits, despite being told the opposite in the tutorial. Fixing this issue will require some improvements to the tutorial pages (Section 7).



**Figure 4.** a,b) Optical images in transmitted light of Mount Dromedary apatites 4649 and 4673 in the crowd-sourcing experiment. c,d) Two-dimensional histogram of the track locations for the two grains, as identified by the citizen scientists.

## 7 Outlook

geochron@home has been in development for 10 years and remains a work in progress. Planned improvements include:

- 190 1. Interactive tutorial pages. To ensure that novice users do not count the tails but the etch pits of fission tracks, we will add a ‘quiz’ to the tutorial pages. Only users who click on the ‘correct’ features in an unlabelled set of images will be allowed to count new samples.
2. Length measurements. The geochron@home archive already contains horizontally confined fission track measurements (Tamer et al., 2025). However, these results must be generated externally (e.g., using Fission Track Studio) and uploaded via a .json file. The virtual microscope currently lacks the functionality to generate length data within geochron@home. This functionality will be added in a future update.
- 195 3. Dpar and Dper. Etch pits are currently stored as simple sets of x- and y-coordinates. In reality, etch pits have a finite length (‘Dpar’) and width (‘Dper’), which serve as useful indicators for the horizontal etch rates along the c-axis and parallel to it (Donelick, 1993). Functions will be added measure and visualise these type of data in geochron@home.
- 200 4. Machine learning. Data science is experiencing an artificial intelligence revolution that has already started to transform the fission track method (Nachtergaele and De Grave, 2020). Convolutional neural networks must be trained with example data. geochron@home is ideally suited for this task. Section 6 showed how the collective wisdom of multiple fission

track analysts can label fission track images with probabilities rather than counts. This data format is close to the form in which data are treated within an AI algorithm.

205 Once trained on historical data, AI algorithms can be used to count fission tracks automatically. Following the model of Fission Track Studio, machine learning can be used to reverse the fission track counting process. Instead of asking users to count the fission tracks in a sample, the software can ask them to check the results proposed by an AI algorithm, and to remove any features that are *not* fission tracks.

210 Regardless of whether fission tracks were counted manually or with a machine, the value of the geochron@home archive remains the same. It is important to document data so that samples can be reanalysed in the future, for example when a new and improved generation of machine learning algorithms becomes available.

These four improvements will be made by ourselves pending additional funding. However, because geochron@home is free and open software, we invite any interested parties to join the effort and extend or improve our code.

## 8 Conclusions

215 This paper introduced geochron@home, a software platform for FAIR fission track analysis. We demonstrated four different applications of this platform using real data. Putting the FAIR data paradigm into practice, all the imagery, counts and source code for this paper are publicly available. Using these resources, the reader can reproduce all the results that were presented in this publication. We hope that other geochronologists will follow this example. FAIR data promises to address the reproducibility crisis in science (Miyakawa, 2020).

220 geochron@home's rich archive of raw data can be reanalysed in the future. We anticipate that the adoption of FAIR data processing workflows will open up new research opportunities. For example, archived pairs of peer-reviewed fission track images and counts could be used to train the next generation of automated machine learning algorithms. Conversely, it is also possible that future improvements in fission track images analysis will be used to update the count data for published datasets, improved their accuracy.

225 Another advantage of the geochron@home wokflow is the separation of image acquisition and image analysis. This separation reduces the hardware requirements for fission track dating. It opens up the possibility to share resources. State-of-the-art digital microscopes are expensive. Using geochron@home, a single microscope can serve multiple users and make fission track analysis more affordable.

230 The fission track method has always been a test bed for new geochronological developments. Because fission track data are imprecise, the fission track community has sollicited the help of statisticians and mathematicians to develop its analytical protocols. Other geochronological communities are still catching up with concepts and tools such as overdispersion, mixture modelling and radial plots, which have been commonplace in fission track dating for decades (Vermeesch, 2019). In a similar

vein, the subjective nature of fission track identification has prompted the fission track community to organise inter-laboratory comparisons and round-robin studies long before other geochronological communities (Miller et al., 1985; Tamer et al., 2025).

235 With the development of geochron@home, fission tracks is once again ahead of the pack in terms of FAIR data analysis. geochron@home currently only stores images and counts. This is enough to reproduce the results of fission track studies using the external detector method, but not for LA-ICP-MS based data. FAIR data processing of LA-ICP-MS data requires a new generation of mass spectrometer data reduction software. We are currently working on this. The development of FAIR ICP-MS data pipelines will not only benefit fission track dating but other chronometers as well, such as in-situ U-Pb, Rb-Sr and Lu-Hf  
240 (Vermeesch et al., 2025).

With the establishment of FAIR data, the era of ‘fiction tracks’ will be gone for good, and geochronology will be well placed to avoid the reproducibility problems that have plagued other fields of science.

245 *Code and data availability.* geochron@home is free software released under the GPL-3 license. The package and its source code are available from <https://github.com/pvermees/geochron-at-home> (last access: April 2, 2025). The raw data (imagery) are available at the geochron@home archive (<https://github.com/pvermees/GaHa>, last access: April 3, 2025). R-scripts to reproduce the figures are provided in the supplementary information (<https://github.com/pvermees/supplements>, last access: April 3, 2025).

*Author contributions.* Pieter Vermeesch designed the study, acquired the funding, counted fission tracks and wrote the paper. Jiangping He created geochron@home. Tim Band expanded geochron@home and wrote the accompanying microscope image acquisition software. Andrew Carter provided the samples, prepared the training data and counted fission tracks. Rex Galbraith derived Equation 2.

250 *Competing interests.* PV is an Associate Editor of *Geochronology*.

*Acknowledgements.* This research has been supported by the Natural Environment Research Council (grant no. NE/T001518/1), awarded to Pieter Vermeesch. We would like to thank the students of GEOL0017 (‘Isotope Geology’) for their contribution to the crowd-sourcing experiment of Section 6.

## References

- 255 Abbott, B. P., Abbott, R., Abbott, T. D., Abernathy, M. R., Acernese, F., Ackley, K., Adams, C., Adams, T., Addesso, P., Adhikari, R. X., et al.: Observation of gravitational waves from a binary black hole merger, *Physical Review Letters*, 116, 061102, 2016.
- Donelick, R. A.: Method of fission track analysis utilizing bulk chemical etching of apatite, uS Patent 5,267,274, 1993.
- Galton, F.: Vox Populi, *Nature*, 75, 450–451, 1907.
- Gleadow, A.: Future developments in fission track thermochronology, in: *Fission track thermochronology and its application to geology*, 260 edited by Malusà, Marco and Fitzgerald, P., chap. 4, Springer, 2019.
- Gleadow, A. J., Gleadow, S. J., Belton, D. X., Kohn, B. P., Krochmal, M. S., and Brown, R. W.: Coincidence mapping – a key strategy for the automatic counting of fission tracks in natural minerals, *Geological Society, London, Special Publications*, 324, 25–36, 2009.
- Green, P.: Comparison of zeta calibration baselines for fission-track dating of apatite, zircon and sphene, *Chemical Geology: Isotope Geo-science Section*, 58, 1–22, 1985.
- 265 Guo, C., Zhang, Z., Lease, R., Malusà, M. G., Chew, D., Lu, H., Wu, L., Xiang, D., Wang, N., Grasemann, B., et al.: Apatite geo-thermochronology and geochemistry constrain oligocene-miocene growth and geodynamics of the northeastern Tibetan Plateau, *Geo-physical Research Letters*, 52, e2024GL113157, 2025.
- Miller, D. S., Duddy, I. R., Green, P. F., Hurford, A. J., and Naeser, C. W.: Results of interlaboratory comparison of fission-track age standards: fission-track workshop – 1984, *Nuclear Tracks and Radiation Measurements*, 10, 383–391, 1985.
- 270 Miyakawa, T.: No raw data, no science: another possible source of the reproducibility crisis, 2020.
- Nachtergaelie, S. and De Grave, J.: AI-Track-tive: automated fission track recognition using computer vision (Artificial Intelligence), *Geochronology Discussions*, 2020, 1–12, 2020.
- Still, M.: *The Definitive Guide to ImageMagick*, Definitive Guide Series, Springer, 1st edn., ISBN 978-1-59059-590-9, <https://www.apress.com/gp/book/9781590595909>, a comprehensive guide to ImageMagick, covering installation, usage, and scripting., 2006.
- 275 Tamer, M. T., Chung, L., Ketcham, R. A., and Gleadow, A. J. W.: The need for fission-track data transparency and sharing, *Geochronology*, 7, 45–58, <https://doi.org/10.5194/gchron-7-45-2025>, 2025.
- Van Ranst, G., Baert, P., Fernandes, A. C., and De Grave, J.: Technical note: TRACKFlow, a new versatile microscope system for fission track analysis, *Geochronology Discussions*, 2019, 1–25, <https://doi.org/10.5194/gchron-2019-13>, 2019.
- Vermeesch, P.: IsoplotR: a free and open toolbox for geochronology, *Geoscience Frontiers*, 9, 1479–1493, 2018.
- 280 Vermeesch, P.: Statistics for fission-track thermochronology, *Fission-track thermochronology and its application to geology*, pp. 109–122, 2019.
- Vermeesch, P., Hopley, P., Roberts, N., and Parrish, R.: Geochronology of Taung and other southern African australopiths, in: One hundred years of *Australopithecus africanus*, edited by Wood, B. A., Grine, F. E., and Smith, H. B., chap. 4, Springer, 2025.
- Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-W., da Silva Santos, L. B., 285 Bourne, P. E., et al.: The FAIR Guiding Principles for scientific data management and stewardship, *Scientific data*, 3, 1–9, 2016.