



DEPARTMENT OF MATHEMATICS

| | | |
|--|--|---|
| Course: Fundamentals of Linear Algebra, Calculus and Statistics | CIE-I (QUIZ & TEST) | Maximum marks: 10+50=60 |
| Course code: 22MA11C | First semester 2022-2023 Chemistry Cycle Branch: AI, BT, CD, CS, CY, IS, SPARK-C | Time: 9.15am to 11.15am Date: 17-01-2023 |

Instructions to candidates:

- Part A must be answered within the first two pages of the Booklet.
- Answer all questions.

| Q.No | PART- A | M | BT | CO |
|------|---|---|----|----|
| 1.1 | The rank of the Matrix $A = \begin{bmatrix} 1 & 1 & 1 \\ a & b & c \\ a^2 & b^2 & c^2 \end{bmatrix}$ for $a = b \neq c$ is _____. | 2 | 2 | 1 |
| 1.2 | The value of p for which the following set of equations will have no solution is _____. $2x + 3y = 5$ $3x + py = 10$ | 2 | 2 | 1 |
| 1.3 | If $A = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$, then eigenvalues of A^{-1} are _____. | 2 | 2 | 2 |
| 1.4 | The transformation of the Lemniscate $r^2 = 2 \cos 2\theta$ in cartesian system is _____. | 2 | 1 | 2 |
| 1.5 | If $x = at^2$ and $y = 2at$, then the radius of curvature ρ for the given curve is _____. | 2 | 2 | 2 |

| Q.No | PART- B | M | BT | CO |
|-------|---|----|----|----|
| 1 | Test the consistency of the following system $3.0x_1 + 2.0x_2 + 2.0x_3 - 5.0x_4 = 8.0$ $0.6x_1 + 1.5x_2 + 1.5x_3 - 5.4x_4 = 2.7$ $1.2x_1 - 0.3x_2 - 0.3x_3 + 2.4x_4 = 2.1$. Solve if the system is consistent. | 10 | 3 | 1 |
| 2 | The currents i_1, i_2, i_3 in the paths of an electrical network follow the linear equations $i_1 - i_2 + i_3 = 0, 3i_1 + 2i_2 = 7, 2i_2 + 4i_3 = 8$. Determine i_1, i_2, i_3 using Gauss-Jordan elimination method. | 10 | 3 | 4 |
| 3 | Employ the Rayleigh's Power method to estimate the dominant eigenvalue and its associated eigen vector for the matrix $A = \begin{bmatrix} 3 & 2 & 3 \\ 2 & 6 & 6 \\ 3 & 6 & 3 \end{bmatrix}$ by taking $[1 \ 1 \ 1]^T$ as initial eigenvector. Perform 6 iterations. (Consider 4 decimal places) | 10 | 2 | 2 |
| 4 (a) | The temperature u_1, u_2, u_3 of a metal plate under some circumstances is given by: $u_1 - 8u_2 + 3u_3 + 4 = 0,$ $2u_1 + u_2 + 9u_3 = 12,$ $8u_1 + 2u_2 - 2u_3 = 8:$ Solve for the temperatures using Gauss- Seidel iterative method. Carry out 3 iterations. (Consider 4 decimal places) | 6 | 3 | 3 |
| 4(b) | Show that the curves $r = a\theta$ and $r = \frac{a}{\theta}$ intersect orthogonally. | 4 | 2 | 2 |
| 5 | Find the circle of curvature of $b^2x^2 + a^2y^2 = a^2b^2$ at a point of its intersection with the y-axis. | 10 | 2 | 3 |

BT-Blooms Taxonomy, CO-Course Outcomes, M-Marks

| Marks Distribution | Particulars | CO1 | CO2 | CO3 | CO4 | L1 | L2 | L3 | L4 | L5 | L6 |
|--------------------|-------------|-----|-----|-----|-----|----|----|----|----|----|----|
| | Test | 14 | 20 | 16 | 10 | 2 | 32 | 26 | - | - | - |
| | Max Marks | | | | | | | | | | |



RV Educational Institutions
RV College of Engineering

Autonomous
Institution Affiliated
to Visvesvaraya
Technological
University, Belagavi

Approved by AICTE,
New Delhi, Accredited
By NAAC, Bengaluru
And NBA, New Delhi

Go, change the world

DEPARTMENT OF MATHEMATICS

| | | |
|---|--|---|
| Course: Fundamentals of Linear Algebra, Calculus and statistics | CIE-II (QUIZ & TEST) | Maximum marks: 10+50=60 |
| Course code: 22MA11C | First semester 2022-2023 Chemistry Cycle Branch: AI, BT, CD, CS, CY, IS, SPARK-C | Time: 1:00PM-3:00PM Date: 21-02-2023 |

Instructions to students:

1. Answer all questions.
2. Part A must be answered in the first two pages of the answer book only.

| Q.No | PART- A (Quiz) | M | BT | CO |
|------|---|---|----|----|
| 1.1 | The coefficient of x^2 in the Maclaurin series expansion of $e^{-\left(\frac{x}{3}\right)}$ is _____. | 2 | L1 | 1 |
| 1.2 | If $z = x \sin(y) + y \cos(x)$ then $\frac{\partial^2 z}{\partial x \partial y} =$ _____. | 2 | L1 | 1 |
| 1.3 | Total differential of the function $u = x^3 e^{y z^2}$ is _____. | 2 | L2 | 2 |
| 1.4 | The critical point of the function $f(x, y) = x^2 + 2x + 9y - 3y^2 + 5$ is _____. | 2 | L1 | 1 |
| 1.5 | If $x = u \sin(v)$ and $y = u \cos(v)$ then $\frac{\partial(u,v)}{\partial(x,y)} =$ _____. | 2 | L2 | 2 |

| Sl. No. | PART -B | M | BT | CO |
|---------|---|----|----|----|
| 1 | Obtain the Maclaurin series expansion of $\log_e(1 + e^x)$ up to the term containing x^4 and hence deduce the expansion of $\frac{1}{e^{-x}(1+e^x)}$. | 10 | L2 | 2 |
| 2. (a) | Find the value of n so that the equation $V = r^n(3 \cos^2(\theta) - 1)$ satisfies the relation $\frac{\partial}{\partial r} \left(r^2 \frac{\partial V}{\partial r} \right) + \frac{1}{\sin(\theta)} \frac{\partial}{\partial \theta} \left(\sin(\theta) \frac{\partial V}{\partial \theta} \right) = 0$. | 06 | L2 | 1 |
| 2. (b) | Find $\frac{du}{dx}$ for $u = \log_e(x^2 + y^2)$, where $x^3 + y^3 + 5xy = 19$. | 04 | L2 | 2 |
| 3. (a) | Using chain rule express $\frac{\partial w}{\partial r}$ and $\frac{\partial w}{\partial s}$ in terms of r and s if $w = x + 2y + z^2$, $x = \frac{r}{s}$, $y = r^2 + \log_e(s)$, $z = 2r$. | 06 | L2 | 2 |
| 3. (b) | The pressure P (in kilopascals), volume V (in liters), and temperature T (in kelvins) of a mole of an ideal gas are related by the equation $PV = 8.31T$. Find the rate at which the pressure is changing when the temperature is 300K and increasing at a rate of 0.1K/s and the volume is 100L and increasing at a rate of 0.2L/s. | 04 | L3 | 4 |
| 4 | Find the shortest and longest distance from the point $P(1, 2, -1)$ to the sphere $x^2 + y^2 + z^2 = 24$ using Lagrange's multiplier method. | 10 | L3 | 3 |
| 5 | If $u = x + y + z$, $uv = y + z$ and $uvw = z$ then find $J = \frac{\partial(u,v,w)}{\partial(x,y,z)}$ and $J' = \frac{\partial(x,y,z)}{\partial(u,v,w)}$. Also show that $JJ' = 1$. | 10 | L3 | 3 |

BT-Blooms Taxonomy, CO-Course Outcomes, M-Marks

| Marks Distribution | Particulars | | CO1 | CO2 | CO3 | CO4 | L1 | L2 | L3 | L4 | L5 | L6 |
|--------------------|-------------|-----------|-----|-----|-----|-----|----|----|----|----|----|----|
| | Test+Quiz | Max Marks | 12 | 24 | 20 | 04 | 06 | 30 | 24 | -- | -- | -- |



RV Educational Institutions
RV College of Engineering

Autonomous
Institution Affiliated
to Visvesvaraya
Technological
University, Belagavi

Approved by AICTE,
New Delhi

| | | |
|---|---|--------------------------------------|
| Course: Fundamentals of Linear Algebra, Calculus and Statistics | DEPARTMENT OF MATHEMATICS IMPROVEMENT TEST | Maximum marks: 10+50=60 |
| Course code: 22MA11C | First semester 2022-2023 Chemistry Cycle Branch: AI, BT, CS, CD, CY, IS, SPARK C | Time: 2pm to 4pm Date: 20-03-2023 |

Instructions to candidates:

- Part A must be answered within the first two pages of the Booklet.
- Answer all questions.

| Q.No | PART- A | M | BT | CO |
|------|--|---|----|----|
| 1.1 | If the correlation co-efficient is zero, then the two regression lines are _____ to each other. | 1 | L1 | 1 |
| 1.2 | If $y = e^{at}$ is the best exponential curve for the data points $(x_i, y_i), i \in \{1, 2, 3, \dots, n\}$, then $a =$ _____. | 2 | L2 | 2 |
| 1.3 | The value of the integral $I = \int_0^2 \int_0^1 e^{x+y} dx dy$ is _____. | 2 | L2 | 1 |
| 1.4 | The first raw moment about the point 20 is 50 then its mean is _____. | 2 | L1 | 1 |
| 1.5 | Let $\sum x = 50, \sum y = 80, \sum xy = 1030, \sum x^2 = 750$ for a dataset (x_i, y_i) , where $i \in \{1, 2, 3, \dots, 10\}$. The best straight line fit for the given data is _____. | 2 | L2 | 2 |
| 1.6 | The skewness of a normal distribution is _____. | 1 | L1 | 1 |

| Q.No | PART -B | | | M | BT | CO | | | | | | | | | | | | | | |
|--------------------|--|--------|-----|------------------|---------|--------|---------------|-----|-----|--------------------|----------------|-----|----|----|----|---|----|---|----|---|
| 1.a | The first four moments about the working mean 28.5 of a distribution are 0.294, 7.144, 42.409 and 454.98. Calculate the moments about the mean, co-efficient of skewness and kurtosis using moments. Also, comment upon the skewness and kurtosis of the distribution. | | | 6 | L2 | 1 | | | | | | | | | | | | | | |
| 1.b | In a partially destroyed laboratory record, only the lines of regression of y on x and x on y are available as $4x - 5y + 33 = 0$ and $20x - 9y - 107 = 0$ respectively. Calculate the mean values of x, y and the coefficient of correlation between x and y . | | | 4 | L3 | 3 | | | | | | | | | | | | | | |
| 2.a | <p>Ten people of various heights were requested to read letters on a car at 25 yards distance. The number of letters correctly read is as given below:</p> <table border="1"><tr><td>Height (in feet)</td><td>5.1</td><td>5.3</td><td>5.6</td><td>5.7</td><td>5.8</td><td>5.9</td></tr><tr><td>No. of letters</td><td>11</td><td>17</td><td>19</td><td>14</td><td>8</td><td>15</td></tr></table> <p>Is there any correlation between heights and visual power?</p> | | | Height (in feet) | 5.1 | 5.3 | 5.6 | 5.7 | 5.8 | 5.9 | No. of letters | 11 | 17 | 19 | 14 | 8 | 15 | 6 | L2 | 2 |
| Height (in feet) | 5.1 | 5.3 | 5.6 | 5.7 | 5.8 | 5.9 | | | | | | | | | | | | | | |
| No. of letters | 11 | 17 | 19 | 14 | 8 | 15 | | | | | | | | | | | | | | |
| 2.b | <p>For two cities Kolkata and Mumbai, prices of commodities are given below:</p> <table border="1"><tr><td>City</td><td>Kolkata</td><td>Mumbai</td></tr><tr><td>Average Price</td><td>65</td><td>67</td></tr><tr><td>Standard Deviation</td><td>2.5</td><td>3.5</td></tr></table> <p>Correlation co-efficient between the prices of commodities in the two cities is 0.8, then find the most likely price in Mumbai corresponding to the price of ₹.70 at Kolkata.</p> | | | City | Kolkata | Mumbai | Average Price | 65 | 67 | Standard Deviation | 2.5 | 3.5 | 4 | L2 | 3 | | | | | |
| City | Kolkata | Mumbai | | | | | | | | | | | | | | | | | | |
| Average Price | 65 | 67 | | | | | | | | | | | | | | | | | | |
| Standard Deviation | 2.5 | 3.5 | | | | | | | | | | | | | | | | | | |

| | | | | | | | | | | | | | | | | | | | | |
|-----|--|------|------|------|------|------|------|----|----|-----|------|------|------|------|------|------|------|----|----|---|
| 3 | <p>The velocity V of a liquid is known to vary with temperature according to a quadratic law $V = a + bT + cT^2$. Find the best values of a, b and c for the following observations:</p> <table><tr><td>T</td><td>1</td><td>2</td><td>3</td><td>4</td><td>5</td><td>6</td><td>7</td></tr><tr><td>V</td><td>2.31</td><td>2.01</td><td>3.80</td><td>1.66</td><td>1.55</td><td>1.47</td><td>1.41</td></tr></table> <p>Calculate V when $T = 9$.</p> | T | 1 | 2 | 3 | 4 | 5 | 6 | 7 | V | 2.31 | 2.01 | 3.80 | 1.66 | 1.55 | 1.47 | 1.41 | 10 | L3 | 3 |
| T | 1 | 2 | 3 | 4 | 5 | 6 | 7 | | | | | | | | | | | | | |
| V | 2.31 | 2.01 | 3.80 | 1.66 | 1.55 | 1.47 | 1.41 | | | | | | | | | | | | | |
| 4.a | <p>The data from an experiment is given below. The variables y and x are connected by the relation $y = ax^b$, where a and b being constants. Fit this equation to the data by finding the values of a and b:</p> <table><tr><td>x</td><td>350</td><td>400</td><td>500</td><td>600</td></tr><tr><td>y</td><td>61</td><td>26</td><td>7</td><td>26</td></tr></table> | x | 350 | 400 | 500 | 600 | y | 61 | 26 | 7 | 26 | 6 | L2 | 2 | | | | | | |
| x | 350 | 400 | 500 | 600 | | | | | | | | | | | | | | | | |
| y | 61 | 26 | 7 | 26 | | | | | | | | | | | | | | | | |
| 4.b | <p>Evaluate the double integral</p> $\int_0^1 \int_0^{\sqrt{1+x^2}} \frac{1}{1+x^2+y^2} dy dx.$ | 4 | L2 | 2 | | | | | | | | | | | | | | | | |
| 5 | <p>Let D be a region bounded by $x = y^2, y = x - 2$</p> <p>i) Sketch the region D in the xy - plane.</p> <p>ii) Evaluate the double integral $\iint_D y \, dA$.</p> | 10 | L3 | 4 | | | | | | | | | | | | | | | | |

| Marks Distribution | Particulars | | CO1 | CO2 | CO3 | CO4 | L1 | L2 | L3 | L4 | L5 | L6 |
|--------------------|-------------|-----------|-----|-----|-----|-----|----|----|----|----|----|----|
| | Quiz & Test | Max Marks | 12 | 20 | 18 | 10 | 4 | 32 | 24 | -- | - | - |

BT-Blooms Taxonomy, CO-Course Outcomes, M-Marks

*****ALL THE BEST*****

RV COLLEGE OF ENGINEERING[®]
(An Autonomous Institution Affiliated to VTU)
1 Semester B. E. Examinations May-2023

(Common to AI & ML, BT, CS, CY, CD and IS)
**FUNDAMENTALS OF LINEAR ALGEBRA, CALCULUS AND
STATISTICS**

Time: 03 Hours

Instructions to candidates:

Maximum Marks: 100

1. Answer all questions from Part A. Part A questions should be answered in first three pages of the answer book only.
2. Answer FIVE full questions from Part B. In Part B question number 2 is compulsory. Answer any one full question from 3 and 4, 5 and 6, 7 and 8, 9 and 10.
3. Use of mathematics Handbook is permitted. Do not write anything on handbook.

PART-A

| | | |
|------|--|----|
| 1.1 | The reduced system of set of linear equations is $\begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \\ 0 \end{bmatrix}$. | |
| 1.2 | Then the solution for the systems is _____. | 01 |
| 1.3 | If two characteristic roots of a singular matrix A of order 3 are 4, 5 then the third characteristic root is _____. | 01 |
| 1.4 | The circle $x^2 + y^2 - 2ax = 0$ in polar form is _____. | 01 |
| 1.5 | The coefficient of $(x - \frac{\pi}{4})$ in the Taylor's series expansion of $\sin x$ is _____. | 01 |
| 1.6 | The curvature of the curve $y = e^x$ at the point where it crosses the y -axis is _____. | 02 |
| 1.7 | The matrices taken for the computation are $A = \begin{bmatrix} 2 & 2 \\ 3 & 0 \\ -4 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 1 \\ 2 & 3 \\ 1 & 6 \end{bmatrix}$, then the rank of the matrix $A - B$. | 02 |
| 1.8 | If the temperature of a thin wire of finite length is $u = e^{-c^2 p^2 t} [a \cos(px) + b \sin(px)]$, where a, b, p and c are constants, then $u_{xx} =$ _____. | 02 |
| 1.9 | For the implicit function $e^x - e^y = 2xy$, $\frac{dy}{dx}$ using partial differentiation is _____. | 02 |
| 1.10 | Evaluate the integral $\int_0^{\frac{\pi}{2}} \int_0^2 r^2 \sin \theta \, dr \, d\theta$. | 02 |
| 1.11 | Sketch the domain of integral $\int_0^1 \int_x^{2-x} \frac{x}{y} \, dy \, dx$ | 02 |
| 1.12 | If the first three moments of a distribution about the value 2 of the variable are 3, 16 and -20, then mean and variance of the distribution is _____. | 02 |
| 1.12 | In a partially destroyed laboratory record of an analysis of a correlation data, the following results were noted: variance of $x = 9$, equations of lines of regression of y on x is $4x - 5y + 33 = 0$ and x on y is $20x - 9y = 107$. For the given data the value of correlation coefficient is _____. | 02 |

| | | | |
|---|---|--|----|
| 2 | a | Show that the equations $\begin{aligned} -2x + y + z &= l \\ x - 2y + z &= m \\ x + y - 2z &= n \end{aligned}$ have a solution only if $l + m + n = 0$. Find all possible solutions when $l = 1, m = 1, n = -2$. | 05 |
| | b | Apply Gauss-Seidel iterative method, to solve the system of equations: $\begin{aligned} x + y + 54z &= 110 \\ 27x + 6y - z &= 85 \\ 6x + 15y + 2z &= 72 \end{aligned}$ | 05 |
| | c | Carry out three iterations using initial solution as $(0, 0, 0)$. Find the dominant eigenvalue and the corresponding eigenvector of the matrix $\begin{bmatrix} 25 & 1 & 2 \\ 1 & 3 & 0 \\ 2 & 0 & -4 \end{bmatrix}$ by Rayleigh power method taking the initial vector as $[1 \ 0 \ 0]^T$. Perform four iterations. | 06 |
| 3 | a | Find the angle of intersection of the curves $r = a \cos \theta$ and $2r = a$. | 08 |
| | b | If ρ_1 and ρ_2 be the radii of curvature at the extremities of two conjugate semi-diameters of an ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, prove that $(\rho_1^{\frac{2}{3}} + \rho_2^{\frac{2}{3}})(ab)^{2/3} = a^2 + b^2.$ | 08 |
| | | OR | |
| 4 | a | Find the curvature and the circle of curvature of the curve $\sqrt{x} + \sqrt{y} = \sqrt{a}$ at the point $(\frac{a}{4}, \frac{a}{4})$. | 08 |
| | b | Obtain the Maclaurin's series expansion of $\log_e(1 + e^x)$ up to the term containing x^4 and hence deduce the series expansion of $\frac{e^x}{1+e^x}$. | 08 |
| 5 | a | i) If $u = x^2 + y^2$, where $x = at^2, y = 2at$, show that $\frac{du}{dt} = 4a^2t(t^2 + 2)$ using partial derivatives. ii) If $u = f\left(\frac{y-x}{xy}, \frac{z-x}{xz}\right)$, find the value of $x^2u_x + y^2u_y + z^2u_z$. | 08 |
| | b | Find the volume of the greatest rectangular parallelepiped that can be inscribed in the ellipsoid, $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$, using Lagrange's method of undetermined multipliers. | 08 |
| | | OR | |
| 6 | a | If u and v are the real and imaginary parts of a complex function $f(z) = u + iv$; where $u = e^{r \cos \theta} \cos(r \sin \theta), v = e^{r \cos \theta} \sin(r \sin \theta)$ then prove that u and v satisfy the Cauchy-Reimann equations $\frac{\partial u}{\partial r} = \frac{1}{r} \frac{\partial v}{\partial \theta}$ and $\frac{\partial v}{\partial r} = -\frac{1}{r} \frac{\partial u}{\partial \theta}$. | 08 |
| | b | Prove that the functions $u = x + y + z, u = x^2 + y^2 + z^2, w = xy + yz + zx$ are functionally dependent using the concept of Jacobians and hence find the relation between them. | 0 |
| 7 | a | Evaluate $\int_0^a \int_{x^2}^{2a-x} xy \, dy \, dx$ by changing the order of integration. | |
| | b | Represent the region of integration graphically. Compute the volume of the tetrahedron formed by the planes $x = 0, y = 0, z = 0$ and $6x + 4y + 3z = 12$ using triple integration. | |

OR

- a A plate is in the form of a positive quadrant of the circle $x^2 + y^2 = 1$, the thickness ρ at any point is constant. Find the co-ordinates of the centre of gravity of the plate. 08
- b Find the area bounded by the cardioid $r = a(1 + \cos\theta)$ above the initial line using double integration. Represent the area graphically. 08

- a The growth of bacteria (y) in a community after x -hours is given by the following table.

| Hours (x) | 0 | 1 | 2 | 3 | 4 | 5 |
|----------------------------|----|----|----|----|-----|-----|
| Number of bacteria (y) | 32 | 47 | 65 | 92 | 132 | 190 |

- Find the best value of a and b in the formula $y = ab^x$ to fit this data and estimate the number of bacteria y at $x = 6$ hours by the method of least squares. 08

- b Physiological tests of intelligence and of engineering ability were applied to 10 students. Here is a record of ungrouped data showing intelligence ratio ($I.R$) and engineering ability ($E.R$). Calculate the coefficient of correlation between intelligence ratio ($I.R$) and engineering ability ($E.R$). Also find the regression line of intelligence ratio ($I.R$) on engineering ability ($E.R$) and engineering ability ($E.R$) on intelligence ratio ($I.R$).

| Student | A | B | C | D | E | F | G | H | I | J |
|---------|-----|-----|-----|-----|-----|----|-----|----|----|----|
| $I.R$ | 105 | 104 | 102 | 101 | 100 | 99 | 98 | 96 | 93 | 92 |
| $E.R$ | 101 | 103 | 100 | 98 | 95 | 96 | 104 | 92 | 97 | 94 |

08

OR

- a If the velocity V (km/hr) and Resistance R (kg/tonne) are related by a relation of the form $R = a + bV^2$, find a and b by the method of least squares with the use of the following data.

| V | 10 | 20 | 30 | 40 | 50 |
|-----|----|----|----|----|----|
| R | 8 | 10 | 15 | 21 | 30 |

- Compute the value of R when $V = 35$. 08

- b The following table gives the distribution of marks in Mathematics of 50 students in an examination. Compute $\mu_1, \mu_2, \mu_3, \mu_4$ for the following distribution. Also find β_1 and β_2 .

| Marks | 0-10 | 10-20 | 20-30 | 30-40 | 40-50 | 50-60 |
|--------------------|------|-------|-------|-------|-------|-------|
| Number of students | 1 | 6 | 10 | 15 | 11 | 7 |

08

USN

| | | | | | | | | | |
|--|--|--|--|--|--|--|--|--|--|
| | | | | | | | | | |
|--|--|--|--|--|--|--|--|--|--|

RV COLLEGE OF ENGINEERING[®]
 (An Autonomous Institution Affiliated to VTU)
 1 Semester B. E. Supplementary Examinations Oct-2023
 (Common to AI & ML, BT, CS, CY, CD and IS)
**FUNDAMENTALS OF LINEAR ALGEBRA, CALCULUS AND
 STATISTICS**

Time: 03 Hours

Maximum Marks: 100

Instructions to candidates:

1. Answer all questions from Part A. Part A questions should be answered in first three pages of the answer book only.
2. Answer FIVE full questions from Part B. In Part B question number 2 is compulsory. Answer any one full question from 3 and 4, 5 and 6, 7 and 8, 9 and 10.
3. Use of mathematics Handbook is permitted. Do not write anything on handbook.

PART-A

| | | |
|------|--|----|
| 1.1 | The Gauss elimination method reduces the system $[A:B] = \begin{bmatrix} 3 & -1 & 4 \\ 2 & 5 & -3 \end{bmatrix}$ to _____. | 01 |
| 1.2 | If two latent roots of a singular matrix A of order 3 are 2,5 then the third latent root is _____. | 01 |
| 1.3 | The point $P = \left(3, \frac{3\pi}{2}\right)$ is located on a polar curve $r = f(\theta)$, the corresponding cartesian coordinates (x,y) is _____. | 01 |
| 1.4 | For the curve $\frac{dx}{d\theta} = -3 \sin 3\theta$, $\frac{dy}{d\theta} = 3 \cos 3\theta$, $\frac{d^2x}{d\theta^2} = -9 \cos 3\theta$ and $\frac{d^2y}{d\theta^2} = -9 \sin 3\theta$, the radius of curvature is _____. | 01 |
| 1.5 | For a period of time a plane is flying along the curve $3x - 2y = 5$, the curvature of the curve is _____. | 01 |
| 1.6 | The coefficient of $\left(x - \frac{\pi}{3}\right)$ in the Taylor's series expansion of $\cos x$ is _____. | 01 |
| 1.7 | The matrices taken for the computation are $A = \begin{bmatrix} 1 & 2 \\ -1 & 0 \\ -4 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & 1 \\ 2 & 3 \\ 1 & -10 \end{bmatrix}$, then the rank of the matrix $A + B$ is _____. | 02 |
| 1.8 | If the temperature of the circular plate is $\theta = t^n e^{-\left(\frac{r^2}{4t}\right)}$, then $\frac{\partial \theta}{\partial t} =$ _____. | 02 |
| 1.9 | For the function $u(x,y) = y \cos x - (x^2 + y^2) = 0$, $\frac{dy}{dx}$ using partial differentiation is _____. | 02 |
| 1.10 | The value of $\int_0^{\pi/2} \int_0^1 r \sin \theta \, dr \, d\theta$ is _____. | 02 |
| 1.11 | Sketch the domain of integral $\int_0^1 \int_{\sqrt{y}}^{2-y} x \, dx \, dy$. | 02 |
| 1.12 | If the first four moments of a distribution about the value 3 of the variable are 2,5,23 and 159, then the fourth moment about the mean is _____. | 02 |
| 1.13 | In a partially destroyed laboratory record of an analysis of a correlation data, the following results were noted: variance of $x = 9$, equations of lines of regression of y on x is $4x - 5y + 33 = 0$ and x on y is $20x - 9y = 107$. For the given data the value of correlation coefficient is _____ and $\bar{x} =$ _____. | 02 |

PART-B

| | | | |
|---|---|--|----|
| 2 | a | Determine the values of μ for which the system $u + v + w = 1$ $2u + v + 4w = \mu$ $4u + v + 10w = \mu^2$ has a solution. Solve the system for each possible cases. | 06 |
| | b | Apply Gauss-Seidel iterative method, to solve the system of equations: $8x + y + z = 8$ $x + 3y + 5z = 5$ $2x + 4y + z = 4$ | 05 |
| | c | Carry out three iterations using initial solution as (0,0,0). Find the dominant eigenvalue and the corresponding eigenvector of the matrix $\begin{bmatrix} 4 & 1 & -1 \\ 2 & 3 & -1 \\ -2 & 1 & 5 \end{bmatrix}$ by Rayleigh power method taking the initial vector as $[1 \ 0 \ 0]^T$. Perform four iterations. | 05 |
| 3 | a | Prove that the angle of intersection of the curves $r = a \log_e \theta$ and $r = a / \log_e \theta$ is given by $\tan^{-1} \left(\frac{2e}{1-e^2} \right)$. | 08 |
| | b | Find the radius of curvature of the curve $y^2 = \frac{4a^2(2a-x)}{x}$ at the point where the curve meets the x-axis. | 08 |
| | | OR | |
| 4 | a | Find the curvature and the circle of curvature of the cissoid $y^2(2-x) = x^3$ at the point (1,1). | 08 |
| | b | Obtain the Maclaurin's series expansion of $\log_e(\sec x)$ up to the term containing x^6 and hence deduce the series expansion of $\tan x$. | 08 |
| 5 | a | i) If $u = \tan^{-1} \left(\frac{y}{x} \right)$, where $x = e^t - e^{-t}$, $y = e^t + e^{-t}$, show that $\frac{du}{dt} = -\frac{2}{e^{2t} + e^{-2t}}$ using partial derivatives. ii) Find $u \frac{\partial f}{\partial u} + v \frac{\partial f}{\partial v} + w \frac{\partial f}{\partial w}$ if $x = u + v + w$, $y = uv + vw + wu$, $z = uvw$ and $f = f(x, y, z)$. | 08 |
| | b | The temperature T at any point (x,y,z) in space is $T = 400xyz^2$. Find the highest temperature on the surface of the unit sphere $x^2 + y^2 + z^2 = 1$ using Lagrange's method of undetermined multipliers. | 08 |
| | | OR | |
| | a | If $u = \frac{1}{r} \{ \psi(r-at) + \phi(r+at) \}$, prove that $\frac{\partial^2 u}{\partial t^2} = \frac{a^2}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial u}{\partial r} \right)$. | 08 |
| | b | If $x = a^u \cos v$ and $y = a^u \sin v$, then find $J' = \frac{\partial(x,y)}{\partial(u,v)}$ and $J = \frac{\partial(u,v)}{\partial(x,y)}$. Also show that $JJ' = 1$. | 08 |
| | a | Change the order of integration and hence evaluate $\int_0^1 \int_x^{\sqrt{2-x^2}} \frac{x}{\sqrt{x^2+y^2}} dy dx$. Represent the region of integration graphically. | 08 |
| | b | Compute the volume of the ellipsoid $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$ using triple integration. | 08 |
| | | OR | |
| | a | A plate is in the form of a quadrant of the circle $x^2 + y^2 = 1$, but of varying thickness at any point given by $\rho = kxy$, where k is a constant. Find the co-ordinates of the centre of gravity of the plate. | 08 |
| | b | Find the area lying inside the cardioid $r = a(1 + \cos \theta)$ and outside the circle $r = a$ using double integration. Represent the region of integration graphically. | 08 |

9

a

The following pair of observations was noted in an experimental work on cosmic rays. Find by the method least squares the best values of a and b for the equation $y = ax^b$ which fits the data.

| | | | | | |
|-----|-----|------|------|------|-------|
| x | 2 | 3 | 4 | 5 | 6 |
| y | 8.3 | 15.4 | 33.1 | 65.2 | 127.4 |

b

Construct the table of values and compute the value of y when $x = 7$.
The experimental values relating centripetal force and radius for a mass travelling at constant velocity in a circle are as shown:

| | | | | | | |
|----------------|----|----|----|----|----|----|
| Radius (x) | 55 | 30 | 15 | 12 | 11 | 9 |
| Force (y) | 5 | 10 | 15 | 20 | 25 | 30 |

Calculate the coefficient of correlation between radius and force. Determine the equations of the regression line of force on radius and regression line of radius on force.

OR

10

a

The distance (in km) of 60 engineers from their residence to their place of work were found as follows:

| | | | | | | |
|------------------|-------|--------|---------|---------|---------|---------|
| Distance (in km) | 0 - 5 | 5 - 10 | 10 - 15 | 15 - 20 | 20 - 25 | 25 - 30 |
| No. of engineers | 8 | 11 | 15 | 12 | 9 | 5 |

Compute the first four moments about the mean and also find the measures β_1 and β_2 for the above distribution and comment on the nature of the distribution.

b

The velocity V of a liquid is known to vary with temperature according to a quadratic law $V = a + bt + cT^2$. Find the best values of a , b and c by least squares method for the following data:

| | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|
| T | 1 | 1.5 | 2 | 2.5 | 3 | 3.5 | 4 |
| V | 1.1 | 1.3 | 1.6 | 2.0 | 2.7 | 3.4 | 4.1 |

Compute the value of V when $T = 5$.