

Coursework 2

36071280

23/10/2021

Question 1

Q1. We first implement the deterministic model:

- Set $R_1 = 60$ and $F_1 = 30$.

```
R1 = 60
F1 = 30
```

- Using existing functions in R, write the necessary for loop to implement the Lotka-Volterra model that will allow you to project the number of foxes and rabbits at the end of a 2-year period i.e. after 103 more weeks.

```
for ( w in 1:(weeks-1))
{
  R1[w+1] = R1[w] + alpha* R1[w] - beta*R1[w] *F1[w]
  F1[w+1] = F1[w] + beta* R1[w]* F1[w] - gamma * F1[w]
}
```

- Print the last few values of the final result of R_t and F_t .

```
tail(R1)
```

```
## [1] 139.82668 107.94989 84.03695 66.03831 52.41642 42.03523
```

```
tail(F1)
```

```
## [1] 2316.447 2262.657 2201.461 2135.603 2067.103 1997.421
```

Question 2

Q2. A stochastic version of the Lotka-Volterra model exists in a similar manner to the stochastic version of the population growth model. In this case, the number of rabbits born is $\text{Binom}(R_t, \alpha)$, the number of rabbits eaten (new foxes) is $\text{Binom}(R_t F_t, \beta)$, and the number of foxes that die is $\text{Binom}(F_t, \gamma)$.

- Set the seed for running your code to 17540

```
#Implement the Stochastic model
```

```
#Setting initial values
```

```
set.seed(17540)
```

```
sto_R = 60
```

```
sto_Fx = 30
```

- Using existing functions in R, write the necessary for loop to implement the stochastic Lotka-Volterra model that will allow you to project the number of foxes and rabbits at the end of a 2-year period i.e. after 103 more weeks, with the same starting values as for the deterministic model.

```
for ( w in 1:(weeks-1))
{
  no_of_foxes_eaten = rbinom(1,sto_R[w]*sto_Fx[w], beta)
  sto_R[w+1] = sto_R[w] + rbinom(1,sto_R[w],alpha) - no_of_foxes_eaten
  sto_Fx[w+1] = sto_Fx[w] + no_of_foxes_eaten - rbinom(1, sto_Fx[w],gamma)
}
```

- Print the last few values of the final result of Rt and Ft

```
tail(sto_R)
```

```
## [1] 145 109 83 54 35 30
```

```
tail(sto_Fx)
```

```
## [1] 2455 2411 2350 2289 2213 2142
```

Question 3

Q3. Now we visualise the results:

- Create a long data frame called LV with three variables; time, group and size. Each row should contain the size at a single time point for one of the four groups generated rabbits and foxes (deterministic model); sto_rabbits and sto_foxes (stochastic model).

```
group_names = c('Rabbit','Fox','sto_Rabbits','sto_Foxes')
LV = data.frame(time=rep(1:weeks,4),
                group=rep(group_names, each=weeks),
                size=c(R1,F1, sto_R,sto_Fx))
```

- Using ggplot() visualise the changes over time for the number of rabbits and foxes for both the deterministic and stochastic version of the Lotka-Volterra model. All four lines should be in one single plot.

```
ggplot(LV) + geom_line(aes(time, size, color=group, linetype=group))+
  scale_linetype_manual(values=c("solid","dashed","solid","dashed"))+
  scale_color_manual(values=c(rep("orange",2), rep("blue",2)))+
  labs(x="Time ->", y="Population Size ->",
       title="Deterministic vs Stochastic model")+
  theme(legend.position = "bottom")
```

Deterministic vs Stochastic model

