

Inapproximability of counting hypergraph colourings

Andreas Galanis [†] Heng Guo [‡] Jiaheng Wang [‡]

[†]University of Oxford [‡]University of Edinburgh

July 31, 2021

Lovász Local Lemma

A set of events, each occurs with probability p and depends on at most D other events.

Lovász Local Lemma

A set of events, each occurs with probability p and depends on at most D other events.

Lemma (Lovász Local Lemma)

If $ep(D + 1) \leq 1$, then there is a non-zero probability that no event happens.

Lovász Local Lemma

A set of events, each occurs with probability p and depends on at most D other events.

Lemma (Lovász Local Lemma)

If $ep(D + 1) \leq 1$, then there is a non-zero probability that no event happens.

The “canonical” problem (originally considered in [EL75]): hypergraph colouring.

Lovász Local Lemma


A set of events, each occurs with probability p and depends on at most D other events.

Lemma (Lovász Local Lemma)

If $ep(D + 1) \leq 1$, then there is a non-zero probability that no event happens.

The “canonical” problem (originally considered in [EL75]): hypergraph colouring.

- K -uniform: K vertices in each hyperedge;
- Δ -degree: each vertex appears in $\leq \Delta$ hyperedges;
- Event: a hyperedge is monochromatic;
- $p = 1/q^{K-1}$, $D = K\Delta - 1$;
- LLL condition: $\Delta \leq \frac{q^{K-1}}{eK}$



Lovász Local Lemma


A set of events, each occurs with probability p and depends on at most D other events.

Lemma (Lovász Local Lemma)

If $ep(D + 1) \leq 1$, then there is a non-zero probability that no event happens.

The “canonical” problem (originally considered in [EL75]): hypergraph colouring.

- K -uniform: K vertices in each hyperedge;
- Δ -degree: each vertex appears in $\leq \Delta$ hyperedges;
- Event: a hyperedge is monochromatic;
- $p = 1/q^{K-1}$, $D = K\Delta - 1$;
- LLL condition: $\Delta \leq \frac{q^{K-1}}{eK}$



Other kinds of LLL-type problems:

- Boolean K -SAT;
- Constraint Satisfaction Problem;
- ...

Algorithmic LLL and Sampling/Counting LLL

Proof of LLL is non-constructive!

Algorithmic LLL and Sampling/Counting LLL

Proof of LLL is non-constructive!

LLL condition \rightarrow construct a solution efficiently?

Algorithmic LLL and Sampling/Counting LLL

Proof of LLL is non-constructive!

LLL condition \rightarrow construct a solution efficiently?

- Algorithmic LLL [MT10]: efficiently *find* a solution when LLL holds.
- Their output is far from being uniform.

Algorithmic LLL and Sampling/Counting LLL

Proof of LLL is non-constructive!

LLL condition \rightarrow construct a solution efficiently?

- Algorithmic LLL [MT10]: efficiently *find* a solution when LLL holds.
- Their output is far from being uniform.

LLL condition \rightarrow approximately sampling a (uniform) solution / counting solutions?

Algorithmic LLL and Sampling/Counting LLL

Proof of LLL is non-constructive!

LLL condition \rightarrow construct a solution efficiently?

- Algorithmic LLL [MT10]: efficiently *find* a solution when LLL holds.
- Their output is far from being uniform.

LLL condition \rightarrow approximately sampling a (uniform) solution / counting solutions?

A lot of progress from algorithmic side! Recent result by [HSW21] (even perfect samplers):

LLL-type problem	Algorithmic bound	LLL condition
Hypergraph Colourings	$\Delta \lesssim q^{K/3}$	$\Delta \lesssim q^K$
Boolean K -SAT	$\Delta \lesssim 2^{0.175K}$	$\Delta \lesssim 2^K$
General Atomic CSPs	$p^{0.175} \Delta \lesssim 1$	$p\Delta \lesssim 1$

Algorithmic LLL and Sampling/Counting LLL

Proof of LLL is non-constructive!

LLL condition \rightarrow construct a solution efficiently?

- Algorithmic LLL [MT10]: efficiently *find* a solution when LLL holds.
- Their output is far from being uniform.

Intractability region of finding a solution vs. LLL?

LLL condition \rightarrow approximately sampling a (uniform) solution / counting solutions?

A lot of progress from algorithmic side! Recent result by [HSW21] (even perfect samplers):

LLL-type problem	Algorithmic bound	LLL condition
Hypergraph Colourings	$\Delta \lesssim q^{K/3}$	$\Delta \lesssim q^K$
Boolean K -SAT	$\Delta \lesssim 2^{0.175K}$	$\Delta \lesssim 2^K$
General Atomic CSPs	$p^{0.175} \Delta \lesssim 1$	$p\Delta \lesssim 1$

Algorithmic LLL and Sampling/Counting LLL

Proof of LLL is non-constructive!

LLL condition \rightarrow construct a solution efficiently?

- Algorithmic LLL [MT10]: efficiently *find* a solution when LLL holds.
- Their output is far from being uniform.

Intractability region of finding a solution vs. LLL?

- For K -SAT, LLL is asymptotically tight [GST16].

LLL condition \rightarrow approximately sampling a (uniform) solution / counting solutions?

A lot of progress from algorithmic side! Recent result by [HSW21] (even perfect samplers):

LLL-type problem	Algorithmic bound	LLL condition
Hypergraph Colourings	$\Delta \lesssim q^{K/3}$	$\Delta \lesssim q^K$
Boolean K -SAT	$\Delta \lesssim 2^{0.175K}$	$\Delta \lesssim 2^K$
General Atomic CSPs	$p^{0.175} \Delta \lesssim 1$	$p\Delta \lesssim 1$

Algorithmic LLL and Sampling/Counting LLL

Proof of LLL is non-constructive!

LLL condition \rightarrow construct a solution efficiently?

- Algorithmic LLL [MT10]: efficiently *find* a solution when LLL holds.
- Their output is far from being uniform.

Intractability region of finding a solution vs. LLL?

- For K -SAT, LLL is asymptotically tight [GST16].

LLL condition \rightarrow approximately sampling a (uniform) solution / counting solutions?

A lot of progress from algorithmic side! Recent result by [HSW21] (even perfect samplers):

LLL-type problem	Algorithmic bound	LLL condition
Hypergraph Colourings	$\Delta \lesssim q^{K/3}$	$\Delta \lesssim q^K$
Boolean K -SAT	$\Delta \lesssim 2^{0.175K}$	$\Delta \lesssim 2^K$
General Atomic CSPs	$p^{0.175} \Delta \lesssim 1$	$p\Delta \lesssim 1$

Intractability region of sampling / counting vs. LLL?

Algorithmic LLL and Sampling/Counting LLL

Proof of LLL is non-constructive!

LLL condition \rightarrow construct a solution efficiently?

- Algorithmic LLL [MT10]: efficiently *find* a solution when LLL holds.
- Their output is far from being uniform.

Intractability region of finding a solution vs. LLL?

- For K -SAT, LLL is asymptotically tight [GST16].

LLL condition \rightarrow approximately sampling a (uniform) solution / counting solutions?

A lot of progress from algorithmic side! Recent result by [HSW21] (even perfect samplers):

LLL-type problem	Algorithmic bound	LLL condition
Hypergraph Colourings	$\Delta \lesssim q^{K/3}$	$\Delta \lesssim q^K$
Boolean K -SAT	$\Delta \lesssim 2^{0.175K}$	$\Delta \lesssim 2^K$
General Atomic CSPs	$p^{0.175} \Delta \lesssim 1$	$p\Delta \lesssim 1$

Intractability region of sampling / counting vs. LLL? \Leftarrow **Main topic of the work.**

Is hardness transition the same for counting and searching?

Is hardness transition the same for counting and searching?



Is hardness transition the same for counting and searching?



Is hardness transition the same for counting and searching?




Is hardness transition the same for counting and searching?



Theorem ([BGGGS16])

If $\Delta \gtrsim 2^{K/2}$, then it is **NP-hard** to sample a satisfying assignment from K -CNF with variable degree $\leq \Delta$, even when there is no negation in the formula (aka monotone).

Is hardness transition the same for counting and searching?



Theorem ([BGGGS16])

If $\Delta \gtrsim 2^{K/2}$, then it is **NP-hard** to sample a satisfying assignment from K -CNF with variable degree $\leq \Delta$, even when there is no negation in the formula (aka monotone).

Not true for K -SAT!



Hypergraph Colouring

Our results



Hypergraph Colouring

Algorithmic bound closer to LLL ...

Our results



Hypergraph Colouring

Algorithmic bound closer to LLL ... Chance for hardness transition to coincide at LLL??

Our results



Hypergraph Colouring


Algorithmic bound closer to LLL ... Chance for hardness transition to coincide at LLL??

Hardness for searching takes place near LLL indeed, again ...

Theorem

Let $q, K \geq 2$ be integers with $(q, K) \neq (2, 2)$. It is **NP**-hard to find a q -colouring on K -uniform simple hypergraphs of maximal degree at most Δ , when $\Delta \geq 2Kq^K \ln q + 2q$.

Our results



Hypergraph Colouring

Algorithmic bound closer to LLL ... Chance for hardness transition to coincide at LLL??

Hardness for searching takes place near LLL indeed, again ...

... but searching and counting do not coincide either! (at least for even q)

Theorem

Let $q \geq 4$ be even, $K \geq 4$ be even, and $\Delta \geq 5q^{K/2}$. It is **NP-hard** to approximate the number of proper q -colourings in n -vertex K -uniform hypergraphs of maximum degree at most Δ , even within a factor of 2^{cn} for some constant $c(q, K) > 0$.

Reduction

q -Spin system

Define q -spin system over graphs:

- Spins: $[q] = \{1, 2, 3, \dots, q\}$.
- Configuration: $\sigma : V \rightarrow [q]$;

q -Spin system

Define q -spin system over graphs:

- Spins: $[q] = \{1, 2, 3, \dots, q\}$.
- Configuration: $\sigma : V \rightarrow [q]$;
- \mathbf{B} : symmetric $q \times q$ matrix with nonnegative entries;

q -Spin system

Define q -spin system over graphs:

- Spins: $[q] = \{1, 2, 3, \dots, q\}$.
- Configuration: $\sigma : V \rightarrow [q]$;
- \mathbf{B} : symmetric $q \times q$ matrix with nonnegative entries;
- Weight of σ : $w(\sigma) := \prod_{(u,v) \in E} \mathbf{B}(\sigma(u), \sigma(v))$;

q -Spin system


Define q -spin system over graphs:

- Spins: $[q] = \{1, 2, 3, \dots, q\}$.
- Configuration: $\sigma : V \rightarrow [q]$;
- \mathbf{B} : symmetric $q \times q$ matrix with nonnegative entries;
- Weight of σ : $w(\sigma) := \prod_{(u,v) \in E} \mathbf{B}(\sigma(u), \sigma(v))$;
- **Partition function**: $Z_{\mathbf{B}}(G) := \sum_{\sigma} w(\sigma)$.

q -Spin system

Define q -spin system over graphs:

- Spins: $[q] = \{1, 2, 3, \dots, q\}$.
- Configuration: $\sigma : V \rightarrow [q]$;
- \mathbf{B} : symmetric $q \times q$ matrix with nonnegative entries;
- Weight of σ : $w(\sigma) := \prod_{(u,v) \in E} \mathbf{B}(\sigma(u), \sigma(v))$;
- **Partition function**: $Z_{\mathbf{B}}(G) := \sum_{\sigma} w(\sigma)$.




Examples:

	Potts model			
\mathbf{B}	$\begin{bmatrix} b & 1 & \cdots & 1 \\ 1 & b & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & b \end{bmatrix}$			

q -Spin system

Define q -spin system over graphs:

- Spins: $[q] = \{1, 2, 3, \dots, q\}$.
- Configuration: $\sigma : V \rightarrow [q]$;
- \mathbf{B} : symmetric $q \times q$ matrix with nonnegative entries;
- Weight of σ : $w(\sigma) := \prod_{(u,v) \in E} \mathbf{B}(\sigma(u), \sigma(v))$;
- **Partition function**: $Z_{\mathbf{B}}(G) := \sum_{\sigma} w(\sigma)$.




Examples:

	Potts model			
\mathbf{B}	$\begin{bmatrix} b & 1 & \cdots & 1 \\ 1 & b & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & b \end{bmatrix}$			

q -Spin system

Define q -spin system over graphs:

- Spins: $[q] = \{1, 2, 3, \dots, q\}$.
- Configuration: $\sigma : V \rightarrow [q]$;
- \mathbf{B} : symmetric $q \times q$ matrix with nonnegative entries;
- Weight of σ : $w(\sigma) := \prod_{(u,v) \in E} \mathbf{B}(\sigma(u), \sigma(v))$;
- **Partition function**: $Z_{\mathbf{B}}(G) := \sum_{\sigma} w(\sigma)$.




Examples:

	Potts model			
\mathbf{B}	$\begin{bmatrix} b & 1 & \dots & 1 \\ 1 & b & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \dots & b \end{bmatrix}$			

q -Spin system

Define q -spin system over graphs:

- Spins: $[q] = \{1, 2, 3, \dots, q\}$.
- Configuration: $\sigma : V \rightarrow [q]$;
- \mathbf{B} : symmetric $q \times q$ matrix with nonnegative entries;
- Weight of σ : $w(\sigma) := \prod_{(u,v) \in E} \mathbf{B}(\sigma(u), \sigma(v))$;
- **Partition function**: $Z_{\mathbf{B}}(G) := \sum_{\sigma} w(\sigma)$.




Examples:

	Potts model	Proper q -colouring		
\mathbf{B}	$\begin{bmatrix} b & 1 & \dots & 1 \\ 1 & b & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \dots & b \end{bmatrix}$	$\begin{bmatrix} 0 & 1 & \dots & 1 \\ 1 & 0 & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \dots & 0 \end{bmatrix}$		

q -Spin system

Define q -spin system over graphs:

- Spins: $[q] = \{1, 2, 3, \dots, q\}$.
- Configuration: $\sigma : V \rightarrow [q]$;
- \mathbf{B} : symmetric $q \times q$ matrix with nonnegative entries;
- Weight of σ : $w(\sigma) := \prod_{(u,v) \in E} \mathbf{B}(\sigma(u), \sigma(v))$;
- **Partition function**: $Z_{\mathbf{B}}(G) := \sum_{\sigma} w(\sigma)$.




Examples:

	Potts model	Proper q -colouring	Ising model	
\mathbf{B}	$\begin{bmatrix} b & 1 & \dots & 1 \\ 1 & b & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \dots & b \end{bmatrix}$	$\begin{bmatrix} 0 & 1 & \dots & 1 \\ 1 & 0 & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \dots & 0 \end{bmatrix}$	$\begin{bmatrix} b & 1 \\ 1 & b \end{bmatrix}$	

q -Spin system

Define q -spin system over graphs:

- Spins: $[q] = \{1, 2, 3, \dots, q\}$.
- Configuration: $\sigma : V \rightarrow [q]$;
- \mathbf{B} : symmetric $q \times q$ matrix with nonnegative entries;
- Weight of σ : $w(\sigma) := \prod_{(u,v) \in E} \mathbf{B}(\sigma(u), \sigma(v))$;
 • ... with external field: $w(\sigma) := \prod_{u \in V} f(\sigma(u)) \prod_{(u,v) \in E} \mathbf{B}(\sigma(u), \sigma(v))$
- **Partition function**: $Z_{\mathbf{B}}(G) := \sum_{\sigma} w(\sigma)$.




Examples:

	Potts model	Proper q -colouring	Ising model	
\mathbf{B}	$\begin{bmatrix} b & 1 & \dots & 1 \\ 1 & b & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \dots & b \end{bmatrix}$	$\begin{bmatrix} 0 & 1 & \dots & 1 \\ 1 & 0 & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \dots & 0 \end{bmatrix}$	$\begin{bmatrix} b & 1 \\ 1 & b \end{bmatrix}$	
			$f = [1, \lambda]$	

q -Spin system

Define q -spin system over graphs:

- Spins: $[q] = \{1, 2, 3, \dots, q\}$.
- Configuration: $\sigma : V \rightarrow [q]$;
- \mathbf{B} : symmetric $q \times q$ matrix with nonnegative entries;
- Weight of σ : $w(\sigma) := \prod_{(u,v) \in E} \mathbf{B}(\sigma(u), \sigma(v))$;
 • ... with external field: $w(\sigma) := \prod_{u \in V} f(\sigma(u)) \prod_{(u,v) \in E} \mathbf{B}(\sigma(u), \sigma(v))$
- **Partition function**: $Z_{\mathbf{B}}(G) := \sum_{\sigma} w(\sigma)$.



Examples:


	Potts model	Proper q -colouring	Ising model	Hard-core Model
\mathbf{B}	$\begin{bmatrix} b & 1 & \dots & 1 \\ 1 & b & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \dots & b \end{bmatrix}$	$\begin{bmatrix} 0 & 1 & \dots & 1 \\ 1 & 0 & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \dots & 0 \end{bmatrix}$	$\begin{bmatrix} b & 1 \\ 1 & b \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$
			$f = [1, \lambda]$	$f = [1, \lambda]$

Reduction for counting hypergraph colourings




- Consider Δ -regular graph G .

Reduction for counting hypergraph colourings




- Consider Δ -regular graph G .
- Each vertex v of $G \rightarrow k$ new vertices H_v in H_G ;

Reduction for counting hypergraph colourings




- Consider Δ -regular graph G .
- Each vertex v of $G \rightarrow k$ new vertices H_v in H_G ; each edge \rightarrow hyperedge.

Reduction for counting hypergraph colourings




- Consider Δ -regular graph G .
- Each vertex v of $G \rightarrow k$ new vertices H_v in H_G ; each edge \rightarrow hyperedge.
- H_G is Δ -regular and $K = 2k$ -uniform.

Reduction for counting hypergraph colourings




- Consider Δ -regular graph G .
- Each vertex v of $G \rightarrow k$ new vertices H_v in H_G ; each edge \rightarrow hyperedge.
- H_G is Δ -regular and $K = 2k$ -uniform.
- Consider a proper q -colouring on H_G :

Reduction for counting hypergraph colourings




- Consider Δ -regular graph G .
- Each vertex v of $G \rightarrow k$ new vertices H_v in H_G ; each edge \rightarrow hyperedge.
- H_G is Δ -regular and $K = 2k$ -uniform.
- Consider a proper q -colouring on H_G :
 - H_v has the same colour $s \in [q] \rightarrow$ assign v with spin s .

Reduction for counting hypergraph colourings



- Consider Δ -regular graph G .
- Each vertex v of $G \rightarrow k$ new vertices H_v in H_G ; each edge \rightarrow hyperedge.
- H_G is Δ -regular and $K = 2k$ -uniform.
- Consider a proper q -colouring on H_G :
 - H_v has the same colour $s \in [q] \rightarrow$ assign v with spin s .
 - H_v has mixed colours \rightarrow assign v with spin 0.

Reduction for counting hypergraph colourings



$$B = \begin{bmatrix} t^2 & t & t & \cdots & t \\ t & 0 & 1 & \cdots & 1 \\ t & 1 & 0 & \cdots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ t & 1 & 1 & \cdots & 0 \end{bmatrix}$$

$$t = (q^k - q)^{1/\Delta}.$$

- Consider Δ -regular graph G .
- Each vertex v of $G \rightarrow k$ new vertices H_v in H_G ; each edge \rightarrow hyperedge.
- H_G is Δ -regular and $K = 2k$ -uniform.
- Consider a proper q -colouring on H_G :
 - H_v has the same colour $s \in [q] \rightarrow$ assign v with spin s .
 - H_v has mixed colours \rightarrow assign v with spin 0.
- It turns out $Z_B(G) = \#\text{HYPERCOL}(H_G)$.

Inapproximability of spin systems

[DFJ02]: Hardness of approximating Hard-core model with $\lambda = 1$ (i.e., $\#IND$), $\Delta \geq 25$.


[DFJ02]: Hardness of approximating Hard-core model with $\lambda = 1$ (i.e., $\#\text{IND}$), $\Delta \geq 25$.

- Gadget: Random (d -regular) bipartite graph $G \sim \mathcal{G}_{n,n,d}$.

History

[DFJ02]: Hardness of approximating Hard-core model with $\lambda = 1$ (i.e., $\# \text{IND}$), $\Delta \geq 25$.


- Gadget: **Random (d -regular) bipartite graph** $G \sim \mathcal{G}_{n,n,d}$.
- Observe: $d < \text{some threshold}$ \rightarrow imbalanced “phase”.



History

[DFJ02]: Hardness of approximating Hard-core model with $\lambda = 1$ (i.e., $\# \text{IND}$), $\Delta \geq 25$.


- Gadget: **Random (d -regular) bipartite graph** $G \sim \mathcal{G}_{n,n,d}$.
- Observe: $d < \text{some threshold} \rightarrow \text{inbalanced "phase"}$.
- $\mathcal{E}(\alpha, \beta) := \text{Expected \# of indset s.t. } \alpha n \text{ in left, } \beta n \text{ in right.}$
- \mathcal{E} takes maximum at $\alpha \neq \beta$.



History


[DFJ02]: Hardness of approximating Hard-core model with $\lambda = 1$ (i.e., $\# \text{IND}$), $\Delta \geq 25$.

- Gadget: **Random (d -regular) bipartite graph** $G \sim \mathcal{G}_{n,n,d}$.
- Observe: $d < \text{some threshold} \rightarrow$ imbalanced “phase”.
- $\mathcal{E}(\alpha, \beta) :=$ Expected # of indset s.t. αn in left, βn in right.
- \mathcal{E} takes maximum at $\alpha \neq \beta$.
- Use this to encode variables in E2LIN2 (**NP**-hard to approx. with factor $11/12$).



[DFJ02]: Hardness of approximating Hard-core model with $\lambda = 1$ (i.e., $\# \text{IND}$), $\Delta \geq 25$.


- Gadget: **Random (d -regular) bipartite graph** $G \sim \mathcal{G}_{n,n,d}$.
- Observe: $d < \text{some threshold} \rightarrow \text{inbalanced "phase"}$.
- $\mathcal{E}(\alpha, \beta) := \text{Expected \# of indset s.t. } \alpha n \text{ in left, } \beta n \text{ in right}$.
- \mathcal{E} takes maximum at $\alpha \neq \beta$.
- Use this to encode variables in E2LIN2 (**NP**-hard to approx. with factor $11/12$).
- Didn't prove **concentration** (only Markov's inequality).



History

[DFJ02]: Hardness of approximating Hard-core model with $\lambda = 1$ (i.e., $\# \text{IND}$), $\Delta \geq 25$.

- Gadget: **Random (d -regular) bipartite graph** $G \sim \mathcal{G}_{n,n,d}$.
- Observe: $d < \text{some threshold}$ \rightarrow imbalanced “phase”.
- $\mathcal{E}(\alpha, \beta) :=$ Expected # of indset s.t. αn in left, βn in right.
- \mathcal{E} takes maximum at $\alpha \neq \beta$.
- Use this to encode variables in E2LIN2 (**NP**-hard to approx. with factor $11/12$).
- Didn't prove **concentration** (only Markov's inequality).




[MWW09, CGL12]: Locally tree-like. \rightarrow (non-)uniqueness threshold of Gibbs measure over \mathbb{T}_Δ .

History

[DFJ02]: Hardness of approximating Hard-core model with $\lambda = 1$ (i.e., $\# \text{IND}$), $\Delta \geq 25$.

- Gadget: **Random (d -regular) bipartite graph** $G \sim \mathcal{G}_{n,n,d}$.
- Observe: $d < \text{some threshold}$ \rightarrow imbalanced “phase”.
- $\mathcal{E}(\alpha, \beta) :=$ Expected # of indset s.t. αn in left, βn in right.
- \mathcal{E} takes maximum at $\alpha \neq \beta$.
- Use this to encode variables in E2LIN2 (**NP**-hard to approx. with factor $11/12$).
- Didn't prove **concentration** (only Markov's inequality).




[MWW09, CGL12]: Locally tree-like. \rightarrow (non-)uniqueness threshold of Gibbs measure over \mathbb{T}_Δ .

- (Mixing time behaves differently, though.)

History

[DFJ02]: Hardness of approximating Hard-core model with $\lambda = 1$ (i.e., $\# \text{IND}$), $\Delta \geq 25$.

- Gadget: **Random (d -regular) bipartite graph** $G \sim \mathcal{G}_{n,n,d}$.
- Observe: $d < \text{some threshold}$ \rightarrow imbalanced “phase”.
- $\mathcal{E}(\alpha, \beta) :=$ Expected # of indset s.t. αn in left, βn in right.
- \mathcal{E} takes maximum at $\alpha \neq \beta$.
- Use this to encode variables in E2LIN2 (**NP**-hard to approx. with factor $11/12$).
- Didn't prove **concentration** (only Markov's inequality).



[MWW09, CGL12]: Locally tree-like. \rightarrow (non-)uniqueness threshold of Gibbs measure over \mathbb{T}_Δ .

- (Mixing time behaves differently, though.)
- Detailed analysis over the second moment.


[Sly10]: Hardness of #IND for $\Delta \geq 6$.

[Sly10]: Hardness of #IND for $\Delta \geq 6$.

- Modify the gadget (nearly d -regular) allowing extra edges outwards.
- Even more tedious analysis over second moment.
- Reduction from MAX-CUT.


[Sly10]: Hardness of #IND for $\Delta \geq 6$.

- Modify the gadget (nearly d -regular) allowing extra edges outwards.
- Even more tedious analysis over second moment.
- Reduction from MAX-CUT.




[Sly10]: Hardness of #IND for $\Delta \geq 6$.

- Modify the gadget (nearly d -regular) allowing extra edges outwards.
- Even more tedious analysis over second moment.
- Reduction from MAX-CUT.
- Gadget phases:




[Sly10]: Hardness of #IND for $\Delta \geq 6$.

- Modify the gadget (nearly d -regular) allowing extra edges outwards.
- Even more tedious analysis over second moment.
- Reduction from MAX-CUT.
- Gadget phases:
 - + $\Pr[v^L \in I] = p^+, \Pr[v^R \in I] = p^-$.



[Sly10]: Hardness of #IND for $\Delta \geq 6$.



- Modify the gadget (nearly d -regular) allowing extra edges outwards.
- Even more tedious analysis over second moment.
- Reduction from MAX-CUT.
- Gadget phases:
 - + $\Pr[v^L \in I] = p^+, \Pr[v^R \in I] = p^-.$
 - $\Pr[v^L \in I] = p^-, \Pr[v^R \in I] = p^+.$



History

[Sly10]: Hardness of #IND for $\Delta \geq 6$.


- Modify the gadget (nearly d -regular) allowing extra edges outwards.
- Even more tedious analysis over second moment.
- Reduction from MAX-CUT.
- Gadget phases:
 - + $\Pr[v^L \in I] = p^+, \Pr[v^R \in I] = p^-$.
 - $\Pr[v^L \in I] = p^-, \Pr[v^R \in I] = p^+$.



History

[Sly10]: Hardness of #IND for $\Delta \geq 6$.


- Modify the gadget (nearly d -regular) allowing extra edges outwards.
- Even more tedious analysis over second moment.
- Reduction from MAX-CUT.
- Gadget phases:
 - + $\Pr[v^L \in I] = p^+, \Pr[v^R \in I] = p^-$.
 - $\Pr[v^L \in I] = p^-, \Pr[v^R \in I] = p^+$.



History


[Sly10]: Hardness of #IND for $\Delta \geq 6$.

- Modify the gadget (nearly d -regular) allowing extra edges outwards.
- Even more tedious analysis over second moment.
- Reduction from MAX-CUT.
- Gadget phases:
 - + $\Pr[v^L \in I] = p^+, \Pr[v^R \in I] = p^-.$
 - $\Pr[v^L \in I] = p^-, \Pr[v^R \in I] = p^+.$
- Sample I and I' respectively.



[Sly10]: Hardness of #IND for $\Delta \geq 6$.


- Modify the gadget (nearly d -regular) allowing extra edges outwards.
- Even more tedious analysis over second moment.
- Reduction from MAX-CUT.
- Gadget phases:
 - + $\Pr[v^L \in I] = p^+, \Pr[v^R \in I] = p^-.$
 - $\Pr[v^L \in I] = p^-, \Pr[v^R \in I] = p^+.$
- Sample I and I' respectively.
- Possibility I and I' are valid at v :



History


[Sly10]: Hardness of #IND for $\Delta \geq 6$.

- Modify the gadget (nearly d -regular) allowing extra edges outwards.
- Even more tedious analysis over second moment.
- Reduction from MAX-CUT.
- Gadget phases:
 - + $\Pr[v^L \in I] = p^+, \Pr[v^R \in I] = p^-.$
 - $\Pr[v^L \in I] = p^-, \Pr[v^R \in I] = p^+.$
- Sample I and I' respectively.
- Possibility I and I' are valid at v :
 - Phases agree: $(1 - (p^+)^2)(1 - (p^-)^2).$



[Sly10]: Hardness of #IND for $\Delta \geq 6$.

- Modify the gadget (nearly d -regular) allowing extra edges outwards.
- Even more tedious analysis over second moment.
- Reduction from MAX-CUT.
- Gadget phases:
 - + $\Pr[v^L \in I] = p^+, \Pr[v^R \in I] = p^-.$
 - $\Pr[v^L \in I] = p^-, \Pr[v^R \in I] = p^+.$
- Sample I and I' respectively.
- Possibility I and I' are valid at v :
 - Phases agree: $(1 - (p^+)^2)(1 - (p^-)^2).$
 - Phases disagree: $(1 - p^+ p^-)^2.$



History

[Sly10]: Hardness of #IND for $\Delta \geq 6$.


- Modify the gadget (nearly d -regular) allowing extra edges outwards.
- Even more tedious analysis over second moment.
- Reduction from MAX-CUT.

• Gadget phases:

$$+ \Pr[v^L \in I] = p^+, \Pr[v^R \in I] = p^-.$$

$$- \Pr[v^L \in I] = p^-, \Pr[v^R \in I] = p^+.$$

- Sample I and I' respectively.
- Possibility I and I' are valid at v :
 - Phases agree: $(1 - (p^+)^2)(1 - (p^-)^2)$.
 - Phases disagree: $(1 - p^+ p^-)^2$.



- (Spins of v 's are almost independent. Phases take $+/-$ with probability almost $1/2$.)

History

[Sly10]: Hardness of #IND for $\Delta \geq 6$.


- Modify the gadget (nearly d -regular) allowing extra edges outwards.
- Even more tedious analysis over second moment.
- Reduction from MAX-CUT.

• Gadget phases:

$$+ \Pr[v^L \in I] = p^+, \Pr[v^R \in I] = p^-.$$

$$- \Pr[v^L \in I] = p^-, \Pr[v^R \in I] = p^+.$$

- Sample I and I' respectively.
- Possibility I and I' are valid at v :
 - Phases agree: $(1 - (p^+)^2)(1 - (p^-)^2)$.
 - Phases disagree: $(1 - p^+ p^-)^2$.



- (Spins of v 's are almost independent. Phases take $+/-$ with probability almost $1/2$.)
- $p^+ \neq p^- \implies (1 - (p^+)^2)(1 - (p^-)^2) < (1 - p^+ p^-)^2$. Neighbour phases prefer to differ.

[SS14, GŠV16]: General antiferromagnetic Ising (2-spin) (with external field).

History

[SS14, GŠV16]: General antiferromagnetic Ising (2-spin) (with external field).

[GŠV15]: General antiferromagnetic q -spin.

[SS14, GŠV16]: General antiferromagnetic Ising (2-spin) (with external field).

[GŠV15]: General antiferromagnetic q -spin.

- Phase $\alpha, \beta \in [0, 1]^q$: fraction of each colour.

[SS14, GŠV16]: General antiferromagnetic Ising (2-spin) (with external field).

[GŠV15]: General antiferromagnetic q -spin.

- Phase $\alpha, \beta \in [0, 1]^q$: fraction of each colour.
- Ψ_1 : the exponential growth of the expectation of $Z_B^{\alpha, \beta}(G)$, $G \sim \mathcal{G}_{n, n, d}$ w.r.t. n .

[SS14, GŠV16]: General antiferromagnetic Ising (2-spin) (with external field).

[GŠV15]: General antiferromagnetic q -spin.

- Phase $\alpha, \beta \in [0, 1]^q$: fraction of each colour.
- Ψ_1 : the exponential growth of the expectation of $Z_B^{\alpha, \beta}(G)$, $G \sim \mathcal{G}_{n, n, d}$ w.r.t. n .
- Ψ_2 : exponent in second moment.

[SS14, GŠV16]: General antiferromagnetic Ising (2-spin) (with external field).

[GŠV15]: General antiferromagnetic q -spin.

- Phase $\alpha, \beta \in [0, 1]^q$: fraction of each colour.
- Ψ_1 : the exponential growth of the expectation of $Z_B^{\alpha, \beta}(G)$, $G \sim \mathcal{G}_{n, n, d}$ w.r.t. n .
- Ψ_2 : exponent in second moment.
- Dominant phase: maximizer of $\Psi_1(\alpha, \beta)$.

[SS14, GŠV16]: General antiferromagnetic Ising (2-spin) (with external field).

[GŠV15]: General antiferromagnetic q -spin.

- Phase $\alpha, \beta \in [0, 1]^q$: fraction of each colour.
- Ψ_1 : the exponential growth of the expectation of $Z_B^{\alpha, \beta}(G)$, $G \sim \mathcal{G}_{n, n, d}$ w.r.t. n .
- Ψ_2 : exponent in second moment.
- Dominant phase: maximizer of $\Psi_1(\alpha, \beta)$.
- Connect Ψ_1 and Ψ_2 for dominant phases. \rightarrow Only need to care Ψ_1 .

[SS14, GŠV16]: General antiferromagnetic Ising (2-spin) (with external field).

[GŠV15]: General antiferromagnetic q -spin.

- Phase $\alpha, \beta \in [0, 1]^q$: fraction of each colour.
- Ψ_1 : the exponential growth of the expectation of $Z_B^{\alpha, \beta}(G)$, $G \sim \mathcal{G}_{n, n, d}$ w.r.t. n .
- Ψ_2 : exponent in second moment.
- Dominant phase: maximizer of $\Psi_1(\alpha, \beta)$.
- Connect Ψ_1 and Ψ_2 for dominant phases. \rightarrow Only need to care Ψ_1 .

Theorem ([GŠV15])

Suppose $\Delta \geq 3$ and \mathbf{B} is an ergodic interaction matrix of an antiferromagnetic spin system. If the dominant phases (α, β) satisfy $\alpha \neq \beta$, are permutation symmetric and Hessian dominant, then it is NP-hard to approximate $Z_B(G)$ on an n -vertex triangle-free Δ -regular graph G .

[SS14, GŠV16]: General antiferromagnetic Ising (2-spin) (with external field).

[GŠV15]: General antiferromagnetic q -spin.

- Phase $\alpha, \beta \in [0, 1]^q$: fraction of each colour.
- Ψ_1 : the exponential growth of the expectation of $Z_B^{\alpha, \beta}(G)$, $G \sim \mathcal{G}_{n, n, d}$ w.r.t. n .
- Ψ_2 : exponent in second moment.
- Dominant phase: maximizer of $\Psi_1(\alpha, \beta)$.
- Connect Ψ_1 and Ψ_2 for dominant phases. \rightarrow Only need to care Ψ_1 .

Theorem ([GŠV15])

Suppose $\Delta \geq 3$ and B is an *ergodic* interaction matrix of an antiferromagnetic spin system. If the dominant phases (α, β) satisfy $\alpha \neq \beta$, are permutation symmetric and Hessian dominant, then it is **NP-hard** to approximate $Z_B(G)$ on an n -vertex triangle-free Δ -regular graph G .

Ergodic: B is irreducible and aperiodic.

[SS14, GŠV16]: General antiferromagnetic Ising (2-spin) (with external field).

[GŠV15]: General antiferromagnetic q -spin.

- Phase $\alpha, \beta \in [0, 1]^q$: fraction of each colour.
- Ψ_1 : the exponential growth of the expectation of $Z_B^{\alpha, \beta}(G)$, $G \sim \mathcal{G}_{n, n, d}$ w.r.t. n .
- Ψ_2 : exponent in second moment.
- Dominant phase: maximizer of $\Psi_1(\alpha, \beta)$.
- Connect Ψ_1 and Ψ_2 for dominant phases. \rightarrow Only need to care Ψ_1 .

Theorem ([GŠV15])

Suppose $\Delta \geq 3$ and B is an ergodic interaction matrix of an *antiferromagnetic* spin system. If the dominant phases (α, β) satisfy $\alpha \neq \beta$, are permutation symmetric and Hessian dominant, then it is **NP-hard** to approximate $Z_B(G)$ on an n -vertex triangle-free Δ -regular graph G .

antiferromagnetic: $\lambda_2(B) < 0$.

[SS14, GŠV16]: General antiferromagnetic Ising (2-spin) (with external field).

[GŠV15]: General antiferromagnetic q -spin.

- Phase $\alpha, \beta \in [0, 1]^q$: fraction of each colour.
- Ψ_1 : the exponential growth of the expectation of $Z_B^{\alpha, \beta}(G)$, $G \sim \mathcal{G}_{n, n, d}$ w.r.t. n .
- Ψ_2 : exponent in second moment.
- Dominant phase: maximizer of $\Psi_1(\alpha, \beta)$.
- Connect Ψ_1 and Ψ_2 for dominant phases. \rightarrow Only need to care Ψ_1 .

Theorem ([GŠV15])

Suppose $\Delta \geq 3$ and \mathbf{B} is an ergodic interaction matrix of an antiferromagnetic spin system. If the dominant phases (α, β) satisfy $\alpha \neq \beta$, are *permutation symmetric* and Hessian dominant, then it is **NP-hard** to approximate $Z_B(G)$ on an n -vertex triangle-free Δ -regular graph G .

Permutation symmetric: dominant phases can be obtained from each other, by permutating spins while leaving \mathbf{B} invariant, or switch α, β , or both.

[SS14, GŠV16]: General antiferromagnetic Ising (2-spin) (with external field).

[GŠV15]: General antiferromagnetic q -spin.

- Phase $\alpha, \beta \in [0, 1]^q$: fraction of each colour.
- Ψ_1 : the exponential growth of the expectation of $Z_B^{\alpha, \beta}(G)$, $G \sim \mathcal{G}_{n, n, d}$ w.r.t. n .
- Ψ_2 : exponent in second moment.
- Dominant phase: maximizer of $\Psi_1(\alpha, \beta)$.
- Connect Ψ_1 and Ψ_2 for dominant phases. \rightarrow Only need to care Ψ_1 .

Theorem ([GŠV15])

*Suppose $\Delta \geq 3$ and B is an ergodic interaction matrix of an antiferromagnetic spin system. If the dominant phases (α, β) satisfy $\alpha \neq \beta$, are permutation symmetric and **Hessian** dominant, then it is NP-hard to approximate $Z_B(G)$ on an n -vertex triangle-free Δ -regular graph G .*

Hessian: the Hessian of Ψ_1 is negative-definite.

From first moment to tree recursion

Ψ_1 has explicit form... but inconvenient to work with!

From first moment to tree recursion

Ψ_1 has explicit form... but inconvenient to work with!

- They introduce a simpler function ($\mathbf{r}, \mathbf{c} \in \mathbb{R}_{\geq 0}^q$):

$$\Phi(\mathbf{r}, \mathbf{c}) := \ln \frac{\mathbf{r}^\top \mathbf{B} \mathbf{c}}{\|\mathbf{r}\|_p \|\mathbf{c}\|_p} \quad \text{where } p = \Delta/(\Delta - 1).$$

From first moment to tree recursion

Ψ_1 has explicit form... but inconvenient to work with!

- They introduce a simpler function ($\mathbf{r}, \mathbf{c} \in \mathbb{R}_{\geq 0}^q$):

$$\Phi(\mathbf{r}, \mathbf{c}) := \ln \frac{\mathbf{r}^\top \mathbf{B} \mathbf{c}}{\|\mathbf{r}\|_p \|\mathbf{c}\|_p} \quad \text{where } p = \Delta/(\Delta - 1).$$

- There's a one-to-one mapping (up to scaling) between critical points of Ψ_1 and Φ ; values are equal (up to a factor Δ) at these points. No local maximal at boundaries for either function.

From first moment to tree recursion

Ψ_1 has explicit form... but inconvenient to work with!

- They introduce a simpler function ($\mathbf{r}, \mathbf{c} \in \mathbb{R}_{\geq 0}^q$):

$$\Phi(\mathbf{r}, \mathbf{c}) := \ln \frac{\mathbf{r}^\top \mathbf{B} \mathbf{c}}{\|\mathbf{r}\|_p \|\mathbf{c}\|_p} \quad \text{where } p = \Delta/(\Delta - 1).$$

- There's a one-to-one mapping (up to scaling) between critical points of Ψ_1 and Φ ; values are equal (up to a factor Δ) at these points. No local maximal at boundaries for either function.

Critical points of Φ satisfies ($d := \Delta - 1$):

$$R_i \propto \left(\sum_{j \in [q]} \mathbf{B}_{ij} C_j \right)^d ; \quad C_j \propto \left(\sum_{i \in [q]} \mathbf{B}_{ij} R_i \right)^d .$$

From first moment to tree recursion

Ψ_1 has explicit form... but inconvenient to work with!

- They introduce a simpler function ($\mathbf{r}, \mathbf{c} \in \mathbb{R}_{\geq 0}^q$):

$$\Phi(\mathbf{r}, \mathbf{c}) := \ln \frac{\mathbf{r}^\top \mathbf{B} \mathbf{c}}{\|\mathbf{r}\|_p \|\mathbf{c}\|_p} \quad \text{where } p = \Delta/(\Delta - 1).$$

- There's a one-to-one mapping (up to scaling) between critical points of Ψ_1 and Φ ; values are equal (up to a factor Δ) at these points. No local maximal at boundaries for either function.

Critical points of Φ satisfies ($d := \Delta - 1$):

$$R_i \propto \left(\sum_{j \in [q]} \mathbf{B}_{ij} C_j \right)^d ; \quad C_j \propto \left(\sum_{i \in [q]} \mathbf{B}_{ij} R_i \right)^d .$$

“Tree recursion”: Recursion for marginals on infinite d -ary tree. R : odd layer; C : even layer.

From first moment to tree recursion

Ψ_1 has explicit form... but inconvenient to work with!

- They introduce a simpler function ($\mathbf{r}, \mathbf{c} \in \mathbb{R}_{\geq 0}^q$):

$$\Phi(\mathbf{r}, \mathbf{c}) := \ln \frac{\mathbf{r}^\top \mathbf{B} \mathbf{c}}{\|\mathbf{r}\|_p \|\mathbf{c}\|_p} \quad \text{where } p = \Delta/(\Delta - 1).$$

- There's a one-to-one mapping (up to scaling) between critical points of Ψ_1 and Φ ; values are equal (up to a factor Δ) at these points. No local maximal at boundaries for either function.

Critical points of Φ satisfies ($d := \Delta - 1$):

$$R_i \propto \left(\sum_{j \in [q]} \mathbf{B}_{ij} C_j \right)^d ; \quad C_j \propto \left(\sum_{i \in [q]} \mathbf{B}_{ij} R_i \right)^d .$$

“Tree recursion”: Recursion for marginals on infinite d -ary tree. R : odd layer; C : even layer.

Jacobian stable fixpoints of tree recursion \iff **Hessian** local maxima of Ψ_1 .

Dominant phase analysis

Proper q -colourings ([GŠV15])

Φ is still not easy to analyse:

Proper q -colourings ([GŠV15])

Φ is still not easy to analyse:

- Each candidate maximizer of Φ is local maxima. \rightarrow Cannot perturb R_i, C_i 's;

Proper q -colourings ([GŠV15])

Φ is still not easy to analyse:

- Each candidate maximizer of Φ is local maxima. \rightarrow Cannot perturb R_i, C_i 's;
- Φ has $2q$ variables.

Proper q -colourings ([GŠV15])

Φ is still not easy to analyse:

- Each candidate maximizer of Φ is local maxima. \rightarrow Cannot perturb R_i, C_i 's;
- Φ has $2q$ variables.

Observe: any fixpoint of the tree recursion for q -colourings has support size ≤ 3 in each side.

Proper q -colourings ([GŠV15])

Φ is still not easy to analyse:

- Each candidate maximizer of Φ is local maxima. \rightarrow Cannot perturb R_i, C_i 's;
- Φ has $2q$ variables.

Observe: any fixpoint of the tree recursion for q -colourings has support size ≤ 3 in each side.

- New variables: $q_i, R_i, C_i, i = 1, 2, 3$.

Proper q -colourings ([GŠV15])

Φ is still not easy to analyse:

- Each candidate maximizer of Φ is local maxima. \rightarrow Cannot perturb R_i, C_i 's;
- Φ has $2q$ variables.

Observe: any fixpoint of the tree recursion for q -colourings has support size ≤ 3 in each side.

- New variables: $q_i, R_i, C_i, i = 1, 2, 3$. Rewrite Φ . \rightarrow Optimization with only 9 variables!

Proper q -colourings ([GŠV15])

Φ is still not easy to analyse:

- Each candidate maximizer of Φ is local maxima. \rightarrow Cannot perturb R_i, C_i 's;
- Φ has $2q$ variables.

Observe: any fixpoint of the tree recursion for q -colourings has support size ≤ 3 in each side.

- New variables: $q_i, R_i, C_i, i = 1, 2, 3$. Rewrite Φ . \rightarrow Optimization with only 9 variables!
- Relaxation: allow q_i 's to take any **nonnegative real**.

Proper q -colourings ([GŠV15])

Φ is still not easy to analyse:

- Each candidate maximizer of Φ is local maxima. \rightarrow Cannot perturb R_i, C_i 's;
- Φ has $2q$ variables.

Observe: any fixpoint of the tree recursion for q -colourings has support size ≤ 3 in each side.

- New variables: $q_i, R_i, C_i, i = 1, 2, 3$. Rewrite Φ . \rightarrow Optimization with only 9 variables!
- Relaxation: allow q_i 's to take any **nonnegative real**. (Need some extra check at boundary)
- **Perturb q_i 's** to jump out local maxima.

Proper q -colourings ([GŠV15])

Φ is still not easy to analyse:

- Each candidate maximizer of Φ is local maxima. \rightarrow Cannot perturb R_i, C_i 's;
- Φ has $2q$ variables.

Observe: any fixpoint of the tree recursion for q -colourings has support size ≤ 3 in each side.

- New variables: $q_i, R_i, C_i, i = 1, 2, 3$. Rewrite Φ . \rightarrow Optimization with only 9 variables!
- Relaxation: allow q_i 's to take any **nonnegative real**. (Need some extra check at boundary)
- **Perturb q_i 's** to jump out local maxima.

(q_1, q_2, q_3) -type fixpoint: $q_i > 0$ entries take $R_i (C_i)$ in $\mathbf{r}(\mathbf{c})$; $R_i \neq R_j, C_i \neq C_j$ for $i \neq j, q_{i,j} > 0$.

Proper q -colourings ([GŠV15])

Φ is still not easy to analyse:

- Each candidate maximizer of Φ is local maxima. \rightarrow Cannot perturb R_i, C_i 's;
- Φ has $2q$ variables.

Observe: any fixpoint of the tree recursion for q -colourings has support size ≤ 3 in each side.

- New variables: $q_i, R_i, C_i, i = 1, 2, 3$. Rewrite Φ . \rightarrow Optimization with only 9 variables!
- Relaxation: allow q_i 's to take any **nonnegative real**. (Need some extra check at boundary)
- **Perturb q_i 's** to jump out local maxima.

(q_1, q_2, q_3) -type fixpoint: $q_i > 0$ entries take $R_i (C_i)$ in $\mathbf{r}(\mathbf{c})$; $R_i \neq R_j, C_i \neq C_j$ for $i \neq j, q_{i,j} > 0$.

Candidate maximizers via perturbation argument:

Proper q -colourings ([GŠV15])

Φ is still not easy to analyse:

- Each candidate maximizer of Φ is local maxima. \rightarrow Cannot perturb R_i, C_i 's;
- Φ has $2q$ variables.

Observe: any fixpoint of the tree recursion for q -colourings has support size ≤ 3 in each side.

- New variables: $q_i, R_i, C_i, i = 1, 2, 3$. Rewrite Φ . \rightarrow Optimization with only 9 variables!
- Relaxation: allow q_i 's to take any **nonnegative real**. (Need some extra check at boundary)
- **Perturb q_i 's** to jump out local maxima.

(q_1, q_2, q_3) -type fixpoint: $q_i > 0$ entries take $R_i (C_i)$ in $\mathbf{r}(\mathbf{c})$; $R_i \neq R_j, C_i \neq C_j$ for $i \neq j, q_{i,j} > 0$.

Candidate maximizers via perturbation argument:

- $(q/2, q/2, 0)$ with $R_1/R_2 = C_2/C_1$.

Proper q -colourings ([GŠV15])

Φ is still not easy to analyse:

- Each candidate maximizer of Φ is local maxima. \rightarrow Cannot perturb R_i, C_i 's;
- Φ has $2q$ variables.

Observe: any fixpoint of the tree recursion for q -colourings has support size ≤ 3 in each side.

- New variables: $q_i, R_i, C_i, i = 1, 2, 3$. Rewrite Φ . \rightarrow Optimization with only 9 variables!
- Relaxation: allow q_i 's to take any **nonnegative real**. (Need some extra check at boundary)
- **Perturb q_i 's** to jump out local maxima.

(q_1, q_2, q_3) -type fixpoint: $q_i > 0$ entries take $R_i (C_i)$ in $\mathbf{r}(\mathbf{c})$; $R_i \neq R_j, C_i \neq C_j$ for $i \neq j, q_{i,j} > 0$.

Candidate maximizers via perturbation argument:

- $(q/2, q/2, 0)$ with $R_1/R_2 = C_2/C_1$.
- $(q, 0, 0)$.

Proper q -colourings ([GŠV15])

Φ is still not easy to analyse:

- Each candidate maximizer of Φ is local maxima. \rightarrow Cannot perturb R_i, C_i 's;
- Φ has $2q$ variables.

Observe: any fixpoint of the tree recursion for q -colourings has support size ≤ 3 in each side.

- New variables: $q_i, R_i, C_i, i = 1, 2, 3$. Rewrite Φ . \rightarrow Optimization with only 9 variables!
- Relaxation: allow q_i 's to take any **nonnegative real**. (Need some extra check at boundary)
- **Perturb q_i 's** to jump out local maxima.

(q_1, q_2, q_3) -type fixpoint: $q_i > 0$ entries take $R_i (C_i)$ in $\mathbf{r}(\mathbf{c})$; $R_i \neq R_j, C_i \neq C_j$ for $i \neq j, q_{i,j} > 0$.

Candidate maximizers via perturbation argument:

- $(q/2, q/2, 0)$ with $R_1/R_2 = C_2/C_1$.
- $(q, 0, 0)$. \rightarrow **Ruled out by checking stability.**

Proper q -colourings ([GŠV15])

Φ is still not easy to analyse:

- Each candidate maximizer of Φ is local maxima. \rightarrow Cannot perturb R_i, C_i 's;
- Φ has $2q$ variables.

Observe: any fixpoint of the tree recursion for q -colourings has support size ≤ 3 in each side.

- New variables: $q_i, R_i, C_i, i = 1, 2, 3$. Rewrite Φ . \rightarrow Optimization with only 9 variables!
- Relaxation: allow q_i 's to take any **nonnegative real**. (Need some extra check at boundary)
- **Perturb q_i 's** to jump out local maxima.

(q_1, q_2, q_3) -type fixpoint: $q_i > 0$ entries take $R_i (C_i)$ in $\mathbf{r}(\mathbf{c})$; $R_i \neq R_j, C_i \neq C_j$ for $i \neq j, q_{i,j} > 0$.

Candidate maximizers via perturbation argument:

- $(q/2, q/2, 0)$ with $R_1/R_2 = C_2/C_1$.
- $(q, 0, 0)$. \rightarrow **Ruled out by checking stability**.

The reason why they can only deal with **even q** .

Our case

Recall:

Proper q -colouring	Our case
$\begin{bmatrix} 0 & 1 & \cdots & 1 \\ 1 & 0 & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & 0 \end{bmatrix}$	$\begin{bmatrix} t^2 & t & t & \cdots & t \\ t & 0 & 1 & \cdots & 1 \\ t & 1 & 0 & \cdots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ t & 1 & 1 & \cdots & 0 \end{bmatrix}$

Our case

Recall:

Proper q -colouring	Our case
$\begin{bmatrix} 0 & 1 & \cdots & 1 \\ 1 & 0 & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & 0 \end{bmatrix}$	$\begin{bmatrix} t^2 & t & t & \cdots & t \\ t & 0 & 1 & \cdots & 1 \\ t & 1 & 0 & \cdots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ t & 1 & 1 & \cdots & 0 \end{bmatrix}$

- \mathbf{r} and \mathbf{c} has support size ≤ 3 respectively (except R_0 and C_0).

Our case

Recall:

Proper q -colouring	Our case
$\begin{bmatrix} 0 & 1 & \cdots & 1 \\ 1 & 0 & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & 0 \end{bmatrix}$	$\begin{bmatrix} t^2 & t & t & \cdots & t \\ t & 0 & 1 & \cdots & 1 \\ t & 1 & 0 & \cdots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ t & 1 & 1 & \cdots & 0 \end{bmatrix}$

- \mathbf{r} and \mathbf{c} has support size ≤ 3 respectively (except R_0 and C_0).
- 11 variables after rewriting Φ .

Our case

Recall:

Proper q -colouring	Our case
$\begin{bmatrix} 0 & 1 & \cdots & 1 \\ 1 & 0 & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & 0 \end{bmatrix}$	$\begin{bmatrix} t^2 & t & t & \cdots & t \\ t & 0 & 1 & \cdots & 1 \\ t & 1 & 0 & \cdots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ t & 1 & 1 & \cdots & 0 \end{bmatrix}$

- \mathbf{r} and \mathbf{c} has support size ≤ 3 respectively (except R_0 and C_0).
- 11 variables after rewriting Φ .

Maximizer behaves differently for $d = \Delta - 1$ above or below some threshold.

Our case

Recall:

Proper q -colouring	Our case
$\begin{bmatrix} 0 & 1 & \cdots & 1 \\ 1 & 0 & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & 0 \end{bmatrix}$	$\begin{bmatrix} t^2 & t & t & \cdots & t \\ t & 0 & 1 & \cdots & 1 \\ t & 1 & 0 & \cdots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ t & 1 & 1 & \cdots & 0 \end{bmatrix}$

- \mathbf{r} and \mathbf{c} has support size ≤ 3 respectively (except R_0 and C_0).
- 11 variables after rewriting Φ .

Maximizer behaves differently for $d = \Delta - 1$ above or below some threshold.

- In fact, [GŠV15] considers Potts with $b < \frac{\Delta - q}{\Delta}$.

Guess the fixpoints! ($d = 5q^k$)

Guess the fixpoints! ($d = 5q^k$)

Support size 3?

Guess the fixpoints! ($d = 5q^k$)

Support size 3? \times

Guess the fixpoints! ($d = 5q^k$)

Support size 3? \times

Support size 2?

Guess the fixpoints! ($d = 5q^k$)

Support size 3? \times

Support size 2?

- $(q/2, q/2, 0)$ with $R_0/R_1 = C_0/C_3$.

Guess the fixpoints! ($d = 5q^k$)

Support size 3? \times

Support size 2?

- $(q/2, q/2, 0)$ with $R_0/R_1 = C_0/C_3$.
- $q = 6, k = 3$:

$\mathbf{r} = 0.9863, 0.0045, 0.0045, 0.0045, 0.0001, 0.0001, 0.0001$;

$\mathbf{c} = 0.9863, 0.0001, 0.0001, 0.0001, 0.0045, 0.0045, 0.0045$.

Guess the fixpoints! ($d = 5q^k$)

Support size 3? \times

Support size 2?

- $(q/2, q/2, 0)$ with $R_0/R_1 = C_0/C_3$.
- **Unique** and stable.

Guess the fixpoints! ($d = 5q^k$)

Support size 3? \times

Support size 2?

- $(q/2, q/2, 0)$ with $R_0/R_1 = C_0/C_3$.
- **Unique** and stable.

Support size 1 (i.e., $(q, 0, 0)$)?

Guess the fixpoints! ($d = 5q^k$)

Support size 3? \times

Support size 2?

- $(q/2, q/2, 0)$ with $R_0/R_1 = C_0/C_3$.
- **Unique** and stable.

Support size 1 (i.e., $(q, 0, 0)$)?

$$\begin{bmatrix} t^2 & t & t & \cdots & t \\ t & 0 & 1 & \cdots & 1 \\ t & 1 & 0 & \cdots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ t & 1 & 1 & \cdots & 0 \end{bmatrix}$$

Guess the fixpoints! ($d = 5q^k$)

Support size 3? \times

Support size 2?

- $(q/2, q/2, 0)$ with $R_0/R_1 = C_0/C_3$.
- **Unique** and stable.

Support size 1 (i.e., $(q, 0, 0)$)?

$$\begin{bmatrix} t^2 & t & t & \cdots & t \\ t & 0 & 1 & \cdots & 1 \\ t & 1 & 0 & \cdots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ t & 1 & 1 & \cdots & 0 \end{bmatrix} \rightarrow \begin{bmatrix} * & * \\ * & * \end{bmatrix}$$

Guess the fixpoints! ($d = 5q^k$)

Support size 3? \times

Support size 2?

- $(q/2, q/2, 0)$ with $R_0/R_1 = C_0/C_3$.
- **Unique** and stable.

Support size 1 (i.e., $(q, 0, 0)$)?

- Behaves like a 2-spin due to the “mixed” colour!

Guess the fixpoints! ($d = 5q^k$)

Support size 3? \times

Support size 2?

- $(q/2, q/2, 0)$ with $R_0/R_1 = C_0/C_3$.
- **Unique** and stable.

Support size 1 (i.e., $(q, 0, 0)$)?

- Behaves like a 2-spin due to the “mixed” colour!
- General antiferro 2-spin: balanced solution (Q^*, Q^*) ,

Guess the fixpoints! ($d = 5q^k$)

Support size 3? \times

Support size 2?

- $(q/2, q/2, 0)$ with $R_0/R_1 = C_0/C_3$.
- **Unique** and stable.

Support size 1 (i.e., $(q, 0, 0)$)?

- Behaves like a 2-spin due to the “mixed” colour!
- General antiferro 2-spin: balanced solution (Q^*, Q^*) ,
- (corresponds to $R_0/R_1 = C_0/C_1$,

$$\mathbf{r} = \mathbf{c} = 0.993, 0.001, 0.001, 0.001, 0.001, 0.001, 0.001.$$

Guess the fixpoints! ($d = 5q^k$)

Support size 3? \times

Support size 2?

- $(q/2, q/2, 0)$ with $R_0/R_1 = C_0/C_3$.
- **Unique** and stable.

Support size 1 (i.e., $(q, 0, 0)$)?

- Behaves like a 2-spin due to the “mixed” colour!
- General antiferro 2-spin: balanced solution (Q^*, Q^*) ,
- (corresponds to $R_0/R_1 = C_0/C_1$, unstable.)

$$\mathbf{r} = \mathbf{c} = 0.993, 0.001, 0.001, 0.001, 0.001, 0.001, 0.001.$$

Guess the fixpoints! ($d = 5q^k$)

Support size 3? \times

Support size 2?

- $(q/2, q/2, 0)$ with $R_0/R_1 = C_0/C_3$.
- **Unique** and stable.

Support size 1 (i.e., $(q, 0, 0)$)?

- Behaves like a 2-spin due to the “mixed” colour!
- General antiferro 2-spin: balanced solution (Q^*, Q^*) ,
- (corresponds to $R_0/R_1 = C_0/C_1$, unstable.)
- ... and inbalanced ones (Q^+, Q^-) , (Q^-, Q^+) in **non-uniqueness region**.

Guess the fixpoints! ($d = 5q^k$)

Support size 3? \times

Support size 2?

- $(q/2, q/2, 0)$ with $R_0/R_1 = C_0/C_3$.
- **Unique** and stable.

Support size 1 (i.e., $(q, 0, 0)$)?

- Behaves like a 2-spin due to the “mixed” colour!
- General antiferro 2-spin: balanced solution (Q^*, Q^*) ,
- (corresponds to $R_0/R_1 = C_0/C_1$, unstable.)
- ... and inbalanced ones (Q^+, Q^-) , (Q^-, Q^+) in **non-uniqueness region**.
- (correspond to $R_0/R_1 \neq C_0/C_1$,

$$\mathbf{r} = 0.9997, 0.0001, 0.0001, 0.0001, 0.0001, 0.0001, 0.0001;$$

$$\mathbf{c} = 0.9732, 0.0045, 0.0045, 0.0045, 0.0045, 0.0045, 0.0045.$$

Guess the fixpoints! ($d = 5q^k$)

Support size 3? \times

Support size 2?

- $(q/2, q/2, 0)$ with $R_0/R_1 = C_0/C_3$.
- **Unique** and stable.

Support size 1 (i.e., $(q, 0, 0)$)?

- Behaves like a 2-spin due to the “mixed” colour!
- General antiferro 2-spin: balanced solution (Q^*, Q^*) ,
- (corresponds to $R_0/R_1 = C_0/C_1$, unstable.)
- ... and inbalanced ones (Q^+, Q^-) , (Q^-, Q^+) in **non-uniqueness region**.
- (correspond to $R_0/R_1 \neq C_0/C_1$, **stable**.)

$$\mathbf{r} = 0.9997, 0.0001, 0.0001, 0.0001, 0.0001, 0.0001, 0.0001;$$

$$\mathbf{c} = 0.9732, 0.0045, 0.0045, 0.0045, 0.0045, 0.0045, 0.0045.$$

Guess the fixpoints! ($d = 5q^k$)

Support size 3? \times

Support size 2? \checkmark

- $(q/2, q/2, 0)$ with $R_0/R_1 = C_0/C_3$.
- **Unique** and stable.

Support size 1 (i.e., $(q, 0, 0)$)?

- Behaves like a 2-spin due to the “mixed” colour!
- General antiferro 2-spin: balanced solution (Q^*, Q^*) ,
- (corresponds to $R_0/R_1 = C_0/C_1$, unstable.)
- ... and imbalanced ones (Q^+, Q^-) , (Q^-, Q^+) in **non-uniqueness region**.
- (correspond to $R_0/R_1 \neq C_0/C_1$, **stable**.)

Use a more careful interpolate-and-perturb argument to show 2-supported is global maxima.

Guess the fixpoints! ($d = 5q^k$)

Support size 3? \times

Support size 2? \checkmark

- $(q/2, q/2, 0)$ with $R_0/R_1 = C_0/C_3$.
- **Unique** and stable.

Support size 1 (i.e., $(q, 0, 0)$)?

- Behaves like a 2-spin due to the “mixed” colour!
- General antiferro 2-spin: balanced solution (Q^*, Q^*) ,
- (corresponds to $R_0/R_1 = C_0/C_1$, unstable.)
- ... and imbalanced ones (Q^+, Q^-) , (Q^-, Q^+) in **non-uniqueness region**.
- (correspond to $R_0/R_1 \neq C_0/C_1$, **stable**.)

Use a more careful interpolate-and-perturb argument to show 2-supported is global maxima.

- $(q, 0, 0)$ with $R_0/R_1 \neq C_0/C_1$ can be regarded as a limit of $(q_1, q_2, 0)$ fixpoint.

Guess the fixpoints! ($d = q^k$)

Support size 3?

Support size 2?

Support size 1?

Guess the fixpoints! ($d = q^k$)

Support size 3? \times

Support size 2?

Support size 1?

Guess the fixpoints! ($d = q^k$)

Support size 3? \times

Support size 2? \times ($(q/2, q/2, 0)$ with $R_0/R_1 = C_0/C_3$ no more exists.)

Support size 1?

Guess the fixpoints! ($d = q^k$)

Support size 3? \times

Support size 2? \times ($(q/2, q/2, 0)$ with $R_0/R_1 = C_0/C_3$ no more exists.)

Support size 1?

- Lies in **uniqueness region**.

Guess the fixpoints! ($d = q^k$)

Support size 3? \times

Support size 2? \times ($(q/2, q/2, 0)$ with $R_0/R_1 = C_0/C_3$ no more exists.)

Support size 1?

- Lies in **uniqueness region**.
- Only solution: (Q^*, Q^*) .

Guess the fixpoints! ($d = q^k$)

Support size 3? \times

Support size 2? \times ($(q/2, q/2, 0)$ with $R_0/R_1 = C_0/C_3$ no more exists.)

Support size 1?

- Lies in **uniqueness region**.
- Only solution: (Q^*, Q^*) .
- Stable in this case.

Guess the fixpoints! ($d = q^k$)

Support size 3? \times

Support size 2? $\times ((q/2, q/2, 0)$ with $R_0/R_1 = C_0/C_3$ no more exists.)

Support size 1?

- Lies in **uniqueness region**.
- Only solution: (Q^*, Q^*) .
- Stable in this case.
- Translation-invariant.

$$\mathbf{r} = \mathbf{c} = 0.984, 0.003, 0.003, 0.003, 0.003, 0.003, 0.003.$$

Guess the fixpoints! ($d = q^k$)

Support size 3? \times

Support size 2? \times ($(q/2, q/2, 0)$ with $R_0/R_1 = C_0/C_3$ no more exists.)

Support size 1?

- Lies in **uniqueness region**.
- Only solution: (Q^*, Q^*) .
- Stable in this case.
- Translation-invariant.

$$\mathbf{r} = \mathbf{c} = 0.984, 0.003, 0.003, 0.003, 0.003, 0.003, 0.003.$$

Dominant phase satisfies $\alpha = \beta$. Cannot apply **[GŠV15]**.

Summary

Future directions

Hardness side: handle odd q .

Future directions

Hardness side: handle odd q .

- More detailed comparison between candidates? (Meanwhile improving **[GŠV15]**?)

Future directions

Hardness side: handle odd q .

- More detailed comparison between candidates? (Meanwhile improving [GŠV15]?)
- New gadgets from graph to hypergraph?

Future directions

Hardness side: handle odd q .

- More detailed comparison between candidates? (Meanwhile improving [GŠV15]?)
- New gadgets from graph to hypergraph?
- Analyse asymptotically when, say, $\Delta \gtrsim q^{0.51K}$?

Future directions

Hardness side: handle odd q .

- More detailed comparison between candidates? (Meanwhile improving [GŠV15]?)
- New gadgets from graph to hypergraph?
- Analyse asymptotically when, say, $\Delta \gtrsim q^{0.51K}$?

Algorithmic side: close the gap between $q^{K/3}$ and $q^{K/2}$.


Future directions

Hardness side: handle odd q .

- More detailed comparison between candidates? (Meanwhile improving [GŠV15]?)
- New gadgets from graph to hypergraph?
- Analyse asymptotically when, say, $\Delta \gtrsim q^{0.51K}$?

Algorithmic side: close the gap between $q^{K/3}$ and $q^{K/2}$.

- Which one is the computational transition threshold? (We guess $1/2$.)



Hypergraph Colouring

Thank you!

arXiv: 2107.05486