

10.2: Sequences

Theorem 10.1: Limits of Sequences from Limits of Functions

Suppose f is a function such that $f(n) = a_n$, for positive integers n . If

$\lim_{x \rightarrow \infty} f(x) = L$, then the limit of the sequence $\{a_n\}$ is also L , where L may be $\pm\infty$.

Example. Determine if the following sequences converge or diverge. If the sequence converges, find its limit.

$$\left\{ e^{2n/(n+2)} \right\}_{n=1}^{\infty}$$

$$\left\{ \frac{(-1)^n}{n} \right\}_{n=1}^{\infty}$$

$$\left\{ \frac{\arctan(n)}{n} \right\}_{n=1}^{\infty}$$

$$\left\{ \frac{e^{-n}}{42 \sin(e^{-n})} \right\}_{n=1}^{\infty}$$

10.2: Limit Laws for Sequences

Assume the sequences $\{a_n\}$ and $\{b_n\}$ have limits A and B , respectively. Then

1. $\lim_{n \rightarrow \infty} (a_n \pm b_n) = A \pm B$
2. $\lim_{n \rightarrow \infty} ca_n = cA$, where c is a real number
3. $\lim_{n \rightarrow \infty} a_nb_n = AB$
4. $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{A}{B}$, provided $B \neq 0$.

Example. Consider the sequences $\{a_n\}$, $\{b_n\}$, $\{c_n\}$, and $\{d_n\}$ where

$$a = \frac{1}{n}, \quad b_n = n, \quad c_n = e^n, \quad \text{and } d_n = \sqrt{n}.$$

Compute the following limits.

A. $\lim_{n \rightarrow \infty} a_n$

B. $\lim_{n \rightarrow \infty} b_n$

C. $\lim_{n \rightarrow \infty} c_n$

D. $\lim_{n \rightarrow \infty} d_n$

E. $\lim_{n \rightarrow \infty} a_nb_n$

F. $\lim_{n \rightarrow \infty} a_nc_n$

G. $\lim_{n \rightarrow \infty} a_nd_n$

True or False: If for some sequence $\{a_n\}$ and $\{b_n\}$, $\lim_{n \rightarrow \infty} a_n = 0$ and $\lim_{n \rightarrow \infty} b_n = \infty$, then $\lim_{n \rightarrow \infty} a_nb_n = 0$.

Definition. (Terminology for Sequences)

- $\{a_n\}$ is **increasing** if $a_{n+1} > a_n$
- $\{a_n\}$ is **nondecreasing** if $a_{n+1} \geq a_n$
- $\{a_n\}$ is **decreasing** if $a_{n+1} < a_n$
- $\{a_n\}$ is **nonincreasing** if $a_{n+1} \leq a_n$
- $\{a_n\}$ is **monotonic** if it is either nonincreasing or nondecreasing (it moves in one direction)
- $\{a_n\}$ is **bounded above** if there is a number M such that $a_n \leq M$, for all relevant values of n
- $\{a_n\}$ is **bounded below** if there is a number N such that $a_n \geq N$, for all relevant values of n .
- If $\{a_n\}$ is bounded above and bounded below, then we say that $\{a_n\}$ is a **bounded** sequence.


Example. Consider the sequence $\{-n^2\}_{n=1}^{\infty}$. What can we say about this sequence?

Theorem 10.3: Geometric Sequences

Let r be a real number. Then

$$\lim_{n \rightarrow \infty} r^n = \begin{cases} 0 & \text{if } |r| < 1 \\ 1 & \text{if } r = 1 \\ \text{does not exist} & \text{if } r \leq -1 \text{ or } r > 1. \end{cases}$$

If $r > 0$, then $\{r^n\}$ is a monotonic sequence. If $r < 0$, then $\{r^n\}$ oscillates.



Example. Determine if the following sequences converge

$$\left\{ \frac{3^{n+1} + 3}{3^n} \right\}$$

$$\{2^{n+1}3^{-n}\}$$

$$\left\{ \frac{(-1)^n}{2^n} \right\}$$

$$\left\{ \frac{75^{n-1}}{99^n} + \frac{5^n \sin(n)}{8^n} \right\}$$

Theorem 10.4: Squeeze Theorem for Sequences

Let $\{a_n\}$, $\{b_n\}$, and $\{c_n\}$ be sequences with $a_n \leq b_n \leq c_n$, for all integers n greater than some index N . If $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} c_n = L$, then $\lim_{n \rightarrow \infty} b_n = L$.

Example. Find the limit of the sequence $b_n = \frac{9 \cos(n)}{n^2 + 1}$.

Theorem 10.5: Bounded Monotonic Sequence

A bounded monotonic sequence converges.

Theorem 10.6: Growth Rates of Sequences

The following sequences are ordered according to increasing growth rates as $n \rightarrow \infty$; that is, if $\{a_n\}$ appears before $\{b_n\}$ in the list, then $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = 0$ and $\lim_{n \rightarrow \infty} \frac{b_n}{a_n} = \infty$:

$$\{(\ln n)^q\} \ll \{n^p\} \ll \{n^p(\ln n)^r\} \ll \{n^{p+s}\} \ll \{b^n\} \ll \{n!\} \ll \{n^n\}$$

Example. Use growth rates to determine which of the following sequences converge.

$$\left\{ \frac{\ln(n^{10})}{0.00001n} \right\}$$

$$\left\{ \frac{n^8 \ln(n)}{n^{8.001}} \right\}$$

$$\left\{ \frac{n!}{10^n} \right\}$$

Definition. (Limit of a Sequence)

The sequence $\{a_n\}$ converges to L provided the terms of a_n can be made arbitrarily close to L by taking n sufficiently large. More precisely, $\{a_n\}$ has the unique limit L if, given any $\varepsilon > 0$, it is possible to find a positive integer N (depending only on ε) such that

$$|a_n - L| < \varepsilon \quad \text{whenever } n > N.$$

If the **limit of a sequence** is L , we say the sequence **converges** to L , written

$$\lim_{n \rightarrow \infty} a_n = L.$$

A sequence that does not converge is said to **diverge**.