2.2 Definitions of Limits


Definition (Briggs). Suppose the function f is defined for all x near a except possibly at a. If f(x) is arbitrarily close to L (as close to L as we like) for all x sufficiently close (but not equal) to a, we write

$$\lim_{x \to a} f(x) = L$$

and say the limit of f(x) as x approaches a equals L.

Note-Most of the time, we can think of the limit as the value of the function if it could be evaluated at a specific point.

Example. Using the graph of f, determine the following values:



•
$$f(2)$$
 and $\lim_{x\to 2} f(x)$

•
$$f(3)$$
 and $\lim_{x\to 3} f(x)$

Definition. (Briggs)


$$f(1) = 2$$

$$\lim_{\chi \to 1} f(\chi) = 1$$

$$f(2) = 5$$

$$\lim_{\chi \to 2} f(\chi) = 3$$

$$f(3) DNE$$



x2 3 5 6 1 4

1. Right-sided limit Suppose f is defined for all x near a with x > a. If f(x) is arbitrarily close to L for all x sufficiently close to a with x > a, we write


$$\lim_{x \to a^+} f(x) = L$$

and say the limit of f(x) as x approaches a from the right equals L.

2. Left-sided limit Suppose f is defined for all x near a with x < a. If f(x) is arbitrarily close to L for all x sufficiently close to a with x < a, we write


$$\lim_{x \to a^{-}} f(x) = L$$

and say the limit of f(x) as x approaches a from the left equals L.



Example. For $f(x) = \frac{x^3 - 8}{4(x - 2)}$, find

- $\lim_{x \to 2^+} f(x) = 3$
- $\bullet \lim_{x \to 2^{-}} f(x) > 3$



Definition. (Briggs) Relationship Between One-Sided and Two-Sided Limits Assume f is defined for all x near a except possibly at a. Then $\lim_{x\to a} f(x) = L$ if and only if $\lim_{x\to a^+} f(x) = L$ and $\lim_{x\to a^-} f(x) = L$.

Example. For f(x) above, is $\lim_{x\to 2} f(x)$ defined? If so, what is it? What is f(2)?

$$\lim_{x\to 2^+} f(x) = \lim_{x\to 2^+} f(x) = 3$$

Example. Consider the graph of


$$g(x) = \frac{2x^2 - 6x + 4}{|x - 1|} = \begin{cases} -2(x - 2) & x < 1\\ 2(x - 2) & x > 1 \end{cases}$$

Find

$$\bullet \lim_{x \to 1^{-}} g(x) = \mathbf{Z}$$

$$\bullet \lim_{x \to 1^+} g(x) = \frown \ \ \$$

$$ullet \lim_{x o 1} g(x)$$
 DNE because




$$\lim_{x\to 1^-} g(x) \neq \lim_{x\to 1^+} g(x)$$

Fall 2018 Class notes

Example. Consider the function

$$h(x) = \frac{x^2 - 81}{2x + 18} = \frac{(x-9)(x+9)}{2(x+9)} = \frac{x-9}{2}, x \neq -9$$

What does this function look like? What is h(-9)? What is $\lim_{x\to -0} h(x)$?




$$\lim_{\chi \to -9} \frac{\chi^{2}-8!}{2\chi+18} = \lim_{\chi \to -9} \frac{\chi-9}{2}$$

$$= \frac{-9-9}{2} = -9$$

Example. The ceiling function is

$$j(x) = \lceil x \rceil$$

where [x] returns the smallest integer greater than or equal to x. In other words, the ceiling function always rounds up. Find the following:



$$\bullet \lim_{x \to 1^+} j(x) = 2$$


•
$$\lim_{x \to 1} j(x)$$
 DNE

• j(1) = 1

•
$$\lim_{x \to 1.5^{-}} j(x) = 2$$

$$\bullet \quad \lim_{x \to 1.5} j(x) = 2$$

 $\lim_{x \to 1.5^{-}} j(x) = 2 \qquad \bullet \quad \lim_{x \to 1.5^{+}} j(x) = 2 \qquad \bullet \quad \lim_{x \to 1.5} j(x) = 2 \qquad \bullet \quad j(1.5) = 2$




Example. Consider the function

$$h(x) = \cos\left(\frac{1}{x}\right)$$

What is $\lim_{x\to 0} h(x)$?

Consider $x = 1/(n\pi)$. As $n \to \infty, x \to 0$, then,

$$\cos\left(\frac{1}{x}\right) = \cos\left(n\pi\right) = \begin{cases} 1 & \text{if } n \text{ is even} \\ -1 & \text{if } n \text{ is odd} \end{cases}$$



Example. Graph an example with the following characteristics:

$$\lim_{x \to -2^{-}} f(x) = -4 \qquad \lim_{x \to -2^{+}} f(x) = 2 \qquad f(-2) = 0$$

$$\lim_{x \to 4} f(x) = 2 \qquad f(4) \text{ DNE}$$

$$\lim_{x \to 8} f(x) = -2 \qquad f(8) = -2$$

