5.2: Logarithmic Functions and Their Properties

Definition.

For $a > 0$ and $a \neq 1$, the **logarithmic function**

$$y = \log_a(x) \qquad \text{(logarithmic form)}$$

has domain $x > 0$, base a, and is defined by

$$a^y = x \qquad \text{(exponential form)}$$

Example. Rewrite the following in exponential form

$4 = \log_2(16)$

$5 = \log_{10}(100,000)$

$\frac{1}{2} = \log_{16}(4)$

$-4 = \log_3\left(\frac{1}{81}\right)$

$-\frac{1}{4} = \log_{625}\left(\frac{1}{5}\right)$

$-\frac{5}{3} = \log_{\frac{1}{8}}(32)$

Example. Simplify the following:

$\log_3(9)$

$\log_4(2)$

Example. Solve the following:

$\log_5(x) = 4$

$\log_8(x) = 1$

$\log_{81}(x) = -\frac{1}{4}$

$\log_{10}(x+4) = 3$

$$\boxed{\begin{array}{ll}\textbf{Common logarithms:} & \log(x) = \log_{10}(x) \\ \textbf{Natural logarithms:} & \ln(x) = \log_e(x)\end{array}}$$

Example (The Rule of 70)**.** If $\$P$ is invested for t years at interest rate r, compounded continuously, then the future value of the investment is given by

$$S = Pe^{rt}.$$

Find the value of t when the investment doubles.

Change of base formula:
If $a > 0$, $b > 0$ with $a \neq 1$ and $b \neq 1$, then

$$\log_b(x) = \frac{\log_a(x)}{\log_a(b)}.$$

Note: This works for any valid base!

$$\text{Base } e: \quad \log_b(x) = \frac{\ln(x)}{\ln(b)} \qquad\qquad \text{Base } 10: \quad \log_b(x) = \frac{\log(x)}{\log(b)}$$

Example. Solve the following

$3^x = 10$

$6.5^x = 5$

Example. Fill in the tables below and graph a^x and $\log_a(x)$ on the same axes.

x	$y = a^x$
-2	
-1	
0	
1	
2	

x	$y = \log_a(x)$
-2	
-1	
0	
1	
2	



Try this!

Example. Graph $\log(-x)$



Example. Graph $\ln(x)$



Example. Graph $-\log_2(-x)$



Example. Evaluate the following:

$f(x) = \ln(x); \quad f\left(e^{-3x}\right)$

$f(x) = 5^x; \quad f(\log_5(10))$

Properties of exponents and logarithms: Assume $a > 0$:

$a^y = x$	$\log_a(x) = y$
$a^1 = a$	$\log_a(a) = 1$
$a^0 = 1$	$\log_a(1) = 0$
$a^x a^y = a^{x+y}$	$\log_a(xy) = \log_a(x) + \log_a(y)$
$\dfrac{a^x}{a^y} = a^{x-y}$	$\log_a\left(\dfrac{x}{y}\right) = \log_a(x) - \log_a(y)$
$a^{xy} = (a^x)^y$	$\log_a(x^y) = y\log_a(x)$
$a^{\log_a(x)} = x$	$\log_a(a^x) = x$