# Math 121 Class notes Fall 2024

Peter Westerbaan

Last updated: November 8, 2024

# **Table Of Contents**

1.1: Solutions of Linear Equations and Inequalities in One Variable
1.3: Linear Functions
1.4: Graphs and Graphing Utilities
1.5: Solutions of Systems of Linear Equations
1.6: Applications of Functions in Business and Economics
4.1: Linear Inequalities in Two Variables
4.2: Linear Programming: Graphical Methods
5.1: Exponential Functions
5.2: Logarithmic Functions and Their Properties
5.3: Equations and Applications with Exponential and Logarithmic Functions
6.1: Simple Interest
6.2: Compound Interest
6.3: Future Values of Annuities

# 1.1: Solutions of Linear Equations and Inequalities in One Variable

### Definition.

A function f is a special relation between x and y such that each input x results in at most one y. The symbol f(x) is read "f of x" and is called the **value of** f **at** x

**Example.** Let  $f(x) = \frac{x^2}{2} + x$ . Evaluate the following:

$$f(1) f(\frac{1}{2})$$

$$f(-2) f(0)$$

## Composite Functions:

Let f and g be functions of x. Then, the **composite functions** g of f (denoted  $g \circ f$ ) and f of g (denoted  $f \circ g$ ) are defined as:

$$(g \circ f)(x) = g(f(x))$$

$$(f \circ g)(x) = f(g(x))$$

**Example.** Let g(x) = x - 1. Find:

$$(g \circ f)(x)$$

$$(f \circ g)(x)$$

## **Operations with Functions:**

Let f and g be functions of x and define the following:

Sum (f+g)(x) = f(x) + g(x) Difference (f-g)(x) = f(x) - g(x) Product  $(f \cdot g)(x) = f(x) \cdot g(x)$  Quotient  $(\frac{f}{g})(x) = \frac{f(x)}{g(x)} \text{ if } g(x) \neq 0$ 

### Definition.

An **expression** is a meaningful string of numbers, variables and operations:

$$3x-2$$

An **equation** is a statement that two quantities or algebraic expressions are equal:

$$3x - 2 = 7$$

A solution is a value of the variable that makes the equation true:

$$3(3) - 2 = 7$$
  
 $9 - 2 = 7$   
 $7 = 7$ 

A **solution set** is the set of ALL possible solutions of an equation:

3x - 2 = 7 only has the solution x = 3,

2(x-1) = 2x - 2 is true for all possible values of x.

## Properties of Equality:

Substitution Property: The equation formed by substituting one expression for an equal expression is equivalent to the original equation:

$$3(x-3) - \frac{1}{2}(4x-18) = 4$$
$$3x - 9 - 2x + 9 = 4$$
$$x = 4$$

**Addition Property:** The equation formed by adding the same quantity to both sides of an equation is equivalent to the original equation:

$$x-4=6$$
  $x+5=12$   $x-4+4=6+4$   $x+5+(-5)=12+(-5)$   $x=7$ 

Multiplication Property: The equation formed by multiplying both sides of an equation by the same *nonzero* quantity is equivalent to the original equation:

$$\frac{1}{3}x = 6$$

$$3\left(\frac{1}{3}x\right) = 3(6)$$

$$x = 18$$

$$5x = 20$$

$$\frac{5x}{5} = \frac{20}{5}$$

$$x = 4$$

# Solving a linear equation:

Using the properties of equality above, we can solve any linear equation in 1 variable:

**Example.** Solve 
$$\frac{3x}{4} + 3 = \frac{x-1}{3}$$

- 1. Eliminate fractions:
- 2. Remove/evaluate parenthesis:
- 3. Use addition property to isolate the variable to one side:
- 4. Use multiplication property to isolate variable:
- 5. Verify solution via substitution:

$$12\left(\frac{3x}{4} + 3\right) = 12\left(\frac{x-1}{3}\right)$$

$$9x + 36 = 4x - 4$$

$$9x + 36 - 36 - 4x = 4x - 4 - 36 - 4x$$

$$\frac{5x}{5} = \frac{-40}{5}$$

$$\underbrace{\frac{3(-8)}{4} + 3}_{-6+3=-3} \stackrel{?}{=} \underbrace{\frac{(-8)-1}{3}}_{\frac{-9}{3}=-3}$$

**Example.** Solve the following:

$$\frac{3x+1}{2} = \frac{x}{3} - 3$$

$$\frac{2x-1}{x-3} = 4 + \frac{5}{x-3}$$

**Example.** Solve -2x + 6y = 4 for y



**Example.** Suppose that the relationship between a firm's profit, P, and the number of items sold, x, can be described by the equation

$$5x - 4P = 1200$$

a) How many units must be produced and sold for the firm to make a profit of \$150?

b) Solve this equation for P in terms of x. Then, find the profit when 240 units are sold.

#### Definition.

An **inequality** is a statement that one quantity is greater than (or less than) another quantity.

### Properties of Inequalities

Substitution Property: The inequality formed by substituting one expression for an equal expression is equivalent to the original inequality:

$$5x - 4x + 2 < 6$$
  
  $x < 4 \implies$  The solution set is  $\{x : x < 6\}$ 

**Addition Property:** The inequality formed by adding the same quantity to both sides of an inequality is equivalent to the original inequality:

$$x-4 < 6$$
  $x+5 \ge 12$   $x-4+4 < 6+4$   $x+5+(-5) \ge 12+(-5)$   $x < 10$   $x \ge 7$ 

**Multiplication Property** The inequality formed by multiplying both sides of an inequality by the same *positive* quantity is equivalent to the original inequality. The direction of the inequality is flipped when multiplying by a *negative* quantity:

$$\frac{1}{3}x > 6 5x - 5 + 5 \le 6x + 20 + 5$$

$$\frac{3}{3}\left(\frac{1}{3}x\right) > \frac{3}{3}(6) \frac{-x}{-1} \le \frac{25}{-1}$$

$$x > 18 x \ge -25$$

# Example. Solve

$$-x + 8 \le 2x - 4$$

first by gathering the x variable on the left, then again on the right. See that the multiplication property holds in both cases. Plot the solution set on a numberline.





**Example.** Plot the following inequalities:

$$x \leq 2$$

$$x > -3$$





### 1.3: Linear Functions

### Definition.

A linear function is a function of the form

$$y = f(x) = ax + b$$

where a and b are constants.

Example. y = -2x + 8



A linear function can be uniquely determined using only two distinct points.

## Definition.

The point(s) where a graph intersects the axes are called intercepts. The x-coordinate of the point where the function intersects the x-axis is called the x-intercepts. The y-coordinate of the point where the function intersects the x-axis is called the y-intercepts.

- To solve for the *y*-intercept:
  - Set x = 0,
  - Solve for y.

- To solve for the *x*-intercept:
  - Set y = 0,
  - Solve for x.

**Example.** Find the intercepts and graph the following lines:

$$3x + 2y = 12$$

$$x = 4y$$





### Definition.

If a nonvertical line passes through the points  $P_1(x_1, y_1)$  and  $P_2(x_2, y_2)$ , its slope, denoted by m, is found using

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{\Delta y}{\Delta x}$$

 $\Delta y$  is "delta y", and represents the change in y  $\Delta x$  is "delta x", and represents the change in x

*Note:* The slope of a vertical line is undefined.

**Example.** Find the slope of the line passing through the points (-2,1) and (5,3).

### *Note:*

- Two distinct nonvertical lines are *parallel* if and only if their slopes are *equal*.
- Two distinct nonvertical lines are *perpendicular* if and only if their slopes are *negative reciprocals*:
  - e.g. If  $\ell_1$  has a nonzero slope m, then  $\ell_2$  is perpendicular if its slope is -1/m.

# Point-slope form

### Definition.

The equation of the line passing through the point  $(x_1, y_1)$  with slope m can be written in the point-slope form:

$$y - y_1 = m(x - x_1)$$

**Example.** Find the equation of each line that passes through the point (-3,4) and has

a slope of 
$$m = \frac{1}{4}$$

the point (-2,1) on the line

a slope of zero (horizontal)

an undefined slope (vertical)

# Slope-intercept form

### Definition.

The slope-intercept form of the equation of a line with slope m and y-intercept b is

$$y = mx + b$$

**Example** (Example 7, p.82). The population of U.S. males, y (in thousands), projected from 2015 to 2060 can be modeled by

$$y = 1125.9x + 142,960$$

where x is the number of years after 2000.

• Find the slope and y-intercept of the graph of this function.

• What does the y-intercept tell us abut the population of U.S. males?

• Interpret the slope as a rate of change.

# Forms of Linear Equations

General form: Ax + By = C

Point-slope form:  $y - y_1 = m(x - x_1)$ 

Slope-intercept form: y = mx + b

Vertical line: x = a

Horizontal line: y = b

# 1.4: Graphs and Graphing Utilities

As graphing calculators are *not* required for this course, we will use Desmos:

desmos.com/calculator

**Example.** For a certain city, the cost C of obtaining drinking water with p percent impurities (by volume) is given by

$$C = \frac{120,000}{p} - 1200$$

The equation for C requires that  $p \neq 0$ , and because p is the percent impurities, we know 0 . Use the restriction on <math>p and a graphing calculator to obtain an accurate graph of the equation.





# 1.5: Solutions of Systems of Linear Equations

$$+ + + + = 18$$
 $+ + + = 14$ 
 $+ + = 2$ 
 $+ + + = 2$ 
 $+ + + = 2$ 

### Definition.

A system of equations is 2 (or more) equations. The ordered pairs (x, y) that satisfies *all* equations in the system are the solutions of the system.

When solving a system of linear equations, there are three possible outcomes:

- 1. No solution (*Inconsistent*),
- 2. Exactly one solution,
- 3. Infinitely many solutions (Dependent).

**Example.** Use graphing to find the solutions to the following systems

$$2x - 6y = -6$$
$$-x + 3y = -3$$



$$4x + 3y = 11$$
$$2x - 5y = -1$$



$$-4x + 3y = -2$$
$$8x - 6y = 4$$



# Equivalent systems result when

- 1. One expression is replaced by an equivalent expression.
- 2. Two equations are interchanged.
- 3. A multiple of one equation is added to another equation.
- 4. An equation is multiplied by a nonzero constant.

### **Substitution Method**

**Example.** Solve the system  $\begin{cases} 2x + 3y = 4 \\ x - 2y = 3 \end{cases}$ 

- 1. Solve one equation for either one of the variables in terms of the other.
- 2. Substitute this expression into the other equation to give the equation in one unknown.
- 3. Solve this equation for the unknown.
- 4. Substitute solution into the equation in Step 1.
- 5. Check the solution (x, y).

$$x = 2y + 3$$

$$2(2y+3) + 3y = 4$$

$$4y + 6 + 3y = 4$$
$$7y = -2 \Rightarrow y = -\frac{2}{7}$$

$$x = 2\left(-\frac{2}{7}\right) + 3 \Rightarrow x = \frac{17}{7}$$

$$2\left(\frac{17}{7}\right) + 3\left(-\frac{2}{7}\right) = 4$$
$$\left(\frac{17}{7}\right) - 2\left(-\frac{2}{7}\right) = 3$$

**Example.** Use the substitution method to solve the system

$$4x + 5y = 18 \tag{1}$$

$$3x - 9y = -12 (2)$$

### Elimination Method

**Example.** Solve the system  $\begin{cases} 2x - 5y = 4 \\ x + 2y = 3 \end{cases}$ 

- 1. Multiply one or both equations by a nonzero number so the coefficients of one of the variables may cancel.
- 2. Add or subtract the equations to eliminate one of the variables.
- 3. Solve for the remaining variable.
- 4. Substitute solution in one of the original equations and solve for the other variable.
- 5. Check the solution (x, y)

$$\Rightarrow \begin{cases} 2x - 5y = 4 \\ -2x - 4y = -6 \end{cases}$$

$$0x - 9y = -2$$

$$\Rightarrow y = \frac{2}{9}$$

$$2x - 5\left(\frac{2}{9}\right) = 4 \implies x = \frac{23}{9}$$

$$2\left(\frac{23}{9}\right) - 5\left(\frac{2}{9}\right) = 4$$
$$\left(\frac{23}{9}\right) + 2\left(\frac{2}{9}\right) = 3$$

**Example.** Use the elimination method to solve the following systems:

$$2x - 6y = -6$$
$$-x + 3y = -3$$

$$4x + 3y = 11$$
$$2x - 5y = -1$$

$$-4x + 3y = -2$$
$$8x - 6y = 4$$

**Example.** A woman has \$500,000 invested in two rental properties. One yields an annual return of 10% on her investment, and the other returns 12% per year on her investment. Her total annual return from the two investments is \$53,000. Let x represent the amount of the 10% investment and y represent the amount of the 12% investment.

- Write an equation that states that the sum of investments is \$500,000.
- What is the annual return on the 10% investment? What about the 12% investment?
- Write an equation that states the sum of the annual return is \$53,000.
- Solve these two equations simultaneously to find how much is invested in each property.

**Example.** A nurse has two solutions that contain different concentrations of a certain medication. One is a 12.5% concentration, and the other is a 5% concentration. How many cubic centimeters of each should she mix to obtain 20 cubic centimeters of an 8% concentration?

**Example.** Using U.S. Bureau of Labor Statistics data for selected years from 1950 and projected to 2050, the number of men M and women W in the workforce (both in millions) can be modeled by the functions

$$M(t) = 0.591t + 37.3$$
 and  $W(t) = 0.786t + 13.1$ 

where t is the number of years after 1940. Find the year these functions predict that there will be equal numbers of men and women in the U.S. workforce.

23



# 1.6: Applications of Functions in Business and Economics

#### Definition.

**Profit** is the difference between the revenue and total cost:

$$P(x) = R(x) - C(x)$$

where

P(x) = profit from sale of x units,

R(x) = total revenue from sale of x units,

C(x) = total cost from production and sale of x units.

*Note*: In general, the symbols used in economics are  $\pi$ , TR and TC respectively.

In general, total revenue is

Revenue = (price per unit)(number of units)

The **total cost** is composed of fixed cost and variable cost:

- Fixed costs (FC) remain constant regardless of the number of units produced.
- Variable costs (VC) are directly related to the number of units produced.

The total cost is given by

Cost = variable costs + fixed costs

**Example.** Suppose a firm manufactures MP3 players and sells them for \$50 each. The costs incurred in the production and sale of the MP3 players are \$200,000 plus \$10 for each player produced and sold. Write the profit function for the production and sale of x players.







**Example.** The ABC company produces widgets which sell at \$25 each. ABC can produce 30 widgets at a total cost of \$2,050, and 180 widgets at a total cost of \$4,300. Find the revenue, cost, and profit functions.

# Definition. (Marginals)

The

- marginal profit  $(\overline{MP})$  is the rate of change in profit...
- marginal cost  $(\overline{MC})$  is the rate of change in costs...
- marginal revenue  $(\overline{MR})$  is the rate of change in revenue...

with respect to the number of units produced and sold. When these functions are linear, the marginals are given by the slope of their respective function.

**Example.** A manufacturer sells widgets for \$10 per unit. The manufacturer's variable costs are \$2.50 per unit, and the total cost of 100 units is \$1,450.

• Find the profit function. What are the marginal revenue, cost and profit?

• Find the break-even point (where R(x) = C(x)). What happens if we sell more or less than the break-even point?





### Definition.

- Market equilibrium occurs when the quantity of a commodity demanded is equal to the quantity supplied.
- The **law of demand** states that the quantity demanded will decrease as the price increases
- The **law of supply** states that the quantity supplied will increase as the price increases

**Example.** Below is a graph containing a supply and demand curve. Find the market equilibrium.



**Example.** Find the market equilibrium point for the following demand and supply functions:

Supply: 
$$p = 2q + 170$$

Demand: 
$$p = -5q + 450$$

**Example.** Using the supply and demand functions above, modify the supply function to include a \$14 tax per unit sold, then find the new market equilibrium point.

**Example.** Retailers will buy 45 Wi-Fi routers from a wholesaler if the price is \$10 each but only 20 if the price is \$60. The wholesaler will supply 56 routers at \$42 each and 70 at \$50 each. Assuming that the supply and demand functions are linear, find the market equilibrium point.

### 4.1: Linear Inequalities in Two Variables

# Properties of Inequalities

Substitution Property: The inequality formed by substituting one expression for an equal expression is equivalent to the original inequality:

$$5x - 4x + 2 < 6$$
  
  $x < 4 \implies$  The solution set is  $\{x : x < 6\}$ 

**Addition Property:** The inequality formed by adding the same quantity to both sides of an inequality is equivalent to the original inequality:

$$x-4 < 6$$
  $x+5 \ge 12$   $x-4+4 < 6+4$   $x+5+(-5) \ge 12+(-5)$   $x < 10$   $x \ge 7$ 

**Multiplication Property** The inequality formed by multiplying both sides of an inequality by the same *positive* quantity is equivalent to the original inequality. The direction of the inequality is flipped when multiplying by a *negative* quantity:

$$\frac{1}{3}x > 6 5x - 5 + 5 \le 6x + 20 + 5$$

$$3\left(\frac{1}{3}x\right) > 3(6) \frac{-x}{-1} \le \frac{25}{-1}$$

$$x > 18 x \ge -25$$

# One Linear Inequality in Two Variables

### Definition.

Consider the inequality y < x:

The line created by this inequality divides the xy-plane into two **half-planes**. We can determine which half-plane is the solution region by selecting any point not on the line as a **test point**.



**Example.** Graph the inequality  $3x - 2y \le 6$ 



# Definition.

A system of inequalities has two or more inequalities in two or more variables. The solution of the system is the intersection of the individual solution sets.

# Example.

$$3x - 2y \ge 4$$
$$x + y - 3 > 0$$



# **Example.** Graph the solution of the system

$$3x-4y \le 12$$
$$2x+5y > 10$$
$$x-8y \ge -16$$



**Example.** CDF Appliances has assembly plants in Atlanta and Fort Worth, where the company produces a variety of kitchen appliances, including a 12-cup coffee maker and a cappuccino machine. The following table shows each factory's assembly capabilities for the two products and the numbers needed to fill orders.

	Atlanta	Fort Worth	Needed
Coffee maker	160/hr	800/hr	At least 64,000
Cappucino machine	$200/\mathrm{hr}$	$200/\mathrm{hr}$	At least 40,000

Write the system of inequalities that describes the number of assembly hours needed at each plant to fill the orders and graph the solution region for the system



**Example.** A farm co-op has 6000 acres available to plant with corn and soybeans. Each acre of corn requires 9 gallons of fertilizer/herbicide and 3/4 hour of labor to harvest. Each acre of soybeans requires 3 gallons of fertilizer/herbicide and 1 hour of labor to harvest. The co-op has available at most 40,500 gallons of fertilizer/herbicide and at most 5250 hours of labor for harvesting. The number of acres of each crop is limited (constrained) by the available resources: land, fertilizer/herbicide, and labor for harvesting. Write the system of inequalities that describes the constraints and graph the solution region for the system.



**Example.** Graph the solution region for the system

$$5x + 2y \le 54$$

$$2x + 4y \le 60$$

$$x \ge 0, \, y \ge 0$$

Then compute the corners of this region.



## 4.2: Linear Programming: Graphical Methods

#### Definition.

Linear programming is an optimization technique that can be used to solve linearly constrained problems:

$$\max F = 3x + y$$
 subject to 
$$x+5y \le 50$$
 
$$2x+ y \le 28$$
 
$$x \ge 0, y \ge 0$$

The **constraints** of a linear program (LP) may be limitations or requirements of the variables. The **objective function** is the function that we wish to optimize (e.g. maximize profit *or* minimize cost).



## Linear programming (graphical method)

- 1. Write the objective function and constraint inequalities from the problem.
- 2. Graph the solution of the constraint system.
- 3. Find the corners of the resulting feasible region.
- 4. Evaluate the objective function at each corner.
- 5. If two corners give the optimal value, then the entire boundary joining these two points optimizes the function.

**Example.** A farm co-op has 6000 acres available on which to plant corn and soybeans. The following table summarizes each crop's requirement for fertilizer/herbicide, harvesting labor hours, and the available amounts of these resources.

	Corn	Soybeans	Available
Fertilizer/herbicide	9 gal/acre	3 gal/acre	40,500 gal
Harvesting labor	3/4  hr/acre	1 hr/acre	$5{,}250~\mathrm{hr}$

Setup the system of inequalities that represents the constraints.



**Example.** Using the linear constraints from above, suppose the co-ops profits per acre are \$240 for corn and \$160 for soybeans. This gives us the following linear program:

$$\max P = 240x + 160y$$
 subject to 
$$x + y \le 6,000$$
 
$$9x + 3y \le 40,500$$
 
$$\frac{3}{4}x + y \le 5,250$$
 
$$x \ge 0, y \ge 0$$

- 1. Find the "corners" of the feasible region
- 2. Evaluate the profit function at the corners

(x, y)	P = 240x + 160y
(0,0)	\$0
(0, 5250)	\$840,000
(3000, 3000)	\$1,200,000
(3750, 2250)	\$1,260,000
(4500, 0)	\$1,080,000



**Example.** Suppose the profits per acre are instead \$300 for corn, and \$100 for soybeans. This gives us the following linear program:

$$\max P = 300x + 100y$$
 subject to 
$$x+ \quad y \leq \quad 6,000$$
 
$$9x+ \quad 3y \leq \quad 40,500$$
 
$$\frac{3}{4}x+ \quad y \leq \quad 5,250$$
 
$$x \geq 0, y \geq \quad 0$$

Evaluate the profit function at the corners. What combination of corn and soy-beans maximizes the profit?

(x, y)	P = 300x + 100y
(0,0)	\$0
(0, 5250)	\$525,000
(3000, 3000)	\$1,200,000
(3750, 2250)	\$1,350,000
(4500, 0)	\$1,350,000



**Example.** Two chemical plants, one at Macon and one at Jonesboro, produce three types of fertilizer: low phosphorus (LP), medium phosphorus (MP), and high phosphorus (HP). At each plant, the fertilizer is produced in a single production run, so the three types are produced in fixed proportions. The Macon plant produces 1 ton of LP, 2 tons of MP, and 3 tons of HP in a single operation and charges \$600 for what is produced in one operation. On the other hand, one operation of the Jonesboro plant produces 1 ton of LP, 5 tons of MP, and 1 ton of HP, and it charges \$1,000 for what it produces in one operation. If a customer needs 100 tons of LP, 260 tons of MP, and 180 tons of HP, how many production runs should be ordered from each plant to minimize costs?

Organize the information from the problem in the following table:

	Macon	Jonesboro	Requirements
Units of LP			
Units of MP			
Units of HP			

What is the objective function?


Write the linear program that we aim to solve below:

**Example.** From above, we get the following linear program

$$\min C = 600x + 1000y$$
 subject to 
$$x + y \ge 100$$
 
$$2x + 5y \ge 260$$
 
$$3x + y \ge 180$$
 
$$x \ge 0, y \ge 0$$

Graph the solution region of the constraints and evaluate the objective function at the corners of the feasible region above.

(x, y)	P = 600x + 1,000y
(0, 180)	\$180,000
(40, 60)	\$84,000
(80, 20)	\$68,000
(130, 0)	\$78,000



## 5.1: Exponential Functions

## **Properties of Exponents:**

• If m is a positive integer, then  $x^m = \underbrace{x \cdot x \dots x}_{m \text{ times}}$ :

$$3^4 = 3 \cdot 3 \cdot 3 \cdot 3 = 81, \quad (-5)^3 = (-5) \cdot (-5) \cdot (-5) = -125$$

• Additivity — If the bases are the same, then  $x^a \cdot x^b = x^{a+b}$  and  $\frac{x^a}{x^b} = x^{a-b}$ :

$$4^3 \cdot 4^2 = 4^{3+2} = 4^5 = 1024,$$
  $\frac{3^{17}}{3^{12}} = 3^{17-12} = 3^5 = 243$ 

• If  $x \neq 0$ , then  $x^0 = 1$ :

$$4^0 = 1,$$
  $(-7)^0 = 1,$   $2024^0 = 1$ 

• Distributive —  $(ab)^m = a^m b^m$  and  $\left(\frac{a}{b}\right)^m = \frac{a^m}{b^m}$ :

$$(3 \cdot 4)^2 = 3^2 \cdot 4^2 = 9 \cdot 16 = 144,$$
  $\left(\frac{4}{5}\right)^2 = \frac{4^2}{5^2} = \frac{16}{25}$ 

• If  $m \neq 0$ , then  $x^{-m} = \frac{1}{x^m}$  and  $x^m = \frac{1}{x^{-m}}$ :

$$2^{-4} = \frac{1}{2^4} = \frac{1}{16},$$
  $\left(\frac{1}{3}\right)^{-2} = 3^2 = 9$ 

• Multiplicity —  $(x^a)^b = x^{ab}$ :

$$\left(3^2\right)^4 = 3^8 = 6561$$

• Fractional exponents —  $x^{1/m} = \sqrt[m]{x}$ :

$$8^{1/3} = \sqrt[3]{8} = 2,$$
  $16^{3/2} = (16^{1/2})^3 = (\sqrt{16})^3 = 4^3 = 64$ 

#### Definition.

An **exponential function** is of the form

$$f(x) = a^x$$

where a > 0 and  $a \neq 1$ .

*Note:* The variable is in the exponent (e.g.  $2^x$  vs  $x^2$ )

**Example.** Suppose a culture of bacteria has the property that each minute, every microorganism splits into two new organisms. The number of microorganisms after x minutes is given by

$$y=2^x$$
.

Fill out the table below, and graph this exponential function (include x < 0).

 $\frac{x \quad y = 2^x}{0}$  1 2 3 4



Example. Graph the exponential function

$$y = \left(\frac{1}{3}\right)$$

$$-10^{\uparrow}$$

$$-8$$

$$-6$$

$$-4$$

2

For an exponential function  $a^x$ , the function is

• increasing if a > 1, and

-2

• decreasing if 0 < a < 1.

**Example.** Graph the exponential function





## **Example. Compound Interest:**

If an initial principal P is invested at a rate r and compounded n times a year, the future value in t years is given by:

$$A = P\left(1 + \frac{r}{n}\right)^{nt}$$

Suppose that \$800 dollars is invested, and is compounded quarterly at a rate of 6%:

$$A = 800(1.015)^{4t}$$

Find the future value after 10 years.

#### Example. Compounded continuously:

A special function that frequently occurs in the context of exponential functions is

$$y = e^x$$

where e = 2.71828... (think irrational number like  $\pi$ ). When an investment is compounded continuously, it's future value is given by

$$A = Pe^{rt}$$

Suppose that we invest \$800, compounded continuously at 6%. Find the future value in 10 years.

## 5.2: Logarithmic Functions and Their Properties

## Definition.

For a > 0 and  $a \neq 1$ , the **logarithmic function** 

$$y = \log_a(x)$$

(logarithmic form)

has domain x > 0, base a, and is defined by

$$a^y = x$$

(exponential form)

Example. Rewrite the following in exponential form

$$4 = \log_2(16)$$

$$5 = \log_{10}(100, 000)$$

$$\frac{1}{2} = \log_{16}(4)$$

$$-4 = \log_3\left(\frac{1}{81}\right)$$

$$-\frac{1}{4} = \log_{625}\left(\frac{1}{5}\right)$$

$$-\frac{5}{3} = \log_{\frac{1}{8}}(32)$$

**Example.** Simplify the following:

$$\log_3(9)$$

$$\log_4(2)$$

**Example.** Solve the following:

$$\log_5(x) = 4$$

$$\log_8(x) = 1$$

$$\log_{81}(x) = -\frac{1}{4}$$

$$\log_{10}(x+4) = 3$$

Common logarithms:  $\log(x) = \log_{10}(x)$ Natural logarithms:  $\ln(x) = \log_e(x)$ 

**Example** (The Rule of 70). If P is invested for t years at interest rate r, compounded continuously, then the future value of the investment is given by

$$S = Pe^{rt}$$
.

Find the value of t when the investment doubles.

Change of base formula:

If a > 0, b > 0 with  $a \neq 1$  and  $b \neq 1$ , then

$$\log_b(x) = \frac{\log_a(x)}{\log_a(b)}.$$

Note: This works for any valid base!

Base 
$$e: \log_b(x) = \frac{\ln(x)}{\ln(b)}$$

Base 10: 
$$\log_b(x) = \frac{\log(x)}{\log(b)}$$

Example. Solve the following


$$3^x = 10$$

$$6.5^x = 5$$

**Example.** Fill in the tables below and graph  $a^x$  and  $\log_a(x)$  on the same axes.


x	$y = a^x$	
<del>-</del> 2		

$$\frac{x \quad y = \log_a(x)}{}$$




Try this!


**Example.** Graph  $\log(-x)$ 



## **Example.** Graph ln(x)



## **Example.** Graph $-\log_2(-x)$



Fall 2024

**Example.** Evaluate the following:

$$f(x) = \ln(x); \quad f(e^{-3x})$$
  $f(x) = 5^x; \quad f(\log_5(10))$ 

## Properties of exponents and logarithms: Assume a > 0:

$$a^{y} = x \qquad \log_{a}(x) = y$$

$$a^{1} = a \qquad \log_{a}(a) = 1$$

$$a^{0} = 1 \qquad \log_{a}(1) = 0$$

$$a^{x}a^{y} = a^{x+y} \qquad \log_{a}(xy) = \log_{a}(x) + \log_{a}(y)$$

$$\frac{a^{x}}{a^{y}} = a^{x-y} \qquad \log_{a}\left(\frac{x}{y}\right) = \log_{a}(x) - \log_{a}(y)$$

$$a^{xy} = (a^{x})^{y} \qquad \log_{a}(x^{y}) = y \log_{a}(x)$$

$$a^{\log_{a}(x)} = x \qquad \log_{a}(a^{x}) = x$$

# 5.3: Equations and Applications with Exponential and Logarithmic Functions

## Solving exponential equations:

**Example.** Solve  $4(25^{2x}) = 312,500$ 

- 1. Isolate the exponential by rewriting the equation with a base raised to a power on one side:  $\frac{4(25^{2x})}{4} = \frac{312,500}{4}$
- 2. Take the logarithm of both sides:  $\ln (25^{2x}) = \ln (78, 125)$
- 3. Use a property of logarithms to remove the variable from the exponent:  $2x \ln{(25)} = \ln{(78, 125)}$
- 4. Solve for the variable:  $x = \frac{\ln{(78, 125)}}{2\ln{(25)}} \approx 1.75$

**Example.** Unless the exponential function uses base e or base 10, it does not matter which logarithm we use. Solve the following exponential equation first using base 10, then using base e:

$$6(4^{3x-2}) = 20$$

**Example.** Suppose the demand function for q thousand units of a certain commodity is given by

$$p = 30\left(3^{-q/2}\right)$$

At what price per unit will the demand equal 4000 units?

How many units, to the nearest thousand units, will be demanded if the price is \$17.31?

<b>Example.</b> A company finds that its daily sales begin to fall after the end of an advertising campaign, and the decline is such that the number of sales is $S = 2000(2^{-0.1x})$ , where $x$ is the number of days after the campaign ends.
How many sales will be made after 10 days after the end of the campaign?
If the company does not want sales to drop below 350 per day, when should it start a new campaign?

**Example.** The population of a certain city was 30,000 in 2000, and 40,500 in 2010. If the formula  $P = P_0 e^{ht}$  applies to the growth of the city's population, what population is predicted for the year 2030?

## Example. The Gompertz equation

$$N = 100(0.03)^{0.2^t}$$

predicts the size of a deer herd on a small island t decades from now.

What is the size of the deer population now (t = 0)?

During what year will the deer population reach or exceed 70?

**Example.** One company's revenue from the sales of computer tablets from 2015 to 2020 can be modeled by the logistic function

$$y = \frac{9.46}{1 + 53.08e^{-1.28x}}$$

where x is the number of years past 2014 and y is in millions of dollars.

Estimate the sales revenue for 2020

During what year will the sales revenue exceed \$4 million?

**Example** (Bonus). Solve the following for x:

$$6^{x-2} = 2^{-3x}$$

## 6.1: Simple Interest

#### Definition.

The **simple interest** I is given by

$$I = Prt$$

where

I = interest (in dollars)

P = principal (in dollars)


r = annual interest rate (as a decimal)

t = time (in years)

**Note:** The time measurements of r and t must agree

From this, the **future value** of simple interest is

$$S = P + I = P + Prt = P(1 + rt)$$



#### Definition.

The **return on investment** (ROI) is the ratio between the gain and cost of an investment:

 $ROI = \frac{\text{Gains on investment}}{\text{Cost of investment}}$ 

The **earned (effective) interest rate** is the equivalent interest rate of the investment when all the fees and dividends are included.

**Example.** Mary Spaulding bought Wind-Gen Electric stock for \$6,125.00. After 6 months, the value of her shares had risen by \$138.00 and dividends totaling \$144.14 had been paid.

Find the return on investment on this investment.

Find the simple interest rate she earned on this investment if she sold the stock at the end of the 6 months.

6.1: Simple Interest 65 Math 121 Class notes

Example.				
To buy a	Treasury bill (T-bill) that What annual simple interest		· -	nust pay
If the bar earn?	nk charges a fee of \$40 to b	uy a T-bill, what is	the actual interest	rate you
6.1: Simple Interest		66	Math 121	Class notes
•				Fall 2024

## 6.2: Compound Interest

**Example.** Suppose you invest \$1,000 at 5% annual interest. With simple interest, you can take 2 approaches:

- 1. Gain interest on *only* your initial investment
- 2. Reinvest the interest gained

Year	Simple interest	Simple interest reinvested
1	\$1,050.00	\$1,050.00
2	\$1,100.00	\$1,102.50
3	\$1,150.00	\$1,157.63
4	\$1,200.00	\$1,215.51
5	1,250.00	$$1,\!276.28$



#### Definition.

Compound interest is a method where the interest for each period is added to the principal before interest is calculated for the next period.

**Example.** Using the example above, derive a formula for the future value of an investment compounded annually.

#### Definition.

When interest is compounded multiple times a year (e.g. quarterly, monthly, etc.), the **nominal interest rate** is the interest rate *per year*.

If P is invested for t years at a nominal interest rate r compounded m times per year, then the **total number of compounding periods** is

$$n = mt$$

the interest rate per compounding period (periodic interest rate) is

$$i = \frac{r}{m}$$

and the future value is

$$S = P(1+i)^n = P\left(1 + \frac{r}{m}\right)^{mt}$$

**Example.** If \$3,000 is invested for 5 years at 9% compounded 4 times a year, how much interest is earned?

**Example.** For the following, identify the annual interest rate, the length in years, the periodic interest rate, and the number of periods:

12% compounded monthly for 7 years

7.2% compounded quarterly for 11 quarters


Frequency	m
Annually	1
Semi-annually	2
Quarterly	4
Monthly	12
Weekly	52
Daily	365

**Example.** Ben and Taylor want to have \$200,000 in Arthur's college fund on his 18th birthday, and they want to know the impact on this goal of having \$10,000 invested at 9.8%, compounded quarterly, on his 1st birthday. To advise Ben and Taylor regarding this, find

the future value of the \$10,000 investment,

the amount of compound interest that the investment earns,

the impact this would have on their goal.



**Example.** Suppose we invest \$1 at a 100% interest rate for 1 year:

$$S = \left(1 + \frac{1.00}{m}\right)^m$$

Compute the future value

Annually

Semi-annually

Monthly

Weekly

Daily

Each minute (m = 525, 600)

#### Definition.

If P is invested for t years at a nominal rate r compounded continuously, then the future value is given by the exponential function

$$S = Pe^{rt}$$

**Example.** Which investment strategy is worth more: \$3,000 for 8 years at

9%, compounded annually

8%, compounded continuously

**Example.** Suppose you invest \$900 at 11.5%, compounded continuously. How long will it take to gain \$700 in interest?

## Definition.

Let r represent the annual (nominal) interest rate for an investment. Then the **annual** percentage yield (APY) is:

Periodic compounding:

$$APY = \left(1 + \frac{r}{m}\right)^m - 1$$

Continuous compounding:

$$APY = e^r - 1$$

#### 6.3: Future Values of Annuities

#### Definition.

- An **annuity** is a financial plan characterized by regular payments (e.g. mortgages, student loans, etc.).
- The sum of all the payments and the interest earned is called the **future value** of the annuity or its future value.
- An **ordinary annuity** or (**annuity immediate**) is an annuity in which payments are made at the *end of each of the equal payment intervals*.

**Example.** Suppose that we invest \$100 at the end of each year for 5 years in an account that pays 10% compounded annually. How much money will you have at the end of the 5 years?

#### Definition.


If R is deposited at the end of each period for n periods in an annuity that earns interest at a rate of i per period, the **future value of the annuity** will be

$$S = R \cdot S_{\overline{n}|i} = R \left\lceil \frac{(1+i)^n - 1}{i} \right\rceil$$

The notation  $S_{\overline{n}|i}$  represents the future value of an ordinary annuity of \$1 per period for n periods with an interest rate of i per period.

regula	nple. Suppose a pair of twins arly make investments of \$2,000 and at age 21:		
	Find the future value if twin a investment accrue compound i		
	Find the future value if twin I the following 36 years.	B waits 8 years before maki	ng regular payments for
6 3. Fut	cure Values of Annuities	75	Math 121 Class notes

Fall 2024



#### Definition.

An **annuity due** differs from an ordinary annuity in that the payments are made at the *beginning of each period*.

If R is deposited at the beginning of each period for n periods in an annuity that earns interest at a rate of i per period, the **future value of the annuity** will be

$$S_{\text{due}} = R \cdot S_{\overline{n}|i}(1+i) = R \left[ \frac{(1+i)^n - 1}{i} \right] (1+i)$$

**Example.** Find the future value of an investment if \$150 is deposited at the beginning of each month for 9 years at an interest rate of 7.2% compounded monthly.

Fall 2024