

## 2.4: Limits

**Example.** Suppose that the position function of a maglev train (in feet) is given by

$$s(t) = 4t^2, \quad (0 \leq t \leq 30)$$

Using the position function, compute the *average* velocity of the train

on the interval  $[t, 2]$

$t$	1.5	1.9	1.99	1.999	1.9999
<hr/>					

on the interval  $[2, t]$

$t$	2.5	2.1	2.01	2.001	2.0001
<hr/>					

What do the tables above suggest about *instantaneous* velocity of the train at  $t = 2$ ?

**Definition. (Limit of a Function)**

The function  $f$  has the **limit**  $L$  as  $x$  approaches  $a$ , written

$$\lim_{x \rightarrow a} f(x) = L$$


if the value of  $f(x)$  can be made as close to the number  $L$  as we please by taking  $x$  sufficiently close to (but not equal to)  $a$ .

**Example.** Using the graph of  $f$ , determine the following values:

$$f(1) \text{ and } \lim_{x \rightarrow 1} f(x)$$

$$f(2) \text{ and } \lim_{x \rightarrow 2} f(x)$$

$$f(3) \text{ and } \lim_{x \rightarrow 3} f(x)$$



**Example.** Find the limit of the following functions at the value specified:

[Graphs](#)

$$f(x) = x^3 \quad \text{at } x = 2$$

$$g(x) = \begin{cases} x + 2, & x \neq 1 \\ 1, & x = 1 \end{cases} \quad \text{at } x = 1$$

$$h(x) = \begin{cases} -1, & x < 0 \\ 1, & x \geq 0 \end{cases} \quad \text{at } x = 0$$

$$j(x) = \frac{1}{(x-1)^2} \quad \text{at } x = 1$$

$$k(x) = 4 \quad \text{at } x = 0$$

### Theorem 1: Properties of Limits

Suppose

$$\lim_{x \rightarrow a} f(x) = L \quad \text{and} \quad \lim_{x \rightarrow a} g(x) = M$$

Then

1.  $\lim_{x \rightarrow a} [f(x)]^r = \left[ \lim_{x \rightarrow a} f(x) \right]^r$  where  $r$  is a positive constant
2.  $\lim_{x \rightarrow a} cf(x) = c \lim_{x \rightarrow a} f(x)$  where  $c$  is a real number
3.  $\lim_{x \rightarrow a} [f(x) \pm g(x)] = \lim_{x \rightarrow a} f(x) \pm \lim_{x \rightarrow a} g(x) = L \pm M$
4.  $\lim_{x \rightarrow a} [f(x)g(x)] = \left[ \lim_{x \rightarrow a} f(x) \right] \left[ \lim_{x \rightarrow a} g(x) \right] = LM$
5.  $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)} = \frac{L}{M}$  provided  $M \neq 0$

**Example.** Use the above theorem to evaluate the following limits:

$$\lim_{x \rightarrow 1} (5x^{3/2} - 2)$$

$$\lim_{x \rightarrow 3} \frac{2x^3 \sqrt{x^2 + 7}}{x + 1}$$

Suppose that  $\lim_{x \rightarrow a} f(x) = 0$  and  $\lim_{x \rightarrow a} g(x) = 0$ . Then

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)}$$

has an **indeterminate form** of  $\frac{0}{0}$ . To evaluate such a limit, we replace the given function with a function that's equivalent everywhere except at  $x = a$ , and then evaluate the limit.

**Example.** Evaluate the following

$$\lim_{t \rightarrow 2} \frac{4t^2 - 16}{t - 2}$$

$$\lim_{h \rightarrow 0} \frac{\sqrt{4 + h} - 2}{h}$$

### Limit of a Function at Infinity

The function  $f$  has the limit  $L$  as  $x$  increases without bound, written

$$\lim_{x \rightarrow \infty} f(x) = L$$

if  $f(x)$  can be made arbitrarily close to  $L$  by taking  $x$  large enough.

The function  $f$  has the limit  $M$  as  $x$  decreases without bound, written

$$\lim_{x \rightarrow -\infty} f(x) = M$$

if  $f(x)$  can be made arbitrarily close to  $M$  by taking  $x$  to be negative and sufficiently large enough in absolute value.

When the above limits exist, the equations  $y = L$  and/or  $y = M$  are called **horizontal asymptotes**.


**Example.** Evaluate the following infinite limits

$$\lim_{x \rightarrow \infty} \frac{2x^2 + 3x - 4}{x^2 - 7x + 1}$$

$$\lim_{x \rightarrow -\infty} \frac{3x^2 + 4}{2x^3}$$

$$\lim_{x \rightarrow \pm\infty} \frac{3x^5 + 2x^3 - 4}{x^4 + 4x^2 - 1}$$

Graph  
 $f(x)$



$$\lim_{x \rightarrow -\infty} \frac{7x^3 - 2}{-x^3 + \sqrt{25x^6 - 4}}$$

$$\lim_{x \rightarrow \infty} \frac{7x^3 - 2}{-x^3 + \sqrt{25x^6 - 4}}$$

**Example.** The company *Custom Office* makes a line of executive desks. It is estimated that the total cost of making  $x$  *Senior Executive Model* desks is

$$C(x) = 100x + 200,000$$

dollars per year. The average cost of making  $x$  desks is given by

$$\overline{C}(x) = \frac{C(x)}{x}$$

Compute  $\lim_{x \rightarrow \infty} \overline{C}(x)$  and interpret the result.

**Theorem 2**

For all  $n > 0$ ,

$$\lim_{x \rightarrow \pm\infty} \frac{1}{x^n} = 0$$

provided that  $\frac{1}{x^n}$  is defined.