
Question			Answer	Marks	AOs	Guidance	
9	(a)	(i)	 One point labelled. eg (1, 12), (2, 6), (3, 4) etc	**B1** **B1dep** **[2]**	**1.1a** **1.1**	General shape for curve with axes as asymptotes One correct point clearly seen	
9	(a)	(ii)	$y = 2x + 8$ drawn (–4, 0) and (0, 8) clearly labelled	**B1** **B1** **[2]**	**1.1a** **1.1**	Line with positive gradient and positive intercept Both intercepts correctly labelled	Allow marks for the line even if the curve is wrong or missing
9	(b)		**DR** $2x + 8 = \dfrac{12}{x}$ $2x^2 + 8x - 12 = 0$ $x = -2 \pm \sqrt{10}$ When $x = -2 + \sqrt{10}$, $y = 2\left(-2 + \sqrt{10}\right) + 8 = 4 + 2\sqrt{10}$ $x = -2 - \sqrt{10}$, $y = 2\left(-2 - \sqrt{10}\right) + 8 = 4 - 2\sqrt{10}$ So coordinates are $\left(-2 + \sqrt{10}, 4 + 2\sqrt{10}\right)$ and $\left(-2 - \sqrt{10}, 4 - 2\sqrt{10}\right)$	**M1** **M1** **A1** **M1** **A1** **[5]**	**1.1a** **1.1a** **1.1** **1.1** **1.1**	Attempt to eliminate one variable Attempt to rewrite into the form $ax^2 + bx + c = 0$ and attempt to solve Both roots seen (allow BC if both **M** marks awarded) Substituting their roots into either equation Both coordinates correct and exact	Allow **SC1** for (–5.16, –2.32) and (1.16, 10.3) BC www If substitution into the curve used, allow $\dfrac{12}{-2 \pm \sqrt{10}}$ for **M1A0** if denominator not rationalised