Question			Answer	Marks	AOs	Guidance	
5	(a)		$\frac{1}{8} \times \frac{1}{7}$	M1	1.1a	or $\frac{6!}{8!}$ oe	
			$\frac{1}{56}$	A1	1.1	CAO accept 0.0179	0.017857142
				[2]			
5	(b)		${}^5C_3$ or ${}^5C_2 \times {}^3C_1$ seen [10, 30]	M1	1.1a	OR 3W 10 ways 2W1M 10x3 = 30 ways 1W2M 5x3 = 15 ways 3M 1 way So $\frac{10+30}{10+30+15+1} = \frac{5}{7}$	Allow M1 if the 2 cases are identified and added, whatever the calculation. Allow method by subtraction from 1. This question can be done by tree diagram, with similar working.
			${}^8C_3$ is denominator [56] {equiv to ${}^8C_5$}	B1	1.1		
			P(3, 0) + P(2, 1) attempted	M1	2.1		
			$\frac{5}{7}$	A1	1.1		
				[4]			
			alternatively $\frac{5}{8} \times \frac{4}{7} \times \frac{3}{6}$ or $3 \times \frac{5}{8} \times \frac{4}{7} \times \frac{3}{6}$ seen	M1	1.1a	$\frac{60}{336}$ or $\frac{5}{28}$ or $\frac{180}{336}$ or $\frac{15}{28}$	Allow both M marks for all denoms 8 – eg $\frac{5}{8} \times \frac{4}{8} \times \frac{3}{8} = \frac{60}{512}$ etc leading to $\frac{15}{32}$
			both correct including ×3	A1	1.1		
			P(3, 0) + P(2, 1) attempted	M1	2.1	Allow even if ×3 missing	
			$\frac{5}{7}$	A1			
				[4]	1.1		