Question	Answer	Marks	AO	Guidance	
13	**DR** $6x^{-\frac{3}{2}} + 2x^{-2}$ seen	**M1**	**3.1a**	allow one slip in one coefficient	
	$\frac{6x^{-\frac{1}{2}}}{-\frac{1}{2}} + \frac{2x^{-1}}{-1}$ **oe**	**M1***	**1.1**	Two correct indices from their $6x^{-\frac{3}{2}} + 2x^{-2}$	ft their $6x^{-\frac{3}{2}} + 2x^{-2}$ for the method marks condone $+c$
		M1*	**1.1**	Both indices correct and at least one coefficient correct from their $6x^{-\frac{3}{2}} + 2x^{-2}$	
		A1	**2.1**	Both terms fully correct	
	$-12 \times 8^{-\frac{1}{2}} - 2 \times 8^{-1} - (-12 - 2)$ **or** $\frac{-12}{\sqrt{8}} - \frac{2}{8} - (-12 - 2)$	**M1dep***	**1.1**	Must show substitution of corrects limits into their integrand	
	$\frac{55}{4} - 3\sqrt{2}$	**A1**	**1.1**	Accept $\frac{55}{4} - \frac{6}{\sqrt{2}}$ or $\frac{55-12\sqrt{2}}{4}$	
		[6]			