				[1]			
10	**(a)**		to account for seasonal variation in **birds oe** or to account for different weather conditions which affects what **birds** are seen **oe**	**B1**	**2.2b**		
				[1]			
10	**(b)**		positive skew **cao**	**B1**	**1.2**		
				[1]			
10	**(c)**		eg in lower quartile (or lower tail) eg much lower than median (or middle) **oe** eg frequency of 3 and others have frequency of 2 eg distribution is skewed (so median more appropriate)	**B1**	**2.4**	NB mode is 9	
				[1]			
10	**(d)**		median is 19.5	**B1**	**1.1**		
			LQ is between 8th and 9th value or UQ is between 24th and 25th value **soi**	**M1**	**1.1**	allow for sight of 11 or 30	
			IQR = 30 – 11 = 19	**A1**	**1.1**		
				[3]			
10	**(e)**		2015: 32 $\leq$ median $\leq$ 34, much bigger than (their 19.5) in 2019 which supports Ben's first statement **oe** 2015: 31 $\leq$ IQR $\leq$ 34 was (much) bigger than (their 19) in 2019 which does **not** support Ben's second statement **oe**	**B1** **B1**	**2.2b** **2.4**	if **B0B0** allow **SC1** for true – median smaller in 2019 **oe and** false – IQR smaller in 2019 **oe**	
				[2]			

Question			Answer	Marks	AOs	Guidance	
10	(f)		eg other distributions are symmetrical (so could be eg Normal) **oe** eg can't tell how values distributed in each quartile **oe**	**B1**	**2.4**		
				[1]			