13	**(a)**		mean = 1.5 BC	**B1**	**1.1**		
			sd = 1.1, 1.10 or awrt 1.096 BC	**B1**	**1.1**		
				[2]			
13	**(b)**		$n = 10$ and $p = 0.15$	**B1**	**3.3**		
				[1]			
13	**(c)**		<table><tr><td>N</td><td>0</td><td>1</td><td>2</td><td>3</td><td>4</td><td>5 +</td></tr><tr><td>O</td><td>19</td><td>35</td><td>28</td><td>13</td><td>5</td><td>0</td></tr><tr><td>E</td><td>19.7</td><td>34.7</td><td>27.6</td><td>13.0</td><td>4.0</td><td>1.0</td></tr></table>	**M1** **M1** **A1**	**3.4** **3.4** **1.1**	use of B(10, p) FT their p to find at least 2 probabilities; may be unsimplified multiplying their binomial probabilities by 100 all correct to 1 dp or better; must add up to 100	must see at least 2 probabilities for **M1** must see at least 2 expected values calculated for **M1**
				[3]			
13	**(d)**		close match between theoretical (or expected) and observed frequencies so model is a good fit	**B1**	**3.5a**	FT dependent on award of **M1M1** in **(c)** and all frequencies calculated	
				[1]			