13 The function $f(x)$ is defined by $f(x) = \sqrt[3]{27-8x^3}$. Jenny uses her scientific calculator to create a table of values for $f(x)$ and $f'(x)$.

x	$f(x)$	$f'(x)$
0	3	0
0.25	2.9954	–0.056
0.5	2.9625	–0.228
0.75	2.8694	–0.547
1	2.6684	–1.124
1.25	2.2490	–1.977
1.5	0	ERROR

(i) Use calculus to find an expression for $f'(x)$ and hence explain why the calculator gives an error for $f'(1.5)$. **[3]**

(ii) Find the first three terms of the binomial expansion of $f(x)$. **[3]**

(iii) Jenny integrates the first three terms of the binomial expansion of $f(x)$ to estimate the value of $\int_0^1 \sqrt[3]{27-8x^3}\,dx$. Explain why Jenny's method is valid in this case. (You do not need to evaluate Jenny's approximation.) **[2]**

(iv) Use the trapezium rule with 4 strips to obtain an estimate for $\int_0^1 \sqrt[3]{27-8x^3}\,dx$. **[3]**

The calculator gives 2.921 174 38 for $\int_0^1 \sqrt[3]{27-8x^3}\,dx$. The graph of $y = f(x)$ is shown in Fig. 13.




Fig. 13

(v) Explain why the trapezium rule gives an underestimate. **[1]**

13(i)	

13(ii)	
13(iii)	

13(iv)

13(v)