- The acceleration due to gravity is denoted by $g\,\mathrm{m\,s^{-2}}$. Unless otherwise instructed, when a numerical value is needed, use $g = 9.8$.

14 The velocity of a car, $v\,\mathrm{m\,s^{-1}}$ at time t seconds, is being modelled. Initially the car has velocity $5\,\mathrm{m\,s^{-1}}$ and it accelerates to $11.4\,\mathrm{m\,s^{-1}}$ in 4 seconds.

In model A, the acceleration is assumed to be uniform.

(i) Find an expression for the velocity of the car at time t using this model. **[3]**

(ii) Explain why this model is not appropriate in the long term. **[1]**

Model A is refined so that the velocity remains constant once the car reaches $17.8\,\mathrm{m\,s^{-1}}$.

(iii) Sketch a velocity-time graph for the motion of the car, making clear the time at which the acceleration changes. **[3]**

(iv) Calculate the displacement of the car in the first 20 seconds according to this refined model. **[3]**

In model B, the velocity of the car is given by

$$v = \begin{cases} 5 + 0.6t^2 - 0.05t^3 & \text{for } 0 \leqslant t \leqslant 8, \\ 17.8 & \text{for } 8 < t \leqslant 20. \end{cases}$$

(v) Show that this model gives an appropriate value for v when $t = 4$. **[1]**

(vi) Explain why the value of the acceleration immediately before the velocity becomes constant is likely to mean that model B is a better model than model A. **[3]**

(vii) Show that model B gives the same value as model A for the displacement at time 20 s. **[3]**

14(i)	
14(ii)	
14(iii)	

14(iv)	
14(v)	

14(vi)	
14(vii)	

Contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.