15 Bottles of Fizzipop nominally contain 330 ml of drink. A consumer affairs researcher collects a random sample of 55 bottles of Fizzipop and records the volume of drink in each bottle.


Summary statistics for the researcher's sample are shown in the table.

n	55
$\sum x$	18535
$\sum x^2$	6247066.6

(a) (i) Calculate the mean volume of drink in a bottle of Fizzipop. **[1]**

(ii) Show that the standard deviation of the volume of drink in a bottle of Fizzipop is 3.78 ml. **[1]**

The researcher uses software to produce a histogram with equal class intervals, which is shown below.



(b) Explain why the researcher decides that the Normal distribution is a suitable model for the volume of drink in a bottle of Fizzipop. **[2]**

(c) Use your answers to parts **(a)** and **(b)** to determine the expected number of bottles which contain less than 330 ml in a random sample of 100 bottles. **[3]**

In order to comply with new regulations, no more than 1% of bottles of Fizzipop should contain less than 330 ml.

The manufacturer decides to meet the new regulations by adjusting the manufacturing process so that the mean volume of drink in a bottle of Fizzipop is increased.

The standard deviation is unaltered.

(d) Determine the minimum mean volume of drink in a bottle of Fizzipop which should ensure that the new regulations are met. Give your answer to **3** significant figures. **[3]**

The mean volume of drink in a bottle of Fizzipop is set to 340 ml. After several weeks the quality control manager suspects the mean volume may have reduced. She collects a random sample of 100 bottles of Fizzipop.

The mean volume of drink in a bottle in the sample is found to be 339.37 ml.

(e) Assuming the standard deviation is unaltered, conduct a hypothesis test at the 5% level to determine whether there is any evidence to suggest that the mean volume of drink in a bottle of Fizzipop is less than 340 ml. **[7]**

15(a)(i)	
15(a)(ii)	

15(b)	
15(c)	
15(d)	

15(e)