

Question	Answer	Mark	Mark scheme	Additional guidance
20	2, 3, 4	M1	for method to solve $3n + 2 \leq 14$ eg $n \leq (14 - 2) \div 3$ oe	This could be shown within an equation rather than an inequality at this stage
		M1	for complete method to rearrange $\frac{6n}{n^2 + 5} > 1$ to the form $an^2 + bn + c (< 0)$	For the 2rd and 3rd M marks condone no ' < 0 ' and condone use of incorrect inequality signs or '='
		M1	for method to begin to solve $n^2 - 6n + 5 (< 0)$ eg $(n \pm 5)(n \pm 1) (< 0)$	Accept $\frac{- -6 \pm \sqrt{(-6)^2 - 4 \times 1 \times 5}}{2 \times 1}$ (condone one sign error)
		M1	(dep on previous M2) for $n > 1$ and $n \leq 4$ or $1 < n < 5$	Must come from correct working Could be shown on a number line
		A1	(dep M4) cao	
			Alternative method	
		M1	for method to solve $3n + 2 \leq 14$ eg $n \leq (14 - 2) \div 3$ oe	This could be shown within an equation rather than an inequality at this stage
			OR for $3 \times 4 + 2 = 14$	
		M3	for trials with 1, 2, 3 and 4 in the quadratic inequality, correctly evaluated	The values from the trials may be given as improper fractions eg $\frac{24}{21}, \frac{18}{14}, \frac{12}{9}, \frac{6}{6}$
		(M2	for trials with three of 1, 2, 3 and 4, correctly evaluated)	
		(M1	for trials with two of 1, 2, 3 and 4, correctly evaluated)	
		A1	(dep M4) cao	