Advanced genetics - 203.305 Microarray - Hands-on data analysis

Dr. Pierre-Yves Dupont, Postdoctoral Researcher

Computer Biology Research Group

p.y.dupont@massey.ac.nz

Planning

- 15.9.14 Introduction (lecture), AHB2.38
- 22.9.14 Paper discussion, AHB2.38 Discussion worksheet
 due
- 23.9.14 From raw data to lists of differentially expressed genes (Step by step analysis of a microarray data set using the R language, **3h lab**, *SC5.10*)
- 29.09.14 Lab discussion (feedback!) and new developments in global gene expression analysis, *AHB2.38*
- 30.09.14 Biological interpretation of microarray data (Gene ontology analysis using the R language + online research of candidate genes, **3h lab**, *SC5.10*)

Microarray studies

- 1. Indroduction
- 2. Microarray technology
- 3. Statistics
- 4. Gene expression databases and MIAME
- 5. Examples of microarray studies (paper discussion topic and lab topic)

Microarray applications

- Gene expression analysis
- Re-sequencing
- SNP-analysis
- DNA-Protein interactions
- Discovery of new transcripts/alternative splice variants

Expression Studies




Figure modified from: Katherine Joyce, Woods Hole Oceanographic Institution

Definitions

- Genome: entire DNA sequence of an organism
- Epigenome: chemical marks of the genome that modify its expression
- Transcriptome: all gene transcripts present in a given cell/tissue at a given time ("snapshot")
- Transcriptomics: global analysis of gene expression = genome-wide expression profiling

Definitions

- cDNA: complementary DNA made from mRNA by the enzyme reverse transcriptase
- EST: Expressed Sequence Tag, small pieces of an expressed gene (cDNA)
- Hybridization: based on complementary molecules, sequences that are able to base-pair with one another. When two complementary sequences find each other, they will lock together, or hybridize (primer annealing, probe-target binding etc).

Genome-wide expression studies - Medical applications

- Cancer research: Cell-cycle monitoring, genetic markers detection
- **Drug development and response**: Treatment-induced expression pattern
- Diagnosis: Disease-associated expression patterns

Genome-wide expression studies -Biological applications

- Development biology: comparison of different developmental stages
- **Ecology**: interactions between organisms (symbiosis, pathogenicity...) or between organisms and environment (temperature, nutrient...)
- Evolution: within and between species variation, hybrids vs. parents, diploids vs. polyploids
- Functional analyses: wild type vs. mutant

Hypothesis generating tool


Microarray studies

- 1. Indroduction
- 2. Microarray technology
- 3. Statistics
- 4. Gene expression databases and MIAME
- 5. Examples of microarray studies (paper discussion topic and lab topic)


What are microarrays?

A microarray is a **solid support** (such as a membrane or glass microscope slide) on which **DNA of known sequence** is deposited in a **grid-like array**.






Microarray analysis principle



Imge from: http://www.scq.ubc.ca/image-bank/


Microarray analysis principle





Imge from: http://www.scq.ubc.ca/image-bank/

It is possible to represent different samples on one microarray using different fluorescent molecules (fluorophores)

- Cyanin 3 (Cy3): green fluorescence (excited at 550nm, emission at 570nm)
- Cyanin 5 (Cy5): red fluorescence (excited at 650nm, emission at 770nm)







Question driven

Goals? Hypothesis? Questions?

Platform

- What technology?
- Source of the gene probes?
- Cross-species hybridization?

- Platform
- Experimental design

- What statistics?
- What analysis software?
- Replication level
- Hybridization scheme

- Platform
- Experimental design
- Laboratory steps

- Sample preparation and labelling
- Hybridization
- Washing
- Image acquisition

- Platform
- Experimental design
- Laboratory steps
- Bioinformatics steps

- Data transformation and normalization
- Analysis of differentially expressed genes (statistical tests, gene ontology, ...)
- Visualization (graphics)
- Data storage (databases, MIAME standards)


- Platform
- Experimental design
- Laboratory steps
- Bioinformatics steps
- Data interpretation

- Answers?
- New hypotheses?
- Follow-up experiments?
- Validation?

Microarray studies

- 1. Indroduction
- 2. Microarray technology
- 3. Statistics
- 4. Gene expression databases and MIAME
- 5. Examples of microarray studies (paper discussion topic and lab topic)

Log Fold Ratio



Expression ratio: $log(rac{Cy5}{Cy3})$

Log Fold Ratio

- Cy3 = Sample1 (Green)
- Cy5 = Sample2 (Red)
- Cy5 > Cy3: higher expression in sample 2
- Cy3 > Cy5: higher expression in sample 1
- M(log fold ratio) = $log_2(\frac{Cy5}{Cy3})$
- ullet M(log fold ratio) = $log_2(extstyle Cy5) log_2(extstyle Cy3)$

Log Fold Ratio

Reminder: $log_2(x)$ is the unique real number y such that: $b^y=x$. For example: $log_2(8)=3$ because $2^3=8$

$log_2({\color{red}Cy5}/{\color{red}Cy3})$	Cy5/Cy3
2	4
1.58	3
1	2
0.58	1.5
0	1
-0.58	0.66
-1	0.5
-1.58	0.33
-2	0.25

Hypothesis testing

T-test

Null hypothesis (H_0): gene A is **not** differentially expressed between two treatments

Mean:

$$\overline{x} = rac{1}{M} \sum_{i=1}^{M} x_i$$
; for gene x in M $replicates$


Variance:

$$S_{x}^{2}=rac{1}{M-1}\sum_{i=1}^{M}(x_{i}-\overline{x}^{2})$$

T-statistic:

$$T_x = rac{\overline{x_{S_1}} - \overline{x_{S_2}}}{\sqrt[2]{rac{S_{x_{S_1}}^2}{M} + rac{S_{x_{S_2}}^2}{N}}}$$

T-test and P-value



T-test is used only to compare two samples.
To compare more, ANOVA (<u>AN</u>alysis <u>O</u>f

<u>VA</u>riance) is used.

Hypothesis testing

T-test

Null hypothesis (H_0): gene A is **not** differentially expressed between two treatments

- 1. Compute the signal to noise ratio (difference of the means or medians) for each gene
- 2. Compute the t-statistic for each gene using the replicates
- 3. Compare t-statistic with the t-distribution
- 4. If t-statistic is more extreme than the critical t-statistic at a chosen significance level (e.g. lpha=0.05) reject the null hypothesis, otherwise accept it. **P-value estimation**

Quiz

- Usually, a p < **0.05** is considered small enough to reject the null hypothesis of no biological effect in favour of the alternative hypothesis of a biological effect.
- P-values are also known under type **1** error the probability of rejecting the null hypothesis when it is actually true (= false positive rate).
- P-value of 0.01 means a false positive rate of 1 %.
- When analyzing multidimensional data sets, p-values need to be adjusted for **multiple testing** .
- Two common p-value adjustment methods are **Bonferroni** and **False Discovery Rate**.

Bonferroni Correction

- If you hypothesize that a specific gene is up-regulated, p < 0.05 is fine.
- If you hypothesize that any of 10,000 genes is up-regulated, with p < 0.05 you can expect to see 5% (500 genes) up-regulated by chance alone.
- To account for the thousands of repeated measurements, some researchers apply a Bonferroni correction.

$$p < (0.05)/10,000 \ {
m or} \ p < 5e^{-6}$$

The Bonferroni correction is generally considered to be **too** conservative and **False Discovery Rate (FDR)** should be used.

False Discovery Rate Benjamini-Hochberg method

Imagine an array with 6400 genes and an experiment where 184 genes are differentially expressed at p=0.01: 64 genes would be expected to appear differentially expressed by chance alone.

FDR = false discovery rate = $rac{64}{184}*100=35\%$

False Discovery Rate

Benjamini-Hochberg method


	Observed Number of genes	Expected number of False Positives	FDR
10^{-2}	184	64	35
10^{-3}	35	6	18
10^{-4}	15	0.6	4

With decreasing p-value, FDR also decreases, but so does the number of differentially expressed genes – choose a p-value which balances both!


Microarray studies

- 1. Indroduction
- 2. Microarray technology
- 3. Statistics
- 4. Gene expression databases and MIAME
- 5. Examples of microarray studies (paper discussion topic and lab topic)


Gene Expression Omnibus (GEO) @ NCBI (http://www.ncbi.nlm.nih.gov/geo/)




Geo Datasets @ NCBI (http://www.ncbi.nlm.nih.gov/gds/)
Geo Profiles @ NCBI (http://www.ncbi.nlm.nih.gov/geoprofiles/)



ArrayExpress @ EBI (http://www.ebi.ac.uk/arrayexpress/)



Expression Atlas @ EBI (http://www.ebi.ac.uk/gxa/)



MIAME Standard

Minimum Information About a Microarray
Experiment that is needed to enable the
interpretation of the results of the
experiment unambiguously and potentially
to reproduce the experiment

http://fged.org/Workgroups/MIAME/miame.html/

MIAME Standard

- 1. Raw data for each hybridisation (CEL or GPR files)
- 2. **Processed** (normalised) **data** (used to draw the conclusions from the study)
- 3. Essential **sample annotation** including experimental factors and their values
- 4. **Experimental design** including sample data relationships (e.g. which hybridizations are technical and biological replicates)
- 5. Sufficient **array annotation** (e.g. gene identifiers, probe sequences)
- 6. Essential **laboratory and data processing protocols** (e.g. normalization method used to obtain the final data)

Microarray studies

- 1. Indroduction
- 2. Microarray technology
- 3. Statistics
- 4. Gene expression databases and MIAME
- 5. Examples of microarray studies (paper discussion topic and lab topic)

Microarray paper discussion

MOLECULAR ECOLOGY

Molecular Ecology (2009) 18, 3227-3239

doi: 10.1111/j.1365-294X.2009.04261.x

Adaptive differences in gene expression associated with heavy metal tolerance in the soil arthropod *Orchesella* cincta

DICK ROELOFS,* THIERRY K. S. JANSSENS,* MARTIJN J. T. N. TIMMERMANS,*
BENJAMIN NOTA,* JANINE MARIËN,* ZOLTÁN BOCHDANOVITS,† BAUKE YLSTRA‡ and
NICO M. VAN STRAALEN*

*Institute of Ecological Science, VU University Amsterdam, De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands, †Department of Clinical Genetics, Section Medical Genomics, VU Medical Center, Van der Boechorststraat 7, 1081 BT Amsterdam, The Netherlands, ‡Microarray Facility CCA, VU Medical Center, De Boelelaan 1117, 1081 HV Amsterdam, The Netherlands

Lab case study

Herbivory in Nicotiana attenuata (Solanaceae)