


This article was downloaded by: [Stockholm University Library]

On: 02 February 2015, At: 14:19

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Communications in Algebra

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lagb20>

Cleft extensions for a hopf algebra generated by a nearly primitive element

Akira Masoka^a

^a Department of Mathematics , Shimane University , Shimane, 690, Japan
Published online: 27 Jun 2007.

To cite this article: Akira Masoka (1994) Cleft extensions for a hopf algebra generated by a nearly primitive element , Communications in Algebra, 22:11, 4537-4559, DOI: [10.1080/00927879408825086](https://doi.org/10.1080/00927879408825086)

To link to this article: <http://dx.doi.org/10.1080/00927879408825086>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

CLEFT EXTENSIONS FOR A HOPF ALGEBRA
GENERATED BY A NEARLY PRIMITIVE ELEMENT

Akira MASUOKA

Department of Mathematics

Shimane University

Matsue, Shimane 690 JAPAN

The concept of cleft extensions, or equivalently of crossed products, for a Hopf algebra is a generalization of Galois extensions with normal basis and of crossed products for a group. The study of these subjects was founded independently by Blattner-Cohen-Montgomery [BCM] and by Doi-Takeuchi [DT1]. Afterward, Doi [D] improved the theory and determined, in particular, the isomorphic classes of cleft extensions for a certain commutative and cocommutative Hopf algebra. In this paper, we determine such classes of those for a non-commutative, non-cocommutative Hopf algebra A_N of free rank N^2 , which is generated by a group-like g

and a $(1, g)$ -primitive x . For example, the cyclic algebra will be seen to be a cleft extension for A_N over the ground ring.

In the course we do not depend on the general results in [D, § 2], but take a direct method using Bergman's results [B].

Throughout we work over a commutative ring R . Algebra, Hopf algebra, linear and \otimes mean R -algebra, Hopf algebra over R , R -linear and \otimes_R , respectively.

§ 1 Preliminaries

In this section, for later use we recall some fundamental definitions and results on cleft extensions. The main reference is [D, § 1].

Let A be a Hopf algebra with coalgebra structure Δ, ε . Fix an algebra C .

A right A -comodule algebra B (with A -comodule structure $\rho : B \rightarrow B \otimes A$) is called an A -cleft extension over C [D, p.3056], if B contains C as coinvariant subalgebra, that is, $C = \{ b \in B \mid \rho(b) = b \otimes 1 \}$, and if there is such an A -comodule map $\phi : A \rightarrow B$ that is invertible under the convolution product $*$ [S, p.69]. In this case, ϕ can be chosen so as to be unitary ($\phi(1) = 1$) [DT1, p.813]. A unitary invertible A -comodule map $A \rightarrow B$ is called a section [D, p.3056].

We call a pair (B, ϕ) of an A -cleft extension B/C and a section ϕ a cleft system for A over C .

1.1 LEMMA. Let (B, ϕ) be a cleft system. A linear map $\phi': A \rightarrow B$ is a section, if and only if there is a unitary invertible linear map $\gamma: A \rightarrow C$ such that $\phi' = \gamma * \phi$.

Proof. See the proof of [D, Lemma 2.1]. ■

In general, an algebra B given an algebra map $C \rightarrow B$ is called a C-ring [B, p.195]. An A -cleft extension over C is a C-ring in an obvious way. An isomorphism $B \rightarrow B'$ between A -cleft extensions over C means an isomorphism of A -comodule C-rings (precisely, of A -comodules and of C-rings).

1.2 DEFINITION. Denote by

$$\text{Cleft}(A, C)$$

the set of isomorphic classes of A -cleft extensions over C .

An A -cleft extension B/C is characterized as an A -Galois extension with normal basis [DT1, Thm.9], by which we mean that B is a right A -comodule algebra with coinvariant subalgebra C such that $b' \otimes b \mapsto b' \rho(b)$, $B \otimes_C B \rightarrow B \otimes A$ is an isomorphism and that there is a left C -module right A -comodule isomorphism $C \otimes A \simeq B$. This isomorphism is given by $c \otimes a \mapsto c\phi(a)$, if ϕ is a section.

1.3 LEMMA. Each A -comodule C -ring map $F : B \rightarrow B'$ between A -cleft extensions over C is an isomorphism.

Proof. Let $\phi : A \rightarrow B$ be a section. Then the composition $\phi' = F\phi : A \rightarrow B'$ is a section, too. The lemma follows from the commutative diagram:

$$\begin{array}{ccc} & & B \\ & \simeq \nearrow & \\ C \otimes A & & \downarrow F \\ & \simeq \searrow & \\ & & B' \end{array}$$

Here the isomorphisms are induced, as above, from ϕ , ϕ' , respectively. ■

A pair (\triangleright, σ) is called a crossed system for A over C [D, p.3055], if

$$\triangleright : A \otimes C \rightarrow C$$

is a measuring action, if

$$\sigma : A \otimes A \rightarrow C$$

is an invertible linear map, and if these satisfy the normal condition

$$1 \triangleright c = c \quad (c \in C)$$

$$\sigma(a, 1) = \varepsilon(a)1 = \sigma(1, a) \quad (a \in A)$$

as well as the twisted module condition, the cocycle condition described in [D, (1), (2)].

From a cleft system (B, ϕ) , a crossed system (\triangleright, σ) is obtained as follows:

$$a \triangleright c = \sum \phi(a_1) c \phi^{-1}(a_2)$$

$$\sigma(a, a') = \sum \phi(a_1) \phi(a'_1) \phi^{-1}(a_2 a'_2)$$

where $\Delta(a) = \sum a_1 \otimes a_2$, as usual.

Conversely, let (\triangleright, σ) be a crossed system and define a bilinear product on the R -module $C \star A = C \otimes A$ by

$$(c \star a)(c' \star a') = \sum c(a_1 \triangleright c') \sigma(a_2, a'_1) \star a_3 a'_2$$

for $c, c' \in C, a, a' \in A$. Then $C \star A$ with A -coaction $c \star a \mapsto \sum (c \star a_1) \otimes a_2$ is a right A -comodule algebra, called a crossed product. Furthermore, $C \star A$ and $a \mapsto 1 \star a, A \rightarrow C \star A$ form a cleft system for A over $C = C \star R$. See [BM, Thm.1.18], [D, Prop.1.1].

1.4 PROPOSITION. These give a 1-1 correspondence between the isomorphic classes of cleft systems and the crossed systems (both for A over C). (An isomorphism of cleft systems means an A -comodule C -ring isomorphism consistent with the sections.)

Proof. This is essentially shown in [D, Prop.1.1; Thm.1.2]. ■

An A -cleft extension B/C is said to be twisted (respectively, smashed), if $B \simeq C \star A$, a crossed product such that \triangleright (respectively, σ) is trivial. See [D, p.3056; p.3059].

§ 2 Cleft Extensions for A_N

We fix an integer N such that $N \geq 2$ and suppose R contains a root ζ of the N -th cyclotomic polynomial over \mathbb{Z} [L, VIII, § 3]. Hence $\zeta^N = 1$ and there

is a ring map $\mathbb{Z}[\zeta_N] \rightarrow R$, sending a primitive N -th root $\zeta_N \in \mathbb{C}$ of 1 to ζ . Note that ζ may not be a primitive N -th root of 1. For example ζ may be 1, if N is a power of a prime p and $\text{ch } R = p$.

2.1 DEFINITION. Let i, n be integers such that $0 \leq i \leq n$. We denote by

$$\binom{n}{i}$$

the element in R got by putting $q = \zeta$ into the Gauss polynomial

$$\binom{n}{i}_q = \frac{(q^n - 1) \dots (q^{n-i+1} - 1)}{(q^i - 1) \dots (q - 1)}.$$

2.2 LEMMA. 1) If $0 < i < N$, then $\binom{N}{i} = 0$.

2) If $0 < i < n$, then $\binom{n}{i} = \binom{n-1}{i-1} + \zeta^i \binom{n-1}{i}$.

Proof. 1) This follows, since $\binom{N}{i}_q = 0$ ($0 < i < N$) if $q = \zeta_N$. 2) This is well-known. ■

2.3 DEFINITION. Denote by

$$A_N \quad \text{or} \quad A_{N, \zeta}$$

the Hopf algebra defined as follows. As an algebra, this is generated by g, x with the relations:

$$r1) g^N = 1; \quad r2) x^N = 0; \quad r3) xg = \zeta gx.$$

The coalgebra structure is determined by:

$$\Delta(g) = g \otimes g, \quad \varepsilon(g) = 1$$

$$\Delta(x) = x \otimes g + 1 \otimes x, \quad \varepsilon(x) = 0$$

(Recall that such an x is called nearly primitive,

or more precisely $(1, g)$ -primitive.) In fact A_N is a Hopf algebra whose antipode is the algebra anti-endomorphism determined by:

$$S(g) = g^{N-1}, \quad S(x) = -\zeta^{-1}g^{N-1}x$$

(We remark that $S^{2N} = 1$.)

To understand A_N adequately, one sees there are Hopf algebra quotients

$$A_{\text{fr}} \rightarrow A_{\infty} \rightarrow A_N.$$

Here

$$A_{\text{fr}} = R\{g, g^{-1}, x\}/(gg^{-1} - 1, g^{-1}g - 1),$$

the (free) Hopf algebra defined in [S, Example-Exercise, p.89],

$$A_{\infty} = A_{\text{fr}}/(xg - \zeta gx)$$

(which is defined for any $\zeta \in R$), and finally

$$A_N = A_{\infty}/(g^N - 1, x^N).$$

Note that in a Hopf algebra an ideal generated by nearly primitives is a Hopf ideal.

2.4 REMARK. 1) A_N is a particular one of Taft's Hopf algebra [T], which is in fact generated by several group-likes g, g_2, \dots, g_n and x as above. A_2 was previously defined by Sweedler.

2) Let q be a unit in R such that $\zeta = q^{-2}$. Then A_{∞} is regarded as a quantized enveloping algebra of the Lie algebra of upper triangular matrices in $\mathfrak{sl}(2)$ and, if $q - q^{-1}$ is a unit, this is a Hopf subalgebra of $U_q(\mathfrak{sl}(2))$.

2.5 LEMMA. 1) A_N is a free R -module with a basis $g^m x^n$, $0 \leq m, n < N$, of N^2 elements.

$$2) \Delta(g^m x^n) = \sum_{i=0}^n \binom{n}{i} g^m x^{n-i} \otimes g^{m+n-i} x^i \quad (0 \leq m, n < N)$$

Proof. 1) This is shown by applying the Bergman Theorem [B, Thm.1.2].

2) If $wz = zw$ in an algebra, then

$$(z + w)^n = \sum_{i=0}^n \binom{n}{i} z^{n-i} w^i.$$

Apply this to the sum $\Delta(x) = x \otimes g + 1 \otimes x$. ■

Part 1) yields that A_N is not cocommutative as a coalgebra and that, if $\zeta \neq 1$, A_N is not commutative as an algebra.

Let E be an algebra. Denote by E^\times the group of units in E .

2.6 LEMMA. A linear map $\gamma : A_N \rightarrow E$ is invertible, if and only if $\gamma(g^m) \in E^\times$ for all $m = 0, 1, \dots, N-1$.

Proof. "Only if". $\gamma(g^m)^{-1}$ is given by $\gamma^{-1}(g^m)$.

"If". By (2.5.2), $\gamma^{-1}(g^m x^n)$ is given inductively on n by

$$-\gamma(g^m)^{-1} \sum_{i=0}^{n-1} \binom{n}{i} \gamma(g^m x^{n-i}) \gamma^{-1}(g^{m+n-i} x^i)$$

or equally by

$$-\sum_{i=1}^n \binom{n}{i} \gamma^{-1}(g^m x^{n-i}) \gamma(g^{m+n-i} x^i) \gamma(g^{m+n})^{-1}. \quad \blacksquare$$

In the following, we fix an algebra C to determine the set $\text{Cleft}(A_N, C)$.

First, let (B, ϕ) be a cleft system for A_N over C and (\triangleright, σ) the corresponding crossed system.

Write:

$$(2.7) \quad G = \phi(g), \quad X = \phi(x)$$

Set:

$$(2.8) \quad \left\{ \begin{array}{l} \alpha(c) = g \triangleright c = GcG^{-1} \\ \delta(c) = x \triangleright c = [X, c]G^{-1} \quad (c \in C) \\ u = G^N \\ a = X^N \\ b = (\sigma(x, g) - \zeta \sigma(g, x)) \sigma(g, g)^{-1} \end{array} \right.$$

Here we use the usual bracket product: $[X, c] = Xc - cX$.

2.9 LEMMA. 1) $\alpha : C \rightarrow C$ is an algebra automorphism.

2) $\delta : C \rightarrow C$ is a $(1, \alpha)$ -derivation, that is, a linear endomorphism such that

$$\delta(cc') = \delta(c)\alpha(c') + c\delta(c') \quad (c, c' \in C).$$

3) $u \in C^x$, $a, b \in C$.

4) The following are fulfilled:

$$R1) \quad G^N = u$$

$$R2) \quad X^N = a$$

$$R3) \quad XG = \zeta GX + bG^2$$

$$R4) \quad Gc = \alpha(c)G \quad (c \in C)$$

$$R5) \quad Xc = cX + \delta(c)G$$

Proof. 1), 2). Easy.

3) By definition, $b \in C$. To show $u, a \in C$, we see through the A_N -coaction

$$u = G^N \mapsto G^N \otimes g^N = u \otimes 1,$$

$$a = x^N \mapsto \sum_{i=0}^N \binom{N}{i} x^{N-i} \otimes g^{N-i} x^i = a \otimes 1 \quad (\text{by (2.2.1)}).$$

4) R1), R2), R4) and R5) follow by the Definitions (2.8). R3) follows by some simple calculation. ■

In turn, take α, δ, b arbitrarily, where $\alpha, \delta \in \text{End } C$, the algebra of linear endomorphisms, and $b \in C$.

2.10 DEFINITION. Let $0 \leq n \leq N$.

1) Define $k_n = k_n(\alpha, \delta, b) \in C$ inductively by:

$$k_0 = 1$$

$$k_n = \delta(k_{n-1}) + k_{n-1}(b + \zeta \alpha(b) + \dots + \zeta^{n-1} \alpha^{n-1}(b))$$

$$(n \geq 1)$$

2) Define $h_n(c) = h_n(\alpha, \delta, b, c) \in \text{End } C$ by:

$$h_0(c) = c, \quad h_1(c) = \delta(c)$$

$$h_n(c) = \delta h_{n-1}(c) + h_{n-1}(c)(b + \zeta \alpha(b) + \dots + \zeta^{n-2} \alpha^{n-2}(b))$$

$$(n \geq 2)$$

Later in (2.15), we will apply Bergman's result [B, Prop.7.1], a generalization of his theorem quoted in the proof of (2.5.1). For this purpose, in the next lemma we regard R3)-R5) as a reduction system for the free algebra generated by elements G, X and the R -module C , that is, the tensor algebra $T(M)$ on $M = RG \oplus RX \oplus C$. Precisely, we regard R5), for example, as the linear endomorphisms r_{P-Q} of $T(M)$ defined as follows, where P, Q are iterated tensor products of RG, RX, C . r_{P-Q} is defined on the summand $P \otimes RX \otimes C \otimes Q$ by $r_{P-Q}(pXcq) = p(cX + \delta(c)G)q$ ($c \in C$,

$p \in P, q \in Q$), while r_{p-Q} is identical on the summands other than $P \otimes RX \otimes C \otimes Q$. See [B, (22), (23)].

2.11 LEMMA. Suppose the reduction system R3)-R5) for $T(M)$ is given. Let $1 \leq n \leq N$.

1) XG^n is reduced to

$$\zeta^n G^n X + (b + \zeta \alpha(b) + \dots + \zeta^{n-1} \alpha^{n-1}(b)) G^{n+1}.$$

2) $X^{n-1}(\zeta GX + bG^2)$ is reduced to

$$\sum_{i=0}^n \zeta^{n-i} \binom{n}{i} k_i G^{i+1} X^{n-i}.$$

In particular, $X^{N-1}(\zeta GX + bG^2)$ to $G^N X + k_N G^{N+1}$.

3) $X^{n-1}(cX + \delta(c)G)$ ($c \in C$) is reduced to

$$\sum_{i=0}^n \binom{n}{i} h_i(c) G^i X^{n-i}.$$

In particular, $X^{N-1}(cX + \delta(c)G)$ to $cX^N + h_N(c)G^N$.

Proof. Induction on n .

1) Case $n = 1$. OK by R3).

Case $n > 1$. Suppose the result for $n-1$ holds.

Then we have the following reductions:

$$\begin{aligned} XG^n &\mapsto \zeta^{n-1} G^{n-1} XG \\ &\quad + (b + \zeta \alpha(b) + \dots + \zeta^{n-2} \alpha^{n-2}(b)) G^n G \\ &\stackrel{R3)}{\mapsto} \zeta^{n-1} G^{n-1} (\zeta GX + bG^2) \\ &\quad + (b + \zeta \alpha(b) + \dots + \zeta^{n-2} \alpha^{n-2}(b)) G^{n+1} \\ &\stackrel{R4)}{\mapsto} \zeta^n G^n X + \zeta^{n-1} \alpha^{n-1}(b) G^{n+1} \\ &\quad + (b + \zeta \alpha(b) + \dots + \zeta^{n-2} \alpha^{n-2}(b)) G^{n+1} \\ &= \zeta^n G^n X + (b + \zeta \alpha(b) + \dots + \zeta^{n-1} \alpha^{n-1}(b)) G^{n+1} \end{aligned}$$

Thus the result for n holds.

2) Case $n = 1$. Trivial.

Case $n > 1$. Suppose the result for $n-1$ holds.

Then:

$$\begin{aligned}
 X^{n-1}(\zeta GX + bG^2) &\mapsto \sum_0^{n-1} \zeta^{n-1-i} \binom{n-1}{i} X k_i G^{i+1} X^{n-1-i} \\
 &\mapsto \sum_0^{n-1} \zeta^{n-1-i} \binom{n-1}{i} (k_i X + \delta(k_i) G) G^{i+1} X^{n-1-i} \\
 \text{R5)} \quad &\mapsto \sum_0^{n-1} \zeta^{n-1-i} \binom{n-1}{i} k_i \\
 1) \quad &\times (\zeta^{i+1} G^{i+1} X + (b + \dots + \zeta^i \alpha^i(b)) G^{i+2}) X^{n-1-i} \\
 &+ \sum_0^{n-1} \zeta^{n-1-i} \binom{n-1}{i} \delta(k_i) G^{i+2} X^{n-1-i} \\
 &= \sum_0^{n-1} \zeta^{n-1-i} \binom{n-1}{i} k_{i+1} G^{i+2} X^{n-1-i} \\
 &+ \sum_0^{n-1} \zeta^n \binom{n-1}{i} k_i G^{i+1} X^{n-i} \quad (\text{by (2.10.1)}) \\
 &= \sum_0^n \zeta^{n-i} \binom{n}{i} k_i G^{i+1} X^{n-i} \quad (\text{by (2.2.2)})
 \end{aligned}$$

Thus the result for n holds.

3) Similar. ■

2.12 DEFINITION. A 5-tuple $\underline{d} = (\alpha, \delta, u, a, b)$ is called cleft data for A_N over C , if the following D0)-D7) are fulfilled:

D0) $\alpha : C \rightarrow C$ is an algebra automorphism, $\delta : C \rightarrow C$ a $(1, \alpha)$ -derivation, and $u \in C^\times$, $a, b \in C$.

D1) $\alpha(u) = u$

D2) $\delta(u) = (b + \zeta \alpha(b) + \dots + \zeta^{N-1} \alpha^{N-1}(b)) u$

D3) $\delta(a) = 0$

D4) $a - \alpha(a) = k_N(\alpha, \delta, b)u$

D5) $\alpha^N(c) = ucu^{-1}$

$$D6) \quad b\alpha^2(c) - \alpha(c)b = (\delta\alpha - \zeta\alpha\delta)(c) \quad (c \in C)$$

$$D7) \quad [a, c] = h_N(\alpha, \delta, b, c)u$$

We denote the set of all such data by

$$\mathcal{D} = \mathcal{D}(A_N, C).$$

2.13 EXAMPLE. 1) Take $\alpha = 1$, $\delta = 0$. (Then $k_N = 0$, $h_n = 0$ ($n > 0$).) $\underline{d} = (1, 0, u, a, b)$ is cleft data, if and only if $u \in Z(C)^*$, $a, b \in Z(C)$, the center of C .

2) Take $u = 1$, $a = b = 0$. (Then $k_N = 0$, $h_n(c) = \delta^n(c)$ ($n > 0$).) $\underline{d} = (\alpha, \delta, 1, 0, 0)$ is cleft data, if and only if α is an algebra automorphism, δ a $(1, \alpha)$ -derivation, $\alpha^N = 1$, $\delta^N = 0$, and $\delta\alpha = \zeta\alpha\delta$.

2.14 DEFINITION. Let $\underline{d} = (\alpha, \delta, u, a, b) \in \mathcal{D}$. Define a pair $(B_{\underline{d}}, \phi_{\underline{d}})$ as follows: $B_{\underline{d}}$ is the C -ring generated by G, X with the relations R1)-R5). $\phi_{\underline{d}}: A_N \rightarrow B_{\underline{d}}$ is the linear map defined by

$$\phi_{\underline{d}}(g^m x^n) = G^m X^n \quad (0 \leq m, n < N).$$

2.15 PROPOSITION. Let $\underline{d} \in \mathcal{D}$.

- 1) $B_{\underline{d}}$ is a free left C -module with a basis
- $$G^m X^n \quad (0 \leq m, n < N).$$

In particular, we have $C \subset B$.

2) $B_{\underline{d}}$ has a right A_N -comodule algebra structure determined by

$$G \mapsto G \otimes g, \quad X \mapsto X \otimes g + 1 \otimes x.$$

The coinvariant subalgebra equals C .

- 3) $\phi_{\underline{d}}$ is a section.

Thus $(B_{\underline{d}}, \phi_{\underline{d}})$ is a cleft system for A_N over C .

Proof. 1) We apply the left version of Bergman [B, Prop.7.1], in which we should replace the base ring \mathbb{Z} by R .

We introduce a total ordering \leq into the set of words consisting of G, X . Define $G < X$. For two words W, W' , set $W < W'$, if either $\text{Length } W < \text{Length } W'$ or if $\text{Length } W = \text{Length } W'$ and W precedes W' under lexicographic ordering. This \leq is consistent with R1)-R5) in the sense of [B, p.198, lines -10, -9] and satisfies the descending chain condition. Let $R')$ be the reduction system for the tensor algebra $T(M)$ ($M = RG \oplus RX \oplus C$) which consists of R1)-R5) and

R6) $c \cdot c' =$ the product of c and c' in C ($c, c' \in C$) (see [B, (21)]). To show 1), by the left version of [B, Prop.7.1] it suffices to prove that all ambiguities of $R')$ are resolvable. There are no inclusion ambiguities. We have the following overlap ambiguities:

$$\begin{aligned} (G^i, G^{N-i}, G^i), & \quad (X^i, X^{N-i}, X^i) \quad (0 < i < N) \\ (X, G, G^{N-1}), & \quad (X^{N-1}, X, G) \\ (G^{N-1}, G, c), & \quad (X^{N-1}, X, c), \quad (X, G, c) \\ & \quad (c, c' \in C) \\ (G, c, c'), & \quad (X, c, c') \end{aligned}$$

Here we omit the ways of reductions, since they seem obvious. For example, by saying (X, G, G^{N-1}) is resolvable we mean that $(\{GX + bG^2\}G^{N-1}$ and Xu can be reduced to the same expression through R1)-R5).

This is resolvable in fact by D2) and (2.11.1). So is (X^{N-1}, X, G) by D4), (2.11.2), (X^{N-1}, X, c) by D7), (2.11.3), (G^i, G^{N-i}, G^i) by D1), (X^i, X^{N-i}, X^i) by D3), (G^{N-1}, G, c) by D5), (X, G, c) by D6), and finally (G, c, c') and (X, c, c') by D0).

2), 3). One sees directly that the algebra map $B_{\underline{d}} \rightarrow B_{\underline{d}} \otimes A_N$ defined in 2) is a well-defined A_N -comodule structure and that $\phi_{\underline{d}}$ is an A_N -comodule map. By (2.6), $\phi_{\underline{d}}$ is invertible since G is invertible in $B_{\underline{d}}$. By 1), $c \otimes a \mapsto c \phi_{\underline{d}}(a)$, $C \otimes A_N \rightarrow B_{\underline{d}}$ is a unitary, C -module and A_N -comodule isomorphism. Hence the coinvariant subalgebra equals C . ■

2.16 EXAMPLE. Take $\alpha = 1$, $\delta = 0$, $u, a \in R^\times$, $b = 0$. Then one sees from (2.13.1), (2.15) that the cyclic algebra $\left(\frac{u, a}{R, \zeta}\right)$ [P, p.284] with a natural A_N -coaction is a (twisted) A_N -cleft extension over R .

2.17 PROPOSITION. Any A_N -cleft extension B over C is isomorphic with $B_{\underline{d}}$ for some $\underline{d} \in \mathcal{D}$.

Proof. Suppose (B, ϕ) is a cleft system. Set G, X as in (2.7), $\underline{d} = (\alpha, \delta, u, a, b)$ as in (2.8). By (2.9), one traces the proof of (2.15.1) in reverse to see that \underline{d} is cleft data. For example, for D2) one sees: $uX + \delta(u)G = Xu$ (by R5))

$$\begin{aligned} &= XG^N \quad (\text{by R1}) \\ &= G^N X + (b + \zeta \alpha(b) + \dots + \zeta^{N-1} \alpha^{N-1}(b)) G^{N+1} \\ &\quad (\text{by R3), R4), (2.11.1)}) \\ &= uX + (b + \zeta \alpha(b) + \dots + \zeta^{N-1} \alpha^{N-1}(b)) uG \quad (\text{by R1}) \end{aligned}$$

By (2.9) again, one has a natural A_N -comodule C -ring map $B_{\underline{d}} \rightarrow B$. This is an isomorphism by (1.3). ■

2.18 DEFINITION. Suppose $\alpha, \delta \in \text{End } C$, $b, t \in C$. Let $0 \leq n \leq N$. Define

$$p_n = p_n(\alpha, \delta, b, t) \in C$$

inductively by

$$p_0 = 1, \quad p_1 = t$$

$$p_n = (t\alpha + \delta)(p_{n-1}) + p_{n-1}(b + \zeta \alpha(b) + \dots + \zeta^{n-2} \alpha^{n-2}(b))$$

$$(n \geq 2)$$

2.19 LEMMA. Let $\underline{d}, \underline{d}', \underline{d}'' \in \mathcal{D}$.

1) Let $F : B_{\underline{d}'} \rightarrow B_{\underline{d}}$ be an isomorphism of A_N -comodule C -rings. Then there is a unique pair $(v, t) \in C^\times \times C$ such that

$$(2.20) \quad F(G') = vG, \quad F(X') = X + tG.$$

2) Let $(v, t) \in C^\times \times C$. Then the C -ring map $F : B_{\underline{d}'} \rightarrow B_{\underline{d}}$ determined by (2.20) is well-defined, if and only if

$$(2.21) \quad \left\{ \begin{array}{l} \alpha'(c) = v\alpha(c)v^{-1} \\ \delta'(c) = \{(t\alpha + \delta)(c) - ct\}v^{-1} \quad (c \in C) \\ u' = v\alpha(v) \dots \alpha^{N-1}(v)u \\ a' = a + p_N(\alpha, \delta, b, t)u \\ b' = \{vb + (t\alpha + \delta)(v) - \zeta v\alpha(t)\}(v\alpha(v))^{-1} \end{array} \right.$$

where $\underline{d} = (\alpha, \dots)$, $\underline{d}' = (\alpha', \dots)$. In this case, F is an A_N -comodule isomorphism.

3) Let $F' : B_{\underline{d}''} \rightarrow B_{\underline{d}'}$, $F : B_{\underline{d}'} \rightarrow B_{\underline{d}}$ be isomorphisms of A_N -comodule C -rings determined as in 1) by

$(w, s), (v, t) \in C^* \times C$, respectively. Then the composition FF' is determined by $(wv, sv + t)$.

Proof. 1) $F_{\underline{d}'}$, $\phi_{\underline{d}}$ are sections $A_N \rightarrow B_{\underline{d}}$. Hence by (1.1) there is a unitary, invertible linear map $\gamma : A_N \rightarrow C$ such that $F_{\underline{d}'} = \gamma * \phi_{\underline{d}}$. Set $v = \gamma(g) \in C^*$, $t = \gamma(x) \in C$. Then one sees easily that (2.20) holds. The uniqueness is obvious.

2) This is verified directly. For example, F is consistent with R2), if and only if $a' = (X + tG)^N$. Since one sees inductively that

$$(X + tG)^n = \sum_{i=0}^n \binom{n}{i} p_i G^i X^{n-i} \quad (0 \leq n \leq N),$$

the 4th condition in (2.21) follows.

3) Easy. ■

The group C^* acts on the additive group C by the right multiplication. So we have the group $C^* \times C$ of semi-direct product with the multiplication

$$(w \times s)(v \times t) = wv \times (sv + t).$$

2.22 PROPOSITION. 1) $C^* \times C$ acts on the set \mathcal{D} from the left with the action

$$\underline{d}' = (v \times t)\underline{d}$$

defined by (2.21).

2) Suppose $d, d' \in \mathcal{D}$. Then $B_{\underline{d}} \cong B_{\underline{d}'}$, if and only if \underline{d} and \underline{d}' are $C^* \times C$ -equivalent.

Proof. 1) If $\underline{d} \in \mathcal{D}$, then \underline{d}' defined by (2.21) is contained in \mathcal{D} . In fact, by the proof of

(2.19.1) \underline{d}' is cleft data coming from a cleft system $(B_{\underline{d}}, \phi)$, where $\phi(g) = vG$, $\phi(x) = X + tG$.

Suppose $\underline{d}'' = (w \times s)\underline{d}'$ in addition. Then by (2.19.2) we have the isomorphisms

$$F' : B_{\underline{d}''} \rightarrow B_{\underline{d}'}, \quad F : B_{\underline{d}'} \rightarrow B_{\underline{d}}.$$

By (2.19.3) the composition FF' is determined by $(w \times s)(v \times t)$. Hence by (2.19.2)

$$\underline{d}'' = ((w \times s)(v \times t))\underline{d},$$

so $C^x \times C$ acts as a group.

2) This follows by (2.19.1), (2.19.2). ■

2.23 THEOREM. $\underline{d} \mapsto B_{\underline{d}}$ gives a 1-1 correspondence between the set $C^x \times C \setminus \mathcal{D}(A_N, C)$ of $C^x \times C$ -orbits in $\mathcal{D}(A_N, C)$ and the set $\text{Cleft}(A_N, C)$ of isomorphic classes of A_N -cleft extensions over C .

Proof. This follows by (2.17), (2.22.2). ■

2.24 PROPOSITION. Let $\underline{d} = (\alpha, \delta, u, a, b) \in \mathcal{D}$.

1) $B_{\underline{d}}$ is twisted, if and only if there exist $v \in C^x$, $t \in C$ such that

$$\begin{aligned} \alpha(c) &= vcv^{-1} \quad (\alpha \text{ is inner}) \\ \delta(c) &= [t, c]v^{-1} \end{aligned}$$

for $c \in C$. (Such δ may be called an inner $(1, \alpha)$ -derivation.)

2) $B_{\underline{d}}$ is smashed, if and only if there exist $v \in C^x$, $t \in C$ such that

$$\begin{aligned} u &= \alpha^{N-1}(v) \dots \alpha(v)v \\ a &= -p_N(\alpha, \delta, b, t)u \end{aligned}$$

$$b = \zeta \alpha(t) + (\delta(v) - vt) \alpha(v)^{-1}.$$

Proof. Let $\underline{d}' \in \mathcal{D}$ and (\triangleright, σ) be the crossed system corresponding to $(B_{\underline{d}'}, \phi_{\underline{d}'})$. Then it is easy to see:

$$\triangleright \text{ is trivial} \Leftrightarrow \alpha' = 1, \delta' = 0$$

$$\sigma \text{ is trivial} \Leftrightarrow u' = 1, a' = b' = 0$$

1) This follows by [D, Thm.2.3], but here we show directly.

By (2.19), the equivalent condition follows by setting $\alpha' = 1, \delta' = 0$ in the first two equations in (2.21) and replacing v, t by v^{-1}, tv^{-1} , respectively (or equally, by setting $\alpha = 1, \delta = 0$ there and deleting the primes).

2) Similarly, set $u' = 1, a' = b' = 0$ in the last three equations in (2.21) and replace v by v^{-1} . ■

It follows from (2.13.1) that any A_N -twisted extension B/C comes from such an extension E over $Z(C)$, that is, $B \simeq C \otimes_{Z(C)} E$. (Note $Z(C) \subset Z(E)$.)

§ 3 Cleft Extensions for A_2

Suppose $N = 2$ in § 2 (then $\zeta = -1$). We had better choose b in (2.8) rather as $b = \sigma(x, g) + \sigma(g, x)$. Under this change, we summarize the results. We also describe in (3.4) all crossed systems.

Define the Hopf algebra A_2 as in (2.3), setting $N = 2, \zeta = -1$. We need not assume anything on R .

A_2 is R -free with a basis $1, g, x, gx$.

3.1 DEFINITION. A 5-tuple $\underline{d} = (\alpha, \delta, u, a, b)$ is called cleft data for A_2 over a fixed algebra C , if the following are fulfilled:

D0) $\alpha : C \rightarrow C$ is an algebra automorphism, $\delta : C \rightarrow C$ a $(1, \alpha)$ -derivation, $u \in C^\times$, $a, b \in C$.

D1) $\alpha(u) = u$ D2) $\delta(u) = b - \alpha(b)$

D3) $\delta(a) = 0$ D4) $\delta(b) = a - \alpha(a)$

D5) $\alpha^2(c) = ucu^{-1}$

D6) $bc - \alpha(c)b = (\alpha\delta + \delta\alpha)(c)u$ ($c \in C$)

D7) $[a, c] = \delta(c)b + \delta^2(c)u$

We denote by $\mathcal{D} = \mathcal{D}(A_2, C)$ the set of all such data.

$C^\times \times C$ acts on \mathcal{D} by

$$\underline{d}' = (v \times t)\underline{d},$$

where

$$\alpha'(c) = v\alpha(c)v^{-1}$$

$$\delta'(c) = \{(t\alpha + \delta)(c) - ct\}v^{-1} \quad (c \in C)$$

$$u' = v\alpha(v)u$$

$$a' = a + tb + (t\alpha + \delta)(t)u$$

$$b' = vb + \{(t\alpha + \delta)(v) + v\alpha(t)\}u.$$

3.2 DEFINITION. Let $\underline{d} \in \mathcal{D}$. Denote by

$$B_{\underline{d}}$$

the A_2 -cleft extension over C defined as follows. As

a C -ring $B_{\underline{d}}$ is generated by G, X with the relations:

$$R1) G^2 = u \quad R2) X^2 = a \quad R3) XG = -GX + b$$

$$R4) \quad Gc = \alpha(c)G \quad R5) \quad Xc = cX + \delta(c)G \quad (c \in C)$$

The A_2 -comodule algebra structure of $B_{\underline{d}}$ is determined by $G \mapsto G \otimes g$, $X \mapsto X \otimes g + 1 \otimes x$.

In fact $B_{\underline{d}}$ is an A_2 -extension over C with a canonical section $\phi_{\underline{d}} : A_2 \rightarrow B_{\underline{d}}$ defined by $\phi_{\underline{d}}(g) = G$, $\phi_{\underline{d}}(x) = X$, $\phi_{\underline{d}}(gx) = GX$.

3.3 THEOREM. $\underline{d} \mapsto B_{\underline{d}}$ gives a 1-1 correspondence between $C^* \times C \setminus \mathcal{D}(A_2, C)$ and $\text{Cleft}(A_2, C)$.

3.4 PROPOSITION. There is a 1-1 correspondence between the set $\mathcal{D} \times C$ and the set of crossed systems for A_2 over C . Here $(\alpha, \delta, u, a, b, s) \in \mathcal{D} \times C$ corresponds to the crossed system (\triangleright, σ) defined as follows:

$$\begin{aligned} g \triangleright c &= \alpha(c), & x \triangleright c &= \delta(c) \\ gx \triangleright c &= \alpha(\delta(c)u + sc - \alpha(c)s) \end{aligned} \quad (c \in C)$$

TABLE OF $\sigma(b, \#)$

$\begin{smallmatrix} b \\ \# \end{smallmatrix}$	g	x	gx
g	u	s	$\alpha(s)$
x	$b + s$	a	$-\alpha(a) + \delta(s)$
gx	$\alpha(b) + s$	$\alpha(a)$	$-\alpha(a)u + \alpha(\delta(s)u) - \alpha(b+s)s$

Proof. For $s \in C$, define a linear map $\gamma_s : A_2 \rightarrow C$ by $\gamma_s(1) = \gamma_s(g) = 1$, $\gamma_s(x) = 0$, $\gamma_s(gx) = s$. Then one sees from the proof of (2.17) that $(\underline{d}, s) \mapsto (B_{\underline{d}}, \gamma_s * \phi_{\underline{d}})$ gives a 1-1 correspondence between $\mathcal{D} \times C$

and the isomorphic classes of cleft systems. Hence the Proposition follows by (1.4) and some direct calculation. ■

ACKNOWLEDGMENTS

I would like to thank Yukio Doi and Mitsuhiro Takeuchi. Professor Doi brought me the problem of determining $\text{Cleft}(A_2, C)$, which motivated me to do this research. Professor Takeuchi gave me valuable comments, reading an earlier manuscript. In particular, He improved Lemma 1.3, dropping the assumption that the antipode of A was bijective. Following his suggestion, I added Remark 2.4 2) and the explanation after Definition 2.3 when I revised the paper.

REFERENCES

- [B] G. Bergman, The diamond lemma for ring theory, Adv. in Math. 29(1978), 178-218.
- [BCM] R. Blattner, M. Cohen and S. Montgomery, Crossed products and inner actions of Hopf algebras, Trans. Amer. Math. Soc. 298(1986), 671-711.
- [BM] R. Blattner and S. Montgomery, Crossed products and Galois extensions of Hopf algebras, Pacific J. of Math. 137(1989), 37-54.
- [D] Y. Doi, Equivalent crossed products for a Hopf algebra, Comm. Algebra 17(1989), 3053-3085.
- [DT1] Y. Doi and M. Takeuchi, Cleft comodule algebras for a bialgebra, Comm. Algebra 14(1986), 801-817.

- [DT2] Y. Doi and M. Takeuchi, Hopf-Galois extensions of algebras, the Miyashita-Ulbrich actions, and Azumaya algebras, *J. Algebra* 121(1989), 488-516.
- [K] M. Koppinen, A duality theorem for crossed products of Hopf algebras, *J. Algebra* 146(1992), 153-174.
- [L] S. Lang, "Algebra", Addison-Wesley, Reading, Massachusetts, 1965.
- [P] R. Pierce, "Associative Algebras", GTM 88, Springer, New York-Heidelberg-Berlin, 1982.
- [S] M. Sweedler, "Hopf Algebras", Benjamin, New York, 1969.
- [T] E. Taft, The order of the antipode of finite-dimensional Hopf algebra, *Proc. Nat. Acad. Sci. USA* 68(1971), 2631-2633.

Received: July 1993

Revised: August 1993