


This article was downloaded by: [University of Toronto Libraries]

On: 06 January 2015, At: 23:16

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Communications in Algebra

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lagb20>

On the structure of relative hopf modules

Yukio Doi ^a

^a Fukui University, Fukui, 910, Japan

Published online: 27 Jun 2007.

To cite this article: Yukio Doi (1983) On the structure of relative hopf modules, Communications in Algebra, 11:3, 243-255, DOI: [10.1080/00927878308822847](https://doi.org/10.1080/00927878308822847)

To link to this article: <http://dx.doi.org/10.1080/00927878308822847>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

ON THE STRUCTURE OF RELATIVE HOPF MODULES

Yukio Doi

Fukui University
Fukui 910, Japan

Let A be a Hopf algebra over a field k . In this paper we study the notions of (A, B) -Hopf modules for an A -comodule algebra B and $[C, A]$ -Hopf modules for an A -module coalgebra C .

In [2], Sweedler has proved for any Hopf algebra H , that the existence of an integral $x : H \longrightarrow k$ with $x(1) = 1$ is equivalent to the complete reducibility of all H -comodules. He also has reduced the structure theorem for H -Hopf modules. Here we give an extension of these results.

Throughout we freely use the terminology and results of [2]. All vector spaces will be over a field k . Map always means k -linear map, and the unadorned tensor product $V \otimes W$ is understood to be $V \otimes_k W$. We use the sigma notation. Thus, if C is a coalgebra, we write $\Delta(c) = \sum c_{(1)} \otimes c_{(2)}$, for $c \in C$. If M is a right C -comodule with comodule structure map $\rho : M \longrightarrow M \otimes C$, we write for $m \in M$,

$$\rho(m) = \sum m_{(0)} \otimes m_{(1)}.$$

Throughout this paper A is a Hopf algebra with antipode S .

Let B be an algebra and a right A -comodule. The comodule structure map will be denoted by $\rho_B : B \longrightarrow B \otimes A$ and for $\rho_B(b)$ we write $\sum b_{(0)} \otimes b_{(1)}$. B is called a right A -comodule algebra if ρ_B is an algebra map.

A is itself a right A -comodule algebra via $\Delta : A \longrightarrow A \otimes A$. More generally, if B is a subalgebra and a right coideal of A then B becomes a right A -comodule algebra. The ground field k has a trivial right A -comodule algebra structure given by

$$u_A : k \longrightarrow A \approx k \otimes A.$$

Definition. Let B be a right A -comodule algebra. M is called a right (A, B) -Hopf module if M is a right A -comodule and a right B -module such that the following diagram commutes

$$\begin{array}{ccccc} M \otimes B & \xrightarrow{\omega} & M & \xrightarrow{\rho} & M \otimes A \\ \downarrow \rho_M \otimes \rho_B & & & & \uparrow \omega_M \otimes M_A \\ M \otimes A \otimes B \otimes A & \xrightarrow{I \otimes T \otimes I} & & & M \otimes B \otimes A \otimes A \end{array}$$

(ω_M is the B -module action of M , ρ_M is the A -comodule structure map of M , M_A is the multiplication in A , T is the twist map).

The diagram can be expressed as

$$\rho_M(mb) = \sum m_{(0)} b_{(0)} \otimes m_{(1)} b_{(1)}$$

for all $m \in M$, $b \in B$.

We note that B is itself a right (A, B) -Hopf module via ρ_B and $M_B : B \otimes B \longrightarrow B$.

Theorem 1. Let B be a right A -comodule algebra where there is a right A -comodule map $\phi : A \longrightarrow B$ with $\phi(1_A) = 1_B$. Then every right (A, B) -Hopf module is injective as an A -comodule.

Proof. Let M be a right (A, B) -Hopf module. If $M \otimes A$ has the right A -comodule structure given by $I \otimes \Delta : M \otimes A \longrightarrow (M \otimes A) \otimes A$ then the comodule structure map $\rho_M : M \longrightarrow M \otimes A$ is an A -comodule map. We show that there is an A -comodule map $\lambda : M \otimes A \longrightarrow M$ with $\lambda \rho_M = I$. Thus M is injective since it is isomorphic to a direct summand of $M \otimes A$, an injective A -comodule.

Define $\lambda : M \otimes A \longrightarrow M$ as the composite

$$\begin{aligned} M \otimes A &\xrightarrow{\rho \otimes I} M \otimes A \otimes A \xrightarrow{I \otimes S \otimes I} M \otimes A \otimes A \xrightarrow{I \otimes M_A} M \otimes A \\ &\xrightarrow{I \otimes \phi} M \otimes B \xrightarrow{\omega} M \end{aligned}$$

so that $\lambda(m \otimes a) = \sum m_{(0)} \phi(S(m_{(1)})a)$ for $m \in M$, $a \in A$.

For any $m \in M$,

$$\begin{aligned} \lambda \rho_M(m) &= \lambda(\sum m_{(0)} \otimes m_{(1)}) = \sum m_{(0)} \phi(S(m_{(1)})m_{(2)}) \\ &= \sum m_{(0)} \varepsilon(m_{(1)}) \phi(1_A) = m \end{aligned}$$

so that $\lambda \rho_M$ is the identity of M .

Next we claim λ is an A -comodule map.

$$\begin{aligned} \rho_M \lambda(m \otimes a) &= \rho_M(\sum m_{(0)} \phi(S(m_{(1)})a)) \\ &= \sum m_{(0)} \phi(S(m_{(2)})a)_{(0)} \otimes m_{(1)} \phi(S(m_{(2)})a)_{(1)} \end{aligned}$$

The condition that ϕ be a right A -comodule map is exactly

$$\rho_B \phi = (\phi \otimes I) \Delta_A \quad \text{or for } a \in A, \quad \sum \phi(a)_{(0)} \otimes \phi(a)_{(1)} = \sum \phi(a_{(1)}) \otimes a_{(2)}.$$

Since the antipode S is an anti-algebra map the above expression equals

$$\begin{aligned} & \sum m_{(0)} \phi(S(m_{(3)})a_{(1)}) \otimes m_{(1)} S(m_{(2)})a_{(2)} \\ &= \sum m_{(0)} \phi(S(m_{(2)})a_{(1)}) \otimes \epsilon(m_{(1)})a_{(2)} \\ &= \sum m_{(0)} \phi(S(m_{(1)})a_{(1)}) \otimes a_{(2)} \\ &= (\lambda \otimes I)(I \otimes \Delta_A)(m \otimes a). \end{aligned}$$

Thus λ is an A -comodule map.

q. e. d.

In case $B = k$, the above result reduces to [2, LEMMA 14.0.2].

Corollary. The following statements concerning a right A -comodule algebra B are equivalent:

- (i) B is an injective A -comodule.
- (ii) There is a right A -comodule map $\phi : A \longrightarrow B$ with $\phi(1_A) = 1_B$.

Proof. Consider the diagram of right A -comodules

$$\begin{array}{ccccc} 0 & \longrightarrow & k & \xrightarrow{u_A} & A \\ & & \downarrow u_B & \swarrow \phi & \\ & & B & & \end{array}$$

If B is an injective A -comodule then the diagram can be completed by an A -comodule map ϕ to a commutative diagram. Thus we have that (i) implies (ii).

Since B may be regarded as a right (A, B) -Hopf module it follows from Theorem 1 that (ii) implies (i). q. e. d.

Let A, A' be Hopf algebras and $f : A' \longrightarrow A$ be a Hopf algebra map. Then A' becomes a right A -comodule algebra via

$$A' \xrightarrow{\Delta} A' \otimes A' \xrightarrow{I \otimes f} A' \otimes A.$$

Theorem 2. Let $f : A' \longrightarrow A$ be a surjective Hopf algebra map. If there is a right A -comodule map $\phi : A \longrightarrow A'$ with $\phi(1_A) = 1_{A'}$, then we have:

- (1) A' is injective as a right A -comodule.
- (2) For any left A -comodule V , the canonical map

$$A' \square_A V \xrightarrow{f \otimes I} A \square_A V \simeq V$$

is surjective, where \square_A denotes the cotensor product over A .

Proof. (1) is clear by Theorem 1 and thus we need only show (2). Since f is an A -comodule map, $\text{Ker } f$ becomes a right (A, A') -Hopf module in a natural way. Thus we have from Theorem 1 that $\text{Ker } f$ is an injective A -comodule. This implies that the sequence

$$0 \longrightarrow \text{Ker } f \longrightarrow A' \xrightarrow{f} A \longrightarrow 0$$

is a split exact sequence of right A -comodules. Cotensoring over A by V yields the exact sequence

$$0 \longrightarrow (\text{Ker } f) \square_A V \longrightarrow A' \square_A V \longrightarrow A \square_A V \longrightarrow 0.$$

q. e. d.

Remark. The above Theorem shows that for surjective Hopf algebra map $f : A' \longrightarrow A$, f is right coflat if and only if it is right faithfully coflat.

We return to the first setting where B is a right A -comodule algebra.

Define the A -invariant subspace of B to be the set

$$B_0 = \{ b \in B \mid \rho_B(b) = b \otimes 1_A \}.$$

It is clear that B_0 is a subalgebra of B .

Let V be a right B_0 -module. Then $V \otimes_{B_0} B$ is a right B -module in the usual way. It is also a right A -comodule with comodule structure map $\rho : v \otimes_{B_0} b \longmapsto \sum v \otimes_{B_0} b_{(0)} \otimes b_{(1)}$ (this is well defined). One easily checks that $V \otimes_{B_0} B$ is a right (A, B) -Hopf module.

Let M be a right (A, B) -Hopf module. Define the set

$$M_0 = \{ m \in M \mid \rho_M(m) = m \otimes 1_A \}.$$

For any $m \in M_0$ and $b \in B_0$ we have $mb \in M_0$ and thus M_0 is a right B_0 -module. Define

$$\alpha : M_0 \otimes_{B_0} B \longrightarrow M$$

by $\alpha(m \otimes_{B_0} b) = mb$ for $m \in M_0$, $b \in B$. It is then an (A, B) -Hopf map, that is, an A -comodule map and a B -module map.

Theorem 3. Let B be a right A -comodule algebra. If there is a right A -comodule map $\phi : A \longrightarrow B$ which is an algebra map then for every right (A, B) -Hopf module M ,

$$\alpha : M_0 \otimes_{B_0} B \longrightarrow M$$

is an isomorphism of (A, B) -Hopf modules.

Proof. Let $P : M \longrightarrow M$ be the composite

$$M \xrightarrow{\rho} M \otimes A \xrightarrow{I \otimes S} M \otimes A \xrightarrow{I \otimes \phi} M \otimes B \xrightarrow{\omega} M.$$

Explicitly $P(m) = \sum m_{(0)} \phi(S(m_{(1)}))$.

We claim $P(M) \subset M_0$:

$$\begin{aligned} \rho_M P(m) &= \sum m_{(0)} \phi(S(m_{(2)}))_{(0)} \otimes m_{(1)} \phi(S(m_{(2)}))_{(1)} \\ &= \sum m_{(0)} \phi(S(m_{(3)})) \otimes m_{(1)} S(m_{(2)}) \\ &= \sum m_{(0)} \phi(S(m_{(2)})) \otimes \epsilon(m_{(1)}) 1_A \\ &= \sum m_{(0)} \phi(S(m_{(1)})) \otimes 1_A = P(m) \otimes 1_A. \end{aligned}$$

Thus P is in fact a map $M \longrightarrow M_0$.

Define $\beta : M \longrightarrow M_0 \otimes_{B_0} B$ by

$$\beta(m) = \sum P(m_{(0)}) \otimes_{B_0} \phi(m_{(1)}).$$

We will show $\alpha\beta = I$ and $\beta\alpha = I$:

$$\begin{aligned} \alpha\beta(m) &= \alpha(\sum m_{(0)} \phi(S(m_{(1)})) \otimes_{B_0} \phi(m_{(2)})) \\ &= \sum m_{(0)} \phi(S(m_{(1)})) \phi(m_{(2)}) \\ &= \sum m_{(0)} \phi(S(m_{(1)}) m_{(2)}) = m. \end{aligned}$$

For $m \in M_0$, $b \in B$,

$$\begin{aligned} \beta\alpha(m \otimes_{B_0} b) &= \beta(mb) = \sum P(mb_{(0)}) \otimes_{B_0} \phi(b_{(1)}) \\ &= \sum mb_{(0)} \phi(S(b_{(1)})) \otimes_{B_0} \phi(b_{(2)}) \end{aligned}$$

since $\sum b_{(0)} \phi(S(b_{(1)})) \in B_0$ for any $b \in B$

$$\begin{aligned}
&= \sum m_{B_0} b_{(0)} \phi(S(b_{(1)})) \phi(b_{(2)}) \\
&= \sum m_{B_0} b_{(0)} \phi(\epsilon(b_{(1)}) 1_B) = m_{B_0} b. \quad \text{q. e. d.}
\end{aligned}$$

In case $B = A$ and $\phi = I$, the above Theorem reduces to [2, THEOREM 4.1.1].

We dualize Theorem 1, 2 and 3.

Let C be a coalgebra which is a right A -module. C is a right A -module coalgebra if the following hold for all $c \in C$, $a \in A$:

$$\begin{aligned}
(1) \quad \Delta(ca) &= \sum c_{(1)} a_{(1)} \otimes c_{(2)} a_{(2)} \\
(2) \quad \epsilon(ca) &= \epsilon(c) \epsilon(a).
\end{aligned}$$

A is itself a right A -module coalgebra via $M_A : A \otimes A \longrightarrow A$. The ground field k has a trivial right A -module coalgebra structure.

Let N be a right C -comodule and a right A -module. N is called a right $[C, A]$ -Hopf module if the following holds for all $n \in N$, $a \in A$:

$$\rho(na) = \sum n_{(0)} a_{(1)} \otimes n_{(1)} a_{(2)}.$$

Suppose that there exists a right A -module map $\psi : C \longrightarrow A$ with $\epsilon_A \psi = \epsilon_C$. For any right $[C, A]$ -Hopf module N , define $\lambda : N \longrightarrow N \otimes A$ as the composite

$$\begin{aligned}
N &\xrightarrow{\rho} N \otimes C \xrightarrow{I \otimes \psi} N \otimes A \xrightarrow{I \otimes \Delta} N \otimes A \otimes A \\
&\xrightarrow{I \otimes S \otimes I} N \otimes A \otimes A \xrightarrow{\omega \otimes I} N \otimes A
\end{aligned}$$

so that $\lambda(n) = \sum n_{(0)} S(\psi(n_{(1)}))_{(1)} \otimes \psi(n_{(1)})_{(2)}$. If $N \otimes A$ has

the right A -module structure given by $(N \otimes A) \otimes A \xrightarrow{I \otimes M} N \otimes A$ then λ is an A -module map with $\omega\lambda = I$. Thus N is a projective A -module since it is isomorphic to a direct summand of $N \otimes A$, a free A -module.

We summarize this in the following theorem :

Theorem 4. Let C be a right A -module coalgebra where there is a right A -module map $\psi : C \longrightarrow A$ with $\varepsilon\psi = \varepsilon$. Then every right $[C, A]$ -Hopf module is a projective A -module.

Remarks. If C is finite dimensional then $\psi(C)$ is a non-zero finite dimensional right ideal of A so that A must be finite dimensional ([2], p.107). In case $C = k$, the above Theorem reduces to [2, THEOREM 5.1.8].

We state without proof the dual of Corollary of Theorem 1 and Theorem 2 :

Corollary 1. Let C be a right A -module coalgebra. The following are equivalent :

- (i) C is a projective A -module.
- (ii) There is a right A -module map $\psi : C \longrightarrow A$ with $\varepsilon\psi = \varepsilon$.

Corollary 2. Let H be a Hopf algebra and A a Hopf subalgebra. If there is a right A -module map $\psi : H \longrightarrow A$ with $\varepsilon\psi = \varepsilon$ then we have :

- (1) H is a projective A -module.
- (2) For any left A -module V , the canonical map

$$V \simeq A \otimes_A V \longrightarrow H \otimes_A V$$

is injective.

Let C be a right A -module coalgebra. If A^+ denotes the kernel of $\varepsilon : A \longrightarrow k$ then CA^+ is a coideal of C . Hence

$\bar{C} = C/CA^+$ has a unique coalgebra structure such that the projection $p : C \longrightarrow \bar{C}$ is a coalgebra map.

Let N be a right $[C, A]$ -Hopf module. Then the map p induces the right \bar{C} -comodule structure of N

$$N \xrightarrow{\rho} N \otimes C \xrightarrow{I \otimes p} N \otimes \bar{C}.$$

NA^+ is then a \bar{C} -subcomodule of N . Thus $\bar{N} = N/NA^+$ has a unique comodule structure $\bar{\rho} : \bar{N} \longrightarrow \bar{N} \otimes \bar{C}$ making the projection $\pi : N \longrightarrow \bar{N}$ a \bar{C} -comodule map, that is, $\bar{\rho}\pi = (\pi \otimes p)\rho_N$.

Note that we have $\pi(na) = \pi(n)\varepsilon(a)$, for $n \in N$, $a \in A$.

Since C has the left \bar{C} -comodule structure induced by

$$C \xrightarrow{\Delta} C \otimes C \xrightarrow{p \otimes I} \bar{C} \otimes C$$

for any right \bar{C} -comodule W , $W \otimes_{\bar{C}} C$ is defined and it is a right $[C, A]$ -Hopf module via

$$\begin{aligned} W \otimes_{\bar{C}} C &\xrightarrow{I \otimes \Delta} W \otimes_{\bar{C}} C \otimes C \\ W \otimes_{\bar{C}} C \otimes A &\xrightarrow{I \otimes \omega} W \otimes_{\bar{C}} C. \end{aligned}$$

For any right $[C, A]$ -Hopf module N , define $\alpha : N \longrightarrow \bar{N} \otimes C$ be the composite

$$N \xrightarrow{\rho} N \otimes C \xrightarrow{\pi \otimes I} \bar{N} \otimes C.$$

It is easy to see that $\alpha(N) \subset \bar{N} \otimes_{\bar{C}} C$. Thus α is in fact a map $N \longrightarrow \bar{N} \otimes_{\bar{C}} C$. α is then a $[C, A]$ -Hopf module map.

In these terms Theorem 3 can be dualized as follows :

Theorem 5. Let C be a right A -module coalgebra. If there is a right A -module map $\psi : C \longrightarrow A$ which is a coalgebra map then

for every right $[C, A]$ -Hopf module N ,

$$\alpha : N \longrightarrow \bar{N} \otimes_{\bar{C}} C$$

is an isomorphism of $[C, A]$ -Hopf modules.

Since the proof of Theorem 3 is not so easily dualized we include a proof of Theorem 5.

Proof. Let $Q : N \longrightarrow N$ denote the composite

$$N \xrightarrow{\rho} N \otimes C \xrightarrow{I \otimes \psi} N \otimes A \xrightarrow{I \otimes S} N \otimes A \xrightarrow{\omega} N$$

so that $Q(n) = \sum n_{(0)} S(\psi(n_{(1)}))$ for $n \in N$.

For $n \in N$ and $a \in A$,

$$\begin{aligned} Q(na) &= \sum n_{(0)} a_{(1)} S(\psi(n_{(1)} a_{(2)})) \\ &= \sum n_{(0)} a_{(1)} S(\psi(n_{(1)}) a_{(2)}) \\ &= \sum n_{(0)} a_{(1)} S(a_{(2)}) S(\psi(n_{(1)})) \\ &= \sum n_{(0)} \epsilon(a) S(\psi(n_{(1)})) = Q(n\epsilon(a)). \end{aligned}$$

Hence Q vanishes on NA^+ . Thus there is a map \bar{Q} making

$$\begin{array}{ccc} N & \xrightarrow{Q} & N \\ & \searrow \pi & \nearrow \bar{Q} \\ & \bar{N} & \end{array}$$

commute. In particular, if we define $Q_0 : C \longrightarrow C$ by $Q_0(c) = \sum c_{(1)} S(\psi(c_{(2)}))$ then Q_0 factors through \bar{C} , that is, there is a map $\bar{Q}_0 : \bar{C} \longrightarrow C$ with $Q_0 = \bar{Q}_0 p$. Note that we have

$$\omega_C(Q_0 \otimes \psi) \Delta_C = I_C.$$

Let $\beta : \bar{N} \square_{\bar{C}} C \longrightarrow N$ denote the composite

$$\bar{N} \square_{\bar{C}} C \xrightarrow{\text{inclusion}} \bar{N} \boxtimes C \xrightarrow{\bar{Q} \boxtimes \psi} N \boxtimes A \xrightarrow{\omega} N.$$

For any $n \in N$

$$\begin{aligned} \beta \alpha(n) &= \beta(\Sigma \pi(n_{(0)}) \boxtimes n_{(1)}) \\ &= \Sigma n_{(0)} S(\psi(n_{(1)})) \psi(n_{(2)}) \\ &= \Sigma n_{(0)} S(\psi(n_{(1)})_{(1)}) \psi(n_{(1)})_{(2)} \\ &= \Sigma n_{(0)} \varepsilon \psi(n_{(1)}) = \Sigma n_{(0)} \varepsilon(n_{(1)}) = n. \end{aligned}$$

For any $\pi(n) \in \bar{N}$, $c \in C$

$$\begin{aligned} \alpha \omega_N(\bar{Q} \boxtimes \psi)(\pi(n) \boxtimes c) &= (\pi \boxtimes I) \rho(Q(n) \psi(c)) \\ &= (\pi \boxtimes I)(\Sigma Q(n)_{(0)} \psi(c)_{(1)} \boxtimes Q(n)_{(1)} \psi(c)_{(2)}) \\ &= (\pi \boxtimes I)(\Sigma n_{(0)} S(\psi(n_{(3)})) \psi(c_{(1)}) \boxtimes n_{(1)} S(\psi(n_{(2)})) \psi(c_{(2)})) \\ &= \Sigma \pi(n_{(0)}) \varepsilon(n_{(3)}) \varepsilon(c_{(1)}) \boxtimes n_{(1)} S(\psi(n_{(2)})) \psi(c_{(2)}) \\ &= \Sigma \pi(n_{(0)}) \boxtimes n_{(1)} S(\psi(n_{(2)})) \psi(c) \\ &= \Sigma \pi(n_{(0)}) \boxtimes Q_0(n_{(1)}) \psi(c). \end{aligned}$$

Let $\Sigma \pi(n) \boxtimes c \in \bar{N} \square_{\bar{C}} C$, thus

$$(\bar{\rho} \boxtimes I)(\Sigma \pi(n) \boxtimes c) = (I \boxtimes p \boxtimes I)(I \boxtimes \Delta)(\Sigma \pi(n) \boxtimes c).$$

Since $\bar{\rho} \pi = (\pi \boxtimes p) \rho_N$ we have

$$(\pi \boxtimes p \boxtimes I)(\rho_N \boxtimes I)(\Sigma n \boxtimes c) = (I \boxtimes p \boxtimes I)(I \boxtimes \Delta)(\Sigma \pi(n) \boxtimes c).$$

Applying $(I \boxtimes \omega_C)(I \boxtimes \bar{Q}_0 \boxtimes \psi)$ to this, we have

$$\Sigma \pi(n_{(0)}) \boxtimes Q_0(n_{(1)}) \psi(c) = \Sigma \pi(n) \boxtimes c.$$

Thus we have shown that $\alpha\beta$ is the identity on $\bar{N} \square_{\bar{C}} C$. q.e.d.

REFERENCES

- [1] Y. Doi, "Homological coalgebra", J. Math. Soc. Japan 35, 1981, 31-50.
- [2] M. E. Sweedler, "Hopf Algebras", Benjamin, New York, 1969.
- [3] M. Takeuchi, "Relative Hopf Modules - Equivalences and Freeness Criteria -", J. Algebra, 60, 1979, 452-471.

Received: October 1981