
a probable contamination: ≥90% identity
other proteins (105)
probable contamination (11)
count
best bacterial hit, % identity
b bacterial-type: ≥35/50 non-fungal hits
single bacterial hit (43)
bacterial-type (62)
non-fungal among 50 best hits
c bacterial-type: margin ≥50 bits
bit-score margin vs fungal hit
d T-DNA candidate: seed identity ≥35%
bacterial-type, other (18)
T-DNA candidate (44)
identity to T-DNA seed, %
e T-DNA candidate: coverage ≥0.4
coverage of T-DNA seed
f T-DNA candidate: ratio ≥0.9
seed id. / best bacterial id.
g non-T-DNA candidate: Rhizobiales hit ≥50%
non-T-DNA candidate (6)
best Rhizobiales hit, % identity
h chs-like proteins: ≥45% identity to seed
native homologues (150)
chs lineage (44)
identity to chs seed, %
i chs-like loci: ≥70% to F. vanettenii
pass via chs seed (15)
pass only via 70% criterion (18)
identity to F. vanettenii, %
j pKerr14a-type fragments: ≥60% identity
not assigned (238)
assigned fragments (13)
identity to nearest intact, %
k vir/tra/rep, test contigs: ≥60% identity
fungal locus contigs (43)
contaminant contigs (11)
best id. to vir/tra/rep protein, %
l vir/tra/rep, Boletales: ≥60% identity
other Boletales (130)
B. harrisonii (1)
max vir/tra/rep id., %