

FUNCTION APPROXIMATION

QUANTITATIVE ECONOMICS 2023

Piotr Żoch

October 23, 2023

WHAT IS IT?

Goal: approximate a complicated function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ by a simpler function $\hat{f} : \mathbb{R}^n \rightarrow \mathbb{R}$.

- We know f only at a finite number of points and we want to approximate it at other points x .
- f is too complicated to work with directly (e.g. non-analytic) and we need to represent on a computer.

I will talk only about the **most basic** ideas.

FUNCTION APPROXIMATION

- What data should be produced and used?
- What family of “simpler” functions should be used?
- What notion of approximation do we use?
- How good can the approximation be?
- How simple can a good approximation be?

Notice **similarities** and **differences** between function approximation and statistical regression.

NOTATION

- Today we will focus on continuous functions $f : \mathbb{R}^n \rightarrow \mathbb{R}$.
- We can represent every continuous function in a particular function space by a linear combination of **basis functions**.
- Analogy: Every vector in a vector space V can be represented by a linear combination of basis vectors.

NOTATION

- Let F be the space of continuous real-valued functions with domain $X \subset \mathbb{R}^n$.
- Define the **inner product** of two functions $f, g \in F$ as

$$\langle f, g \rangle = \int_X f(x)g(x)w(x)dx$$

where $f, g, w \in F$, w is a **weighting function**.

- $\{F, \langle \cdot, \cdot \rangle\}$ is an inner product space.
- We want to approximate a **known** function $f : X \rightarrow \mathbb{R}$ in $\{F, \langle \cdot, \cdot \rangle\}$.

NOTATION

- Let $\hat{f}(\cdot, \beta)$ be a parametric approximation of f . We have

$$\hat{f}(x, \beta) = \sum_{j=0}^J \beta_j \phi_j(x)$$

- $\phi_j(x)$ are **basis functions**. Write $\Phi_J = \{\phi_0, \phi_1, \dots, \phi_J\}$.
 - $\beta = [\beta_0, \beta_1, \dots, \beta_J]$ is a vector of coefficients.
 - J is the order of interpolation.
- We want to find β such that $\hat{f}(\cdot, \beta)$ is a "good" approximation of f .
- Define the residual function $r(x, \beta) = f(x) - \hat{f}(x, \beta)$. We want to make it small in some sense.

SPECTRAL METHODS

SPECTRAL METHODS

- **Spectral methods:** basis functions are non-zero on the entire domain of f .
- Polynomial interpolation: basis functions are polynomials
- Fourier series: basis functions are sines and cosines

INTERPOLATION

- Suppose we know f at $N = J + 1$ points $\{x_i\}_{i=0}^J$. We call these points **interpolation nodes**.
- Note: we have the **same number** of points as basis functions.
- Let r_i be the residual at x_i . We have

$$\begin{bmatrix} r_0 \\ \vdots \\ r_J \end{bmatrix} = \begin{bmatrix} f(x_0) \\ \vdots \\ f(x_J) \end{bmatrix} - \begin{bmatrix} \phi_0(x_0) & \cdots & \phi_0(x_J) \\ \vdots & \ddots & \vdots \\ \phi_J(x_0) & \cdots & \phi_J(x_J) \end{bmatrix} \begin{bmatrix} \beta_0 \\ \vdots \\ \beta_J \end{bmatrix}$$

- Abuse notation and write it as

$$r = f - \Phi\beta$$

INTERPOLATION

- The idea of **interpolation** is to find β such that $r_i = 0$ for all i .
- The unknowns are the coefficients β .
- This is equivalent to solving the linear system of equations

$$\Phi\beta = f$$

with an obvious solution $\beta = \Phi^{-1} f$.

REGRESSION

- If we have more points than basis functions, $M > J + 1$ we cannot interpolate.
- We have to define a loss function $L(\cdot, r)$ and minimize it.
- If we use the squared loss function

$$L(x, r) = \sum_{i=0}^M r(x_i, \beta)^2,$$

we get the **least squares** problem with solution

$$\beta = \left(\Phi^T \Phi \right)^{-1} \Phi^T f$$

WEIERSTRASS THEOREM

Theorem

Let $f : [a, b] \rightarrow \mathbb{R}$ be a continuous function. Then for every $\varepsilon > 0$ there exists a polynomial p such that

$$\sup_{x \in [a, b]} |f(x) - p(x)| \leq \varepsilon.$$

- We can approximate any continuous function on a compact set by a **polynomial** as closely as we want.

BASIS FUNCTION CHOICE

- Which basis functions should we use?
- The most natural choice seems to be **monomials**: $\phi_j(x) = x^j$.
- Problem: consecutive monomials are very similar to each other – does x^{10} really add much to x^9 ?
- The resulting matrix Φ is a **Vandermonde matrix**, which is **ill-conditioned**.

BASIS FUNCTION CHOICE

- If Φ look like a diagonal matrix, then Φ^{-1} is easy to compute.
- Intiuition: use basis functions that give us **different** information about f .
- **Orthogonal polynomials** are a good choice: $\langle \phi_i, \phi_j \rangle = 0$ for $i \neq j$.

ORTHOGONAL POLYNOMIALS

- For orthogonal polynomials, we have

$$\beta_j = \int_X f(x)\phi_j(x)w(x)dx$$

- Intuition: use basis functions that give us **different** information about f .
- **Orthogonal polynomials** are a good choice: $\langle \phi_i, \phi_j \rangle = 0$ for $i \neq j$.
- Examples: Legendre, Chebyshev, Hermite, Laguerre, Jacobi polynomials.

Chebyshev Polynomials

- Chebyshev polynomials $T_n(x) : [-1, 1] \rightarrow \mathbb{R}$ are given by

$$T_0(x) = 1$$










$$T_1(x) = x$$

$$T_{j+1}(x) = 2xT_j(x) - T_{j-1}(x)$$

- Let $X = [-1, 1]$ and $w(x) = \frac{1}{\sqrt{1-x^2}}$. We have

$$\langle T_i, T_j \rangle = \int_{-1}^1 T_i(x) T_j(x) \frac{1}{\sqrt{1-x^2}} dx = \begin{cases} 0 & i \neq j \\ \pi & i = j = 0 \\ \frac{\pi}{2} & i = j \neq 0 \end{cases}$$

Chebyshev POLYNOMIALS



Chebyshev Nodes

- But it is not the only nice property of Chebyshev polynomials.
- **Chebyshev nodes** are roots of Chebyshev polynomials on $[-1, 1]$:

$$x_i = \cos\left(\frac{2i-1}{2n}\pi\right) \quad \text{for } i = 1, \dots, n.$$

It can be verified that T_n equals 0 at these points.

- Chebyshev nodes are not equally spaced, they are **clustered** at the endpoints of the interval.

NODES CHOICE

- Why are these nodes useful?
- We know that at the interpolation nodes x_i we have $r_i = 0$.
- We want to make the residual as small as possible at other points x .
- A silly choice of nodes (e.g. equidistant) can lead to a large residual at other points, even with a high order of interpolation – **Runge phenomenon**.
- **Minmax approximation**: polynomial approximation using Chebyshev nodes minimizes the maximum absolute error on $[-1, 1]$.

FINITE ELEMENTS METHOD - NEXT WEEK
