SRG Origins

From Geometry to Recursive Operators

A Step-by-Step Mathematical History of Constants,
TGMO, RPCO, REFU, and the Early SRG Models


Hilmir Frímann Halldórsson

v0.1 · 11 October 2026 · Final-review edition

This paper traces a mathematical history: geometry and constants, three recursive operators, and their later adaptation into SRG. Four introductory pages lead to complete proofs and source references. Historical definitions, deductions, later diagnostic choices and unresolved questions remain distinct.

Walkthrough 1 of 4: begin with two circles

Derived geometry. Draw two circles of radius one length unit, with their centres one unit apart. Their upper intersection is one unit from both centres. Joining these three points makes an equilateral triangle.



$R = AB = AC = BC = 1$

$h = \sqrt{3}/2$

$r = h/3 = \sqrt{3}/6$

Circle centres: A, B. Triangle inradius: r. Units: length.

The area is both half base times height and inradius times semiperimeter. Consequently

$$\boxed{\left(\frac{r}{R}\right)^2 = \frac{1}{12}.}$$


Here R is the *original circle radius*, also the triangle's side. It is not the triangle's circumradius; that denominator would give $1/4$. Uniform resizing preserves the ratios.

Four constants, four constructions. π is circumference divided by diameter. $e = \lim(1+1/n)^n$ comes from finer and finer compounding. $\varphi = (1+\sqrt{5})/2$ is a pentagon's diagonal-to-side ratio and a Fibonacci-ratio limit. $\zeta(3) = 1 + 1/2^3 + 1/3^3 + \cdots$ is the convergent inverse-cube sum. One circle picture does not uniquely select their product.

Reading labels. Historical: found in a source. **Derived**: proved under stated assumptions. **Later diagnostic**: a measurement selected during later analysis. **Illustrative**: example values selected for explanation. Full derivations: technical sections 3–5.

Walkthrough 2 of 4: one loop, three questions

Later diagnostic; derived calculation; illustrative six-site example. Place six sites on a loop of circumference one. A field is six numbers. The neighbour difference is $\Delta u_j = u_{j+1} + u_{j-1} - 2u_j$, with indices wrapping around. Choose the positive operator $A_6 = 6^2(-\Delta)$.



Constant mode: 0 (removed)

Nonzero squared frequencies:

36, 108, 144, 108, 36

The repeated values count both directions around the loop. Keep the *whole* nonzero spectrum. Ask three questions about this same list:

1. **Add inverses.**
$$Z_6(1) = \frac{2}{36} + \frac{2}{108} + \frac{1}{144} = \frac{35}{432} \approx 0.081019.$$
At N sites the exact answer is $(1 - N^{-2})/12$. As the loop is refined this approaches $\boxed{+1/12}$.

2. **Add inverse powers $3/2$.**
$$Z_6(3/2) = \frac{2}{36^{3/2}} + \frac{2}{108^{3/2}} + \frac{1}{144^{3/2}} \approx 0.011620.$$
The limiting answer is $\boxed{\zeta(3)/(4\pi^3)}$. Squared frequencies grow like k^2; the power $-3/2$ produces k^{-3}.

3. **Add frequencies and subtract a bulk integral.**
$$S_6 = \frac{1}{4\pi} \sum_{k=1}^{5} \sqrt{\lambda_{6,k}} = \frac{3(2+\sqrt{3})}{\pi}, \qquad S_6 - \frac{36}{\pi^2} \approx -0.083717.$$
The limiting remainder is $\boxed{-1/12}$, with the subtraction fixed.

Why a negative answer? The samples of a concave sine arch lie below its continuous area. Every frequency is positive; the negative number is the difference after subtracting a larger quantity. It does not make the ordinary sum $1 + 2 + 3 + \cdots$ convergent or negative.

The triangle ratio and limiting inverse trace give the same positive number by different measurements. The negative finite part comes from a third measurement. These choices do not invalidate other regularizations. Technical section 5 proves the formulas and states the scale and domains.

Walkthrough 3 of 4: how the historical lock enters

Historical definition and interpretation. The July fusion manuscript, *Symbolic Recursion and SRG Fusion* (source P01, pp.3–4), writes

$$\pi e\varphi = \frac{\zeta(3)}{\gamma}, \qquad \boxed{\gamma_* = \frac{\zeta(3)}{\pi e\varphi} \approx 0.0869947475.}$$

Appendix A, pp.19–20, gives the author's design interpretation: circle geometry and growth, golden packing or growth patterns, discrete lattice sums, and a scaling/calibration factor. It reports approximately 0.087 and distinguishes the relation from a first-principles physical derivation. This rationale is part of the source history.

Later diagnostic choice. The loop calculation supplies $\zeta(3)/(4\pi^3)$. To recover the historical target, the October analysis explicitly chooses the multiplier

$$\underbrace{\frac{4\pi^2}{e\varphi}}_{\text{selected normalization}} \underbrace{\frac{\zeta(3)}{4\pi^3}}_{\text{loop limit}} = \frac{\zeta(3)}{\pi e\varphi}.$$

The equality is exact. The multiplier imports the historical target; the loop did not independently select it. No independently specified `GammaEstimate` or saved independent coupling measurement was recovered. One later R/D engine uses approximately 0.08699 as an input.

A genuine classical bridge. There is an exact finite-interval identity,

$$\boxed{\zeta(3) = 10 \int_0^{\log\varphi} t^2 \coth t \, dt.}$$

Here $\coth t = (e^t + e^{-t})/(e^t - e^{-t})$. Although $\coth t$ grows near zero, $t^2 \coth t$ tends to zero; the integral converges. Its golden endpoint makes logarithmic terms cancel in the trilogarithm identity proved in technical section 4.4. This is established mathematics, with classical attribution, not a new project discovery or an independent selection of the lock's denominator.

Three statements to keep separate. The formula was historically written. The identities used to evaluate it are provable. Whether a separately defined physical or native-state measurement selects it is a further question. Technical section 6 follows each of these statements to the papers and saved records.

Symbols matter. The exponential base e, Euler–Mascheroni's constant $\gamma_{\mathrm{EM}} \approx 0.5772$, and the solved lock target $\gamma_* \approx 0.0870$ are different numbers. A July glossary conflates the last two; the discrepancy remains documented, without changing the historical source.


Walkthrough 4 of 4: from a field now to a field next

Historical January composition; illustrative parameters. An operator is a rule transforming an input. On a periodic grid, opposite edges are neighbours. Begin with

$$\Theta = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \qquad \mathcal{L}\Theta = \begin{pmatrix} 0 & 1 & 0 \\ 1 & -4 & 1 \\ 0 & 1 & 0 \end{pmatrix}.$$

TGMO supplies the neighbour difference. REFU composes it with a nonlinear contribution reading the *same incoming grid*:

$$P = \Theta + .1\mathcal{L}\Theta + .1\tanh\Theta = \begin{pmatrix} 0 & .1 & 0 \\ .1 & c & .1 \\ 0 & .1 & 0 \end{pmatrix}, \quad c = .6 + .1\tanh 1 \approx .6761594.$$



Here $a = b = 1$, $h = \kappa = .1$, $s = 1$, H is off, $L = 9$ and $\varepsilon = 10^{-9}$. The five positive entries are on inactive compression branches and each gains ε. **Paper mode turns all four zero corners into one; signed mode keeps them zero.** Thus the mode changes the next state even in this example.

January's names are *Toroidal Grid Modulation Operator*, *Recursive Phase-Compression Operator* and *Recursive Field Update*. The July prose and standalone files used some names for different functions. March programs explicitly adapted them to moving particles, neighbour graphs and resonance values. An R/D pair adds two real fields with unequal gains; the names alone imply neither time reversal nor complex conjugation.

Recursion uses a result in a subsequent update. A clock-prescribed H curve does not itself accumulate state memory. July's unresolved implicit temporal equation, January's fixed-grid theorems and March's particle results retain their own assumptions. Technical sections 7–8 give the formulas; section 9 identifies what remains open. No source step or simulation was run for this illustration.

Technical exposition and source history

The following sections retain the accepted lineage report's mathematical equation numbers (1)–(29). The source-and-notation addendum corrects three SRG-I locators, the named torus angles and Tower II's inactive-row guards. The author's Appendix A design explanation and public source map are included. The accepted report and all original source bytes remain unchanged.

How to read the history. The July fusion manuscript P01 uses Symbolic Resonance Geometry; the March paper P18 uses Symbolic Resonant Geometry. The shared acronym does not erase changes in state or update rules. Here a *state* is the mathematical input to an update, a *stencil* is a fixed neighbour-difference formula, a *spectrum* is the list of an operator's eigenvalues, and *compression* is a pointwise amplitude transformation. The technical definitions below make those meanings precise.

Contents

1. A worked route a reader can follow **6**

2. Evidence conventions and chronology **7**

3. Elementary geometry, φ, and e **8**

4. From repeated integration to $\zeta(3)$ **10**

5. The three loop observables and the two signs of $1/12$ **12**

6. The lock equation: what was printed, assigned, or measured **14**

7. TGMO, REFU and RPCO: precise operator lineage **16**

8. The March adaptations **19**

9. Claim disposition and remaining definition **21**

10. Reading the evidence responsibly **22**

11. Source keys and reading map **23**

1. A worked route a reader can follow

Draw two circles of radius one whose centres are one unit apart. A point where they intersect is one unit from either centre. Those three points therefore form an equilateral triangle. Its height is $\sqrt{3}/2$; its inscribed circle has radius $\sqrt{3}/6$. Squaring that radius gives $1/12$. No infinite series or physical model is needed.

Now consider the quite different question of what patterns fit around a loop. A smooth wave can fit once, twice, three times, and so on. The squared frequencies grow like $1^2, 2^2, 3^2, \ldots$. Taking inverse powers $3/2$ turns those squared frequencies into $1, 1/8, 1/27, \ldots$. Their convergent sum is $\zeta(3)$. A finite cycle makes this calculation precise, including its scale and the pair of wave directions.

A third calculation adds positive frequencies and subtracts their continuous bulk approximation. The finite remainder tends to $-1/12$. Its sign comes from the subtraction: the sampled sine arch lies below the continuous arch. It is not a negative ordinary sum of positive numbers.

These results fit within one mathematical account, but they are different measurements. The historical SRG lock,

$$\pi e\varphi = \frac{\zeta(3)}{\gamma}, \qquad \gamma_* = \frac{\zeta(3)}{\pi e\varphi} = 0.086994747517035002497523657 48\ldots, \tag{1}$$

introduces another choice. The loop calculation supplies $\zeta(3)/(4\pi^3)$. To obtain (1), the October analysis multiplies by $4\pi^2/(e\varphi)$. That multiplier carries the historical target into the new diagnostic. The recovered sources do not determine it from the equilateral triangle or from a measured field statistic.

Finally, a numerical operator answers a different question: "given the current field, what is its next value?" In the January composition, neighbouring values first contribute a discrete Laplacian; a nonlinear contribution reads the same incoming field; a prescribed function $H(t)$ multiplies the result; compression is last. Section 7 works through an actual 3×3 mathematical example without executing a source step.

Main finding. The exact mathematical connections are substantial and provable. The historical record does not establish one unchanged recursion running from circle geometry to the lock to all later SRG implementations. It records several definitions, some incompatible descriptions, and explicit later adaptations. This report preserves those distinctions.

2. Evidence conventions and chronology

H means recovered historical definition or claim, **D** a deduction with assumptions stated here, **A** a choice introduced in later analysis, and **U** an unresolved specification or provenance gap. "Historical claim" does not certify its truth.

Page references below are physical PDF pages, counted from one. P01–P19 retain the accepted R0 source identifiers. SOURCE_INDEX.md gives human-readable titles, public access routes where verified, exact source hashes and access limits. The source-key map at the end of this paper resolves every abbreviation used below. Code citations count the first line as one, including a possible BOM. Historical source programs were read as text, not executed.

29 June 2025, internal date. *Source:* P08/P09. *Mathematical development:* Golden-scaled nonlinear damping, phase/tensor feedback modules.

Undated cover; June 2025 metadata is only a locator. *Source:* P13, p.5. *Mathematical development:* A distinct "Master Recursive Dual-Phase Collapse Operator".

11 July 2025. *Source:* P07, pp.3–6. *Mathematical development:* Logarithmic RPCO, spatial TGMO formula, REFU and prescribed H.

13 July 2025. *Source:* P06, pp.4,6. *Mathematical development:* Earliest internally dated lock equation located in this bounded primary set; inconsistent identification of its γ.

14 July 2025. *Source:* P03, pp.2,5,7–8. *Mathematical development:* Explicit resonance rule, consistency relation and implicit temporal trail rule.

15 July 2025. *Source:* P01, pp.3–4; P05. *Mathematical development:* Lock claimed empirical; patch supplies a different position engine and standalone operators.

22 July 2025, cover. *Source:* Rainman. *Mathematical development:* Geometric analogies and a proposed cycle/period construction.

6 September 2025. *Source:* P19. *Mathematical development:* Run-dependent coupling claim, lock penalty, unspecified GammaEstimate.

25 January 2026, cover. *Source:* JANUARY, pp.2–3 and accompanying code. *Mathematical development:* Fully specified fixed-grid composition with two RPCO modes.

14 March 2026, internal headers. *Source:* P18 and March programs. *Mathematical development:* Particle adaptation, R/D coupling, assigned frequency tower and static zeta predictions.

9 October 2026. *Source:* Received Geometry–Zeta Investigation 01. *Mathematical development:* Auxiliary normalized loop and three distinct spectral observables.

October 2026. *Source:* Accepted F01, SRG-I, OC01–02. *Mathematical development:* Source recovery and mathematical analysis; no retroactive alteration of historical equations.

P06 cites earlier work that was not recovered as a target manuscript by the named-reference follow-up. Therefore "earliest located" is not a claim of absolute authorship priority. Dates are not inferred from the current copy time.

3. Elementary geometry, φ, and e

3.1 Circles and angle projections [D; H comparison: Rainman pp.3–4]

A circle is $(x, y) = R(\cos\theta, \sin\theta)$. Its angle gives separate horizontal and vertical projections. For example,

$$\cos 45^\circ = \sin 135^\circ = \frac{\sqrt{2}}{2}, \quad \cos 45^\circ \sin 135^\circ = \frac{1}{2}. \tag{2}$$

For equal circles centred at $(0, 0)$ and $(d, 0)$, subtract $x^2 + y^2 = R^2$ and $(x - d)^2 + y^2 = R^2$. For $0 < d \leq 2R$ their intersection has

$$x = d/2, \qquad y = \pm\sqrt{R^2 - d^2/4}. \tag{3}$$

Thus the midpoint is forced by symmetry. The choice $d = R$ gives the canonical vesica and a 60° equilateral triangle; $d = \sqrt{2}R$ gives a different, 45° intersection geometry. They should not be substituted for one another.

Let a be the equilateral triangle's side. Its altitude is $h = \sqrt{3}a/2$. Area equals both $ah/2$ and inradius times semiperimeter $3ar/2$, so $r = h/3$. Consequently

$$\left(\frac{r}{a}\right)^2 = \frac{1}{12}. \quad \text{For the canonical two-circle triangle, } a = R. \tag{4}$$

Here R denotes the original circles' radius, **not** the triangle's circumradius. The latter is $a/\sqrt{3}$, and the inradius-to-circumradius squared is $1/4$. This notation distinction is essential.

Uniform resizing does not change (4). In particular $\gamma_* - 1/12 = 0.00366141418\ldots \neq 0$. The accepted October geometry note distinguishes its side-one triangle from the octagon edge and full shell height; none are silently identified here.

Rainman's comparison $x = 1/2 \leftrightarrow \Re s = 1/2$ (p.3) assigns the same coordinate to two settings. It provides no map relating circle intersections to zeros of the analytic zeta function. An equal coordinate is not a proof about zeros.

3.2 Golden ratio: a particular proportion and a genuine recurrence [D]

For four consecutive vertices of a regular pentagon, Ptolemy's cyclic-quadrilateral identity gives $d^2 = s^2 + sd$, with side s and diagonal d. Dividing by s^2 gives $q^2 = q + 1$, whose positive root is $\varphi = (1 + \sqrt{5})/2$. The identity also follows by drawing the intersecting diagonals and using the resulting similar triangles. This is additional pentagonal geometry; (3) alone does not force it.

For completeness, the algebraic recursion $F_{n+1} = F_n + F_{n-1}$, $F_0 = 0, F_1 = 1$, has characteristic roots φ and $\psi = -\varphi^{-1}$. Direct substitution and the two initial conditions prove

$$F_n = \frac{\varphi^n - \psi^n}{\sqrt{5}}, \qquad \frac{F_{n+1}}{F_n} = \varphi\frac{1 - (\psi/\varphi)^{n+1}}{1 - (\psi/\varphi)^n} \longrightarrow \varphi. \tag{5}$$

The convergence follows from $|\psi/\varphi| < 1$, not merely from assuming a limit and solving its equation.

3.3 Exponential growth and decay [D; H use: P07 p.6]

Compounding a unit growth rate in n equal pieces gives $(1 + 1/n)^n$. The integral definition of logarithm yields $x/(1 + x) \le \log(1 + x) \le x$ for $x > 0$. Hence $n/(n + 1) \le n \log(1 + 1/n) \le 1$, proving the limit e. The series $\exp t = \sum_{k\ge0} t^k/k!$ differentiates to itself; $\exp(-bt)$ solves $u' = -bu$. This explains the mathematical role of exponentials in prescribed decay curves. It does not select the product $\pi e\varphi$ as a physical coupling.

The base e, Euler–Mascheroni constant γ_{EM}, and the lock value γ_* are three different numbers.

4. From repeated integration to $\zeta(3)$

4.1 A convergent construction [D; H: Rainman p.23]

For $0 \le x < 1$, the geometric series gives $f_0(x) = x/(1-x) = \sum_{n\ge1} x^n$. Define $(If)(x) = \int_0^x f(t)\,dt/t$. Termwise integration is justified by uniform convergence on compact subintervals, or by positivity. Each integration divides the nth coefficient by n:

$$I^k f_0(x) = \sum_{n\ge1} \frac{x^n}{n^k} = \mathrm{Li}_k(x), \qquad \lim_{x\uparrow1} I^3 f_0(x) = \sum_{n\ge1} \frac{1}{n^3} = \zeta(3). \tag{6}$$

The endpoint follows by monotone convergence; $\sum n^{-3}$ is finite by the integral comparison test. This is repeated application of an integration operator. It is not an evolving SRG state, and the choice of three integrations does not derive three spatial dimensions.

Writing all integrations gives

$$I(a) = \int_0^a \frac{dx_1}{x_1} \int_0^{x_1} \frac{dx_2}{x_2} \int_0^{x_2} \frac{dx_3}{1-x_3} = \mathrm{Li}_3(a), \quad 0 < a \le 1. \tag{7}$$

This validates the scalar iterated integral on Rainman p.23 at $a = 1$, with the orientation chosen to agree with positive iterated integration. A differential-form presentation must choose the matching chain orientation.

4.2 What changes on the scaled domain [D correction to H: Rainman p.24]

Page 24 sets $a = m_{\max}\gamma$ and $x_i = au_i$. Both the domain **and the differential form** must transform:

$$\frac{dx_3}{1-x_3}\frac{dx_2}{x_2}\frac{dx_1}{x_1} = \frac{a\,du_3}{1-au_3}\frac{du_2}{u_2}\frac{du_1}{u_1}. \tag{8}$$

The factor a in the numerator is required. The result is $\mathrm{Li}_3(a)$, not generally $\zeta(3)$. At $a = 1/2$, (7) is $0.5372131936080402\ldots$, versus $\zeta(3) = 1.202056903159594\ldots$. For $0 < a < 1$ strict inequality follows term by term from (6). For $a > 1$ the pole at $x_3 = 1$ obstructs the ordinary integral. A principal value or analytic continuation would require a newly specified prescription.

Using $dx_3/(a - x_3)$ instead would make the pulled-back form standard. That is a **new scale-dependent measure**, not a consequence of the historical formula. This report does not adopt that completion.

4.3 The proposed geometric bridge [H/U]

Rainman p.18 and p.20 define their construction using $\gamma = \gamma_*$ at the outset. They cannot then serve as an independent derivation selecting that same constant.

Page 20 chooses an affine translation by γ inside projective space. Its mth power is a translation by $m\gamma$. In the concrete p.21 map $T_\gamma([x : y : z]) = [x + \gamma z : y : z]$, the forward image of $x = 0$ satisfies $x - \gamma z = 0$. The printed $x + \gamma z = 0$ is its preimage instead. Substituting $(0, y, z)$ into the displayed transformation proves the sign distinction directly.

Nor is the asserted three-dimensional integration chain on p.24 derived from the intersections described there. A curve intersected with a hyperplane in the p.21 example gives points; the text

has not supplied a map and dimension/orientation argument producing the real ordered simplex in X^3 used on p.23. Non-singularity of a matrix of selected classes alone does not prove they span the intended ambient space: the single vector $(1, 0)$ has non-singular Gram matrix [1] but does not span $\mathbb{R}^2$. These are specific gaps in the proposed bridge, not a rejection of the valid integral (7).

4.4 An exact golden-ratio relation [D; classical, reused by October analysis]

Let $q = \varphi^{-1}$ and $\ell = \log \varphi$, so $q + q^2 = 1$. The following identity genuinely connects the golden split with the trilogarithm:

$$\mathrm{Li}_3(q^2) = \frac{4}{5}\zeta(3) - \frac{2\pi^2}{15}\ell + \frac{2}{3}\ell^3. \tag{9}$$

Here is a full derivation, independent of the proposed vesica chain. For $0 < x < 1/2$, differentiating the defining series proves $\mathrm{Li}_s'(x) = \mathrm{Li}_{s-1}(x)/x$ and $\mathrm{Li}_1(x) = -\log(1-x)$. Differentiating and taking the limit at zero then prove the two dilogarithm identities

$$\begin{aligned} \mathrm{Li}_2(x) + \mathrm{Li}_2(1-x) &= \zeta(2) - \log x \log(1-x), \\ \mathrm{Li}_2(x) + \mathrm{Li}_2(-x/(1-x)) &= -\tfrac{1}{2}\log^2(1-x). \end{aligned} \tag{10}$$

Put $L = \log(1-x)$. The derivative of $\mathrm{Li}_3(x) + \mathrm{Li}_3(1-x) + \mathrm{Li}_3(-x/(1-x))$ reduces by (10) to $[-\zeta(2) + (\log x)L]/(1-x) - L^2/[2x(1-x)]$. That is exactly the derivative of $\zeta(3) + \zeta(2)L - \frac{1}{2}\log x\, L^2 + \frac{1}{6}L^3$. Both have limit $\zeta(3)$ at zero, fixing the constant. Set $x = q^2$. The series also gives $\mathrm{Li}_3(q) + \mathrm{Li}_3(-q) = \frac{1}{4}\mathrm{Li}_3(q^2)$ by cancelling odd powers. Substitution, $L = -\ell$ and $\log x = -2\ell$, proves (9).

These are standard polylogarithm identities; definitions and the dilogarithm identities can be checked against NIST DLMF section 25.12. The derivation above specifies the real domain and integration constant.

The companion identity is $\mathrm{Li}_2(q^2) = \pi^2/15 - \ell^2$: apply (10) at q^2, and use $\mathrm{Li}_2(q) + \mathrm{Li}_2(-q) = \frac{1}{2}\mathrm{Li}_2(q^2)$. Thus the exact finite-interval forms are

$$\int_0^\ell t^2 \coth t\, dt = \frac{\zeta(3)}{10}, \qquad \int_0^\ell t \coth t\, dt = \frac{\pi^2}{20}. \tag{11}$$

To verify, expand $\coth t = 1 + 2\sum_{n\geq 1} e^{-2nt}$ and use positivity to integrate termwise. The first integral becomes $\ell^3/3 + \zeta(3)/2 - \mathrm{Li}_3(q^2)/2 - \ell\,\mathrm{Li}_2(q^2) - \ell^2\,\mathrm{Li}_1(q^2)$. Since $\mathrm{Li}_1(q^2) = \ell$, (9) cancels the logarithmic terms and leaves $\zeta(3)/10$. The second is $\ell^2/2 + \zeta(2)/2 - \mathrm{Li}_2(q^2)/2 - \ell\,\mathrm{Li}_1(q^2) = \pi^2/20$. These classical Hjortnaes identities are also treated by Markov (2024).

Equation (9) rearranges exactly to

$$\gamma_* = \frac{\frac{5}{4}\mathrm{Li}_3(\varphi^{-2}) + \frac{\pi^2}{6}\ell - \frac{5}{6}\ell^3}{\pi e \varphi}. \tag{12}$$

This is an exact evaluation route, not an independent selection of the denominator. For $z = \varphi^{-2}$, truncating the trilogarithm after K terms gives a positive error bounded by $5z^{K+1}/[4\pi e\varphi(K+1)^3(1-z)]$, by bounding every omitted denominator below by $(K+1)^3$ and summing the geometric tail.

5. The three loop observables and the two signs of 1/12

These calculations independently rederive the accepted October investigation. Their operator scale, zero-mode removal, exponent and bulk subtraction are **A**, not recovered July choices.

5.1 Spectrum and exact finite inverse trace

For $N \geq 3$ define $(\Delta_N u)_j = u_{j+1} + u_{j-1} - 2u_j$ with periodic indices and $A_N = N^2(-\Delta_N)$. Substituting $u_j = e^{2\pi i k j/N}$ gives

$$\lambda_{N,k} = 4N^2 \sin^2(\pi k/N), \quad k = 0, \ldots, N-1. \quad Z_N(p) = \sum_{k=1}^{N-1} \lambda_{N,k}^{-p}. \tag{13}$$

The constant mode $k = 0$ is deliberately excluded. The factor N^2 chooses spacing $1/N$, i.e. unit circumference. For circumference $\mathcal{L}$, eigenvalues are divided by $\mathcal{L}^2$ and $Z_N(p)$ is multiplied by $\mathcal{L}^{2p}$.

Factorization of $z^N - 1$ into its N roots, with $z = e^{2ix}$, gives a nonzero constant times $\prod_{k=0}^{N-1} \sin(x + k\pi/N) = c_N \sin Nx$. The constant is immaterial to the logarithmic derivative. Differentiating once and then once more gives

$$\sum_{k=0}^{N-1} \cot(x + k\pi/N) = N \cot Nx, \qquad \sum_{k=0}^{N-1} \csc^2(x + k\pi/N) = N^2 \csc^2 Nx. \tag{14}$$

Remove the $k = 0$ term and let $x \to 0$. The Taylor expansion $\csc^2 x = x^{-2} + 1/3 + O(x^2)$ cancels the divergent terms, leaving $\sum_{k=1}^{N-1} \csc^2(\pi k/N) = (N^2 - 1)/3$. Therefore

$$Z_N(1) = \frac{1 - N^{-2}}{12}, \qquad \lim Z_N(1) = \frac{1}{12}. \tag{15}$$

For example $Z_3(1) = 2/27$, not $1/12$. Equality of the limiting number with (4) does not identify a three-node graph with a filled triangle.

5.2 The inverse-cube frequency sum

For $1 \leq k \leq N/2$, concavity gives $\sin(\pi k/N) \geq 2k/N$, so $\lambda_{N,k} \geq 16k^2$. Pair k and $N - k$. For fixed k, $\lambda_{N,k} \to (2\pi k)^2$, while the paired inverse powers are bounded by $2(16k^2)^{-p}$. This is summable for $p > 1/2$. Dominated convergence, extending absent terms by zero, proves

$$\lim_{N\to\infty} Z_N(p) = \frac{2\zeta(2p)}{(2\pi)^{2p}}, \qquad \lim Z_N(3/2) = \frac{\zeta(3)}{4\pi^3}. \tag{16}$$

The possible unpaired midpoint for even N tends to zero and does not alter the limit. Taking N through multiples of three gives the same full-spectrum result.

By contrast, restricting to states repeated every three nodes keeps only $k = 0, N/3, 2N/3$. The two nonzero eigenvalues are $3N^2$, hence the restricted sum is $2(3N^2)^{-p} \to 0$. With the unscaled operator $-\Delta_N$ they instead stay equal to 3. Neither case equals the full refinement in (16).

The historically normalized October diagnostic is

$$\gamma_N = \frac{4\pi^2}{e\varphi} Z_N(3/2) \longrightarrow \gamma_*. \tag{17}$$

The spectral exponent and prefactor are explicit analytical choices. No preceding geometric identity uniquely selects them. For example choosing $p = 1$ produces (15) instead, and changing circumference changes the uncorrected trace.

5.3 A different observable produces a negative finite part

Define the half-total normalized frequency sum

$$S_N = \frac{1}{4\pi} \sum_{k=1}^{N-1} \sqrt{\lambda_{N,k}} = \frac{N}{2\pi} \sum_{k=1}^{N-1} \sin(\pi k/N). \tag{18}$$

The finite geometric sum of $e^{ik\theta}$ has imaginary part

$$\sum_{k=1}^{N-1} \sin(k\theta) = \sin((N-1)\theta/2) \sin(N\theta/2) / \sin(\theta/2).$$

At $\theta = \pi/N$ this is $\cot(\pi/(2N))$. Replacing the same finite dispersion by its continuous integral gives

$$S_N = \frac{N}{2\pi} \cot \frac{\pi}{2N}, \quad S_N^{\text{bulk}} = \frac{N}{2\pi} \int_0^N \sin(\pi x/N)\, dx = \frac{N^2}{\pi^2}. \tag{19}$$

Divide the Taylor series for cosine by that for sine to obtain $\cot x = x^{-1} - x/3 - x^3/45 - 2x^5/945 + O(x^7)$. Then

$$S_N - S_N^{\text{bulk}} = -\frac{1}{12} - \frac{\pi^2}{720N^2} - \frac{\pi^4}{30240N^4} + O(N^{-6}). \tag{20}$$

The sign has a finite geometric explanation: the trapezoidal sum for the strictly concave sine arch lies below its integral; both endpoints are zero. All terms in S_N remain positive. The negative quantity is the **difference** in (20).

This equals the usual analytically continued value $\zeta(-1)$, but the ordinary series $\sum n$ diverges. DLMF section 25.6 records the continued integer values. As another direct comparison,

$$\sum_{n \geq 1} n e^{-tn} = \frac{e^{-t}}{(1 - e^{-t})^2} = t^{-2} - \frac{1}{12} + O(t^2). \tag{21}$$

Changing to half-integer modes gives $-\frac{d}{dt}[1/(2\sinh(t/2))] = t^{-2} + 1/24 + O(t^2)$. Thus the mode family and subtraction matter. A further oscillator factor $1/2$ halves a coefficient; it does not change the inverse moment in (16) into an energy sum.

Equations (4), (15), (16) and (20) are compatible identities. None says that γ_* replaces $-1/12$ or establishes a physical string model.

6. The lock equation: what was printed, assigned, or measured

6.1 July sources contain a real discrepancy

P06 physical p.4 Eq.(2) prints (1). Its p.6 glossary identifies γ as **Euler–Mascheroni's constant**, approximately 0.577. Those two statements cannot both be exact: substituting $\gamma_{\mathrm{EM}} = 0.5772156649\ldots$ does not satisfy (1). This report uses the new disambiguating notation γ_* for the solved target and γ_{EM} for the classical constant; it does not repair the manuscript.

P01 pp.3–4 Eq.(1) repeats the equation, calls γ a scaling constant of order 10^{-1}, describes calibration, and explicitly says the relation was not derived from first principles. Its claim of observation is historical evidence **of the claim**, not a surviving estimator or measurement receipt. P05 p.2 and the position reconstruction instead assign GAMMA = 0.577; it is unused in that engine.

6.1a The author's design interpretation

The fusion manuscript's Appendix A, physical pp.19–20, supplies a positive historical explanation. It associates the circle constant and the exponential base with geometry and growth, the golden ratio with packing or growth patterns, the zeta value with discrete lattice sums, and gamma with the scale linking those ingredients. It reports approximately 0.087 as a calibration value. These are the author's documented interpretations; the appendix itself distinguishes empirical or analytical tools from a first-principles physical derivation.

That makes the formula intelligible as a design relation. It does not make each factor uniquely forced by one diagram, or supply an independent gamma estimator. The exact arithmetic in (1), the historical interpretation, and the separate later spectral normalization (17) therefore have different evidential roles. The original appendix also includes physical speculation; this companion does not adopt it as a proved result.

6.2 September's proposed estimator is not an implementation

P19 p.8 describes γ as inferred from spectra and adjusted to satisfy the lock. Page 13 calls $\widehat{\gamma}^{(t)} \leftarrow \mathrm{GammaEstimate}(\{\Phi_a\}, \mathrm{spectra}, \zeta(3))$, without defining the function. The p.12 value 0.10 occurs in a table explicitly labelled **"Run Stamp (example)"**, with a placeholder commit and incomplete parameter arrays. It is not an independently verified run measurement.

An independently testable estimator would need an explicit map from state/spectral data to a number, its units/normalization and the raw inputs. A lock penalty or tuning procedure that drives a residual toward zero is not, by itself, an independent measurement of the target. Those missing specifications remain **U**.

6.3 March saved outputs distinguish predictions from observations

The inert source srg_resonance_lock_analysis.py, line 42, computes gamma_derived as zeta3/lock_product. Lines 443–448 label the odd-zeta extension as a hypothesis. Its saved resonance_lock_results.json contains $\gamma_{\mathrm{SRG}} = 0.08699474751703501$, the formula itself, and six non-finite eigenmode lock times. This is evidence of a saved formula evaluation and unsuccessful listed searches, not six coupling measurements.

Tower II lines 289–318 explicitly compute the zeta argument from the **assigned** frequency, put the result in **gamma_predicted**, and separately collect lock_time, final_R and final_coherence. The received CSV has 90 rows: 45 formula predictions and 45 blanks. All finite predictions agree with the indicated formula to the precision of the saved numbers; every blank is at a negative

tower index. These counts come from the accepted lineage investigation's static audit of existing records, which is reused here without replay. No trajectory was regenerated.

Infinity is different: its assigned 0.08699 actually multiplies the R/D cross term (section 8). Conversely Seed Explorer's 0.0870 is stored but unused. One cannot infer use merely from a parameter name.

6.4 The other "constant one" candidates

The informal phrase does not uniquely name a recovered equation. The prior bounded search found no literal occurrence in 278 supplied Markdown/text files; that private file inventory is not republished here. It would be unjustified to identify it with the July equation merely because both contain constants. Several distinguishable candidates survive:

July lock. *Exact source and meaning:* P06 p.4 Eq.(2), P01 p.3 Eq.(1). *Finding:* $\pi e\varphi\gamma = \zeta(3)$; explicit historical target, conflicting γ definitions.

Dual-phase master operator. *Exact source and meaning:* P13 p.5. *Finding:* A tensor/phase/floor/compression expression, not the scalar lock.

A literal equation equal to one. *Exact source and meaning:* srg_resonance_lock_analysis.py, lines 293–327. *Finding:* $1 = (1 + \lambda H/H_{\max})(1 + h\mu + \kappa/s)$, an approximate single-mode multiplier condition.

Frequency master equation. *Exact source and meaning:* P18 p.7 Eq.(12). *Finding:* $\pi e\varphi\,\gamma(\omega) = \zeta(2\log_\varphi(\omega/\omega_0) + 3)$.

DieGlocke lock ratio. *Exact source and meaning:* DieGlocke_Sim_Core.py, lines 6–14. *Finding:* $\pi e\varphi\,\gamma_{\mathrm{EM}}\zeta(3)$, different algebra from the July lock.

The multiplier equation follows formally by linearizing $\kappa\tanh(\Theta/s) \approx (\kappa/s)\Theta$ on an eigenmode $\mathcal{L}\Theta = \mu\Theta$, **and** treating compression as identity on the chosen branch. For the actual $\varepsilon > 0$ RPCO, the inactive nonzero branch is not exactly identity. The approximation is not a proof of (1), contains no $\zeta(3)$, and does not supply an estimator. The source's $(1, 1)$ "fundamental" omits lower nonconstant modes $(1, 0)$ and $(0, 1)$ of the two-dimensional periodic grid. Its assigned polyhedral names are not spectral identifications proved there.

DieGlocke uses rhs = 1/(GAMMA ZETA3), then returns lhs/rhs. Setting that ratio to one would require $\gamma = 1/(\pi e\varphi\zeta(3))$, not (1). Moreover GAMMA is actually assigned 0.5772156649, and the provided Run file neither imports nor calls the ratio. Its time loop freshly samples a product of prescribed fields; earlier composite states are stored for diagnostics, not fed into the next composite (Run lines 37–46). The radial phases, 0.244, 0.122, and modulation gamma 0.3 are explicit inputs, not recovered lock measurements.

P13's master operator includes $\nabla\nabla\cdot\Phi - \Delta\Phi$, floor quantization and feedback, with complex phase $\Phi = e^{i\theta}T$. Its floor/domain and tensor-feedback typing are unresolved, as already recorded in SRG-I section 6.4 (physical pp.12–13). It is a distinct historical proposal, not a substitute solver selected here.

Bounded provenance conclusion. No implemented independent GammaEstimate or saved independent gamma measurement was recovered in the searched sets. No native rule selecting the inverse exponent 3/2 and normalization $4\pi^2/(e\varphi)$ was found there. The latter first appears in this inspected chain as an explicitly declared October diagnostic choice. The informal phrase "the weird constant one" was part of the provenance request, not a manuscript title. Its exact referent remains unidentified; the competing equations above are now documented without asking the owner to reconstruct them from memory.

7. TGMO, REFU and RPCO: precise operator lineage

7.1 Earlier names did not denote one unchanged function

P08 p.1 gives the golden stability map $(\varphi/\sqrt{3})\tanh x$. Its derivative has absolute value at most $\varphi/\sqrt{3} < 1$, and its output is bounded by that same constant. This is a valid scalar damping statement. It is not the later threshold compressor or either logarithmic RPCO.

P07 pp.3–6 uses "Toroidal Gradient Memory Operator" and "Recursive Entangled Field Update." It prints a weighted spatial Laplacian and the positive-output logarithmic compressor. Its claim that that compressor retains negative signs contradicts its exponential formula.

P05 p.1 and the surviving standalone scripts instead implement:

- TGMO.py lines 5–14: $(x, y, z) = ((2+r\cos\theta)\cos(\pi/4), (2+r\cos\theta)\sin(\pi/4), r\sin\theta)$. This is a sampled minor circle at fixed major angle, repeated through the chosen number of rotations; it is not a field Laplacian.
- REFU.py lines 5–13: the prescribed scalar curve $H(t)$ below, not a full next-state update.
- RPCO.py lines 5–13: $\sin(5\,\mathrm{atan2}(Y, X))e^{-0.05(X^2+Y^2)}$, a sampled spatial phase field, not a compressor applied to evolving amplitudes.

P11 pp.1–2 correctly distinguishes a constant radius from a field. $\Delta r = 0$ for constant r, whereas $\Delta\sqrt{x^2+y^2} = 1/\sqrt{x^2+y^2}$ away from the origin in two dimensions. The latter is singular at the origin. Promoting a constant radius to a spatial field changes the mathematical object; the alternatives are not interchangeable repairs.

P03 pp.7–8 is also not January's spatial stencil: its trail definition is $\nabla_T^2\Theta(t) = \Theta(t+1) - 2\Theta(t) + \Theta(t-1)$. Substitution in the printed update gives $y = C_\lambda(y+b)$, $b = \Theta(t-1) - \Theta(t) + F(t)$, for a finite scalar state and frozen feedback. Its compressor is $C_\lambda(x) = x$ below λ and x/φ at or above λ.

The complete branches, reusing accepted SRG-I section 3.2, Theorem 3.1 (physical pp.4–6), are

$$\{y : y = C_\lambda(y+b)\} = \begin{cases}(-\infty, \lambda), & b = 0\\ \varnothing, & b \neq 0\end{cases} \cup \begin{cases}\{\varphi b\}, & \varphi^2 b \geq \lambda\\ \varnothing, & \varphi^2 b < \lambda.\end{cases} \tag{22}$$

Proof: on the first branch $y = y + b$, requiring $b = 0$; on the second $(1-\varphi^{-1})y = \varphi^{-1}b$, giving $y = \varphi b$ and input $\varphi^2 b$. This exhibits both no-solution and nonunique cases. No intended root-selection solver is supplied by these sources. Replacing the temporal term by a spatial difference would be a new model choice.

The July position reconstruction, P05 p.2 / recursive_field_evolve.py, uses a fixed directed five-neighbour graph from the starting positions: $Y = X + 0.05\Delta_{\text{graph}}X + 0.15\tanh(\Delta_{\text{graph}}X)$, then $Z = 0.95Y + 0.05\overline{Y} + \xi$ and a small prescribed sinusoidal scaling of Z's third coordinate. It is a concrete alternative evolution, with stochastic position noise, not an implementation of (22).

7.2 January's complete fixed-grid composition [H]

For a finite real two-dimensional field, JANUARY pp.2–3 and tgmo.py lines 4–46 define

$$\mathcal{L}\Theta_{ij} = a^{-2}(\Theta_{i+1,j} + \Theta_{i-1,j} - 2\Theta_{ij}) + b^{-2}(\Theta_{i,j+1} + \Theta_{i,j-1} - 2\Theta_{ij}). \tag{23}$$

Indices wrap on both axes. The optional third axis has a separate second difference, reflecting by default. The normalized loop (13) shares the one-axis stencil but adds a positive sign reversal, N^2 scaling, zero-mode removal and a spectral observable. January does not compute (17).

An embedded torus needs an additional geometric convention. In January's helper, θ is the major-circle angle and ϕ the minor-circle angle, with $a > b > 0$:

$$X(\theta, \phi) = ((a + b\cos\phi)\cos\theta, (a + b\cos\phi)\sin\theta, b\sin\phi).$$

Taking the squared lengths and dot product of its coordinate derivatives gives the metric $\mathrm{diag}((a + b\cos\phi)^2, b^2)$. The metric Laplacian, defined by $|g|^{-1/2}\partial_i(|g|^{1/2}g^{ij}\partial_j f)$, is consequently

$$\Delta_g f = \frac{f_{\theta\theta}}{(a + b\cos\phi)^2} + \frac{f_{\phi\phi}}{b^2} - \frac{\sin\phi}{b(a + b\cos\phi)} f_\phi.$$

The accepted lineage report uses the consistently exchanged angle names for this comparison; the present edition follows the January helper (tgmo.py lines 62–76). Neither convention makes its constant-coefficient stencil (23) the induced metric operator. SRG-I section 5.3, physical p.9, Eq.(5.7), gives the related comparison. "Toroidal Grid Modulation Operator" names the actual January periodic stencil.

For (23)–(24), positive grid scales a, b, a nonzero finite s, and finite real coefficients and inputs avoid undefined divisions; $H_{\max}$ is fixed, positive and finite. In the app's reference callbacks $D(\Theta, t) = \Theta$ and $f(D) = \kappa\tanh(D/s)$. With a fixed finite positive $H_{\max}$,

$$\begin{aligned} P(\Theta) &= \Theta + h\mathcal{L}\Theta + \kappa\tanh(\Theta/s), \\ Q_t(\Theta) &= m_t P(\Theta), \quad m_t = 1 + \lambda H(t)/H_{\max}, \\ \Theta^+ &= C_{\mathrm{mode}}(Q_t(\Theta)), \\ H(t) &= 2e^{-0.5t}\cos t + e^{-0.3t}\sin t + 1.5e^{-0.7t} + e^{-0.4t}. \end{aligned} \tag{24}$$

This order is explicit in refu.py lines 109–138. Both spatial and nonlinear terms read the same incoming Θ; H multiplies before RPCO. Only the spatial term is multiplied by h. Calling the scheme "Euler style" does not establish a continuous-time limit with every other per-step operation held fixed.

Here H is prescribed by the clock, not an accumulator of past states. Any state-driven memory claim needs an additional defined carrier and update. The library samples $\max|H|$ and adds a guard (lines 25–31). The rollout starts at $t = 0$. A custom H requires both its matching Hmax and compute_global_Hmax=False; automatic normalization otherwise evaluates the original H and overwrites the supplied value. Accepted OC01's correction is retained.

7.3 The two compressors and their symmetry [D from H]

Let $u = e^{-L}$, $v = e^{L}$, $L > 0, \varepsilon > 0$. Moving the exponential through the monotone clamp gives exactly

$$C_p(x) = \begin{cases} \mathrm{clip}(x + \varepsilon, u, v), & x > 0, \\ 1, & x = 0, \\ \mathrm{clip}((|x| + \varepsilon)^{-1}, u, v), & x < 0, \end{cases} \tag{25}$$

$$C_s(x) = \begin{cases} \mathrm{sgn}(x)\,\mathrm{clip}(|x| + \varepsilon, u, v), & x \neq 0, \\ 0, & x = 0. \end{cases}$$

These are refu.py lines 37–69 / JANUARY p.3. They imply the exact reciprocal pairing $C_p(a)C_p(-a) = 1$ for $a > 0$, since clipping in logarithmic space commutes with sign reversal. Signed mode instead has $C_s(-a) = -C_s(a)$. Neither identity is time reversal or an inverse next-state map; a plateau loses magnitude information.

An inactive signed magnitude grows by ε. P18 p.4 Eq.(2) prints an outside-sign formula, but its prose claims $-a \mapsto 1/a$. Without epsilon or clipping, that displayed formula returns $-a$. January's paper mode has the reciprocal negative branch; the March engines implement signed mode. Substituting one for the other would alter the historical recurrence.

The OC01 corrections remain: signed mode is discontinuous at zero for every $L > 0, \varepsilon > 0$; paper mode is continuous there exactly when $\varepsilon = 1$, and then is C^1 but not C^2 locally. Indeed it is $1 + x$ on the positive side and $(1 - x)^{-1}$ on the negative side. Multiplication by a zero gain makes the complete map constant, so discontinuity claims about the compressor do not automatically transfer to that case. Diagnostic samples and update counts remain distinct.

7.4 A concrete 3×3 mathematical step

Choose a periodic grid, $a = b = 1, h = 0.1, \kappa = 0.1, s = 1$, H off, $L = 9, \varepsilon = 10^{-9}$. These are declared example parameters, not all app defaults. With

$$\Theta = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \mathcal{L}\Theta = \begin{pmatrix} 0 & 1 & 0 \\ 1 & -4 & 1 \\ 0 & 1 & 0 \end{pmatrix},$$

the nonlinear term is nonzero only at the centre, because it reads Θ, not an already diffused state. Thus

$$P(\Theta) = \begin{pmatrix} 0 & 0.1 & 0 \\ 0.1 & c & 0.1 \\ 0 & 0.1 & 0 \end{pmatrix}, \quad c = 0.6 + 0.1 \tanh 1 = 0.6761594155955765\ldots. \tag{26}$$

The centre and four edge entries lie on inactive positive branches. Paper RPCO returns $\begin{pmatrix} 1 & 0.1+\varepsilon & 1 \\ 0.1+\varepsilon & c+\varepsilon & 0.1+\varepsilon \\ 1 & 0.1+\varepsilon & 1 \end{pmatrix}$; signed RPCO returns the same positive entries but keeps all four corners **zero**. This is a full algebraic evaluation of the stated example; no field-step function, field rollout or random initializer was executed.

Accepted OC01's paper-cap result and OC02's signed floor patterns/capture sets are later deductions from (24)–(25), with their stated parameter assumptions. They are not claims found in July manuscripts and do not transfer automatically to March's particle state.

8. The March adaptations

8.1 Documentary connection and substantive changes

Seed Explorer line 29 explicitly describes the operators as "from your refu.py and tgmo.py, adapted for glyph particles." Lines 53–57 distinguish its position/resonance particle state from a fixed grid. This is direct documentary lineage; it does not assert identical operators.

For a directed neighbour set N_i, the position difference is $\Delta_i X = |N_i|^{-1} \sum_{j \in N_i} (X_j - X_i)$. The normalized resonance weights are proportional to $(d_{ij} + 10^{-8})^{-1}$. Seed Explorer lines 91–169 rebuild the Euclidean graph, update $X_i \leftarrow X_i + 0.05\Delta_i X + 0.15 \tanh(\Delta_i X)$, mix R_i toward its weighted neighbour mean, multiply by H, move positions toward a resonance-weighted centre, apply a prescribed 0.244 sinusoid, then apply signed RPCO. Position noise follows. Its pass count increments before the first update, so its first H argument is one, unlike January's zero. There is no toroidal wrapping of these Cartesian positions.

July reconstructed engine. *Precisely supported change:* Position-only state, graph fixed before loop, unweighted centre, third-coordinate sinusoid; assigned GAMMA unused. *Primary locator:* P05 p.2; early/recursive_field_evolve.py.

Seed Explorer. *Precisely supported change:* Rebuilt kNN graph, separate R, weighted resonance mixing and fusion, H and signed amplitude RPCO. *Primary locator:* lines 29,53–79,91–169.

Infinity. *Precisely supported change:* Adds real D field, unequal R/D diffusion strengths, actual assigned-gamma cross coupling, shifted H for D. *Primary locator:* lines 91,114–137,162–195.

Condensate Breaking. *Precisely supported change:* Separate branch with optional horizon, multifrequency and breathing compression; gamma stored. *Primary locator:* lines 61–99,155–196.

Golden Tower. *Precisely supported change:* Fixed assigned golden-spaced bands; log-frequency matching, sinusoid per band, additive golden-ratio resonance mixing, band-local fusion. *Primary locator:* lines 72–94,122–179.

Tower II. *Precisely supported change:* Adds configurable stronger mixing over $\varphi^{\pm 1}, \varphi^{\pm 2}$ and sum/difference match tests; optional amplitude-dependent cap; same stage ordering. *Primary locator:* lines 135–213.

These are selected source variants, not a claim that every archived program belongs to one linear sequence.

8.2 R/D cross coupling uses the assigned gamma

For a row with neighbours, Infinity's default coefficients give

$$\begin{aligned} R_i' &= R_i + 0.15(\overline{R}_i - R_i) + 0.01 c_p D_i, \\ D_i' &= D_i + 0.075(\overline{D}_i - D_i) + 0.01 c_p R_i, \\ c_p &= 0.08699 \sin(0.244p + \pi/2). \end{aligned} \tag{27}$$

Both right sides read incoming R/D, and rows without neighbours skip this coupling. Subsequent gains use $H(p)$ and $H(p + \pi/2)$ separately. A time shift of a decaying curve changes its envelope as well as its trigonometric phase. These real fields are not defined as complex conjugates. Equal cross entries alone do not establish sector-exchange symmetry: the diffusion coefficients and later gains differ.

8.3 Tower II's actual composition and static master equation

Tower II lines 140–213 fix the graph and distances for that pass and move positions. At the resonance stage, a row with no selected neighbours, or with unnormalized frequency/distance weight sum below 10^{-10}, is skipped: $R_i^{\text{mix}} = R_i$. Its sine and additive mixing are both skipped. This skips only that row's resonance-mixing stage; the subsequent H gain, fusion, RPCO and position-noise stages still follow. For an active row that passes both guards, the source constructs

$$R_i^{\text{mix}} = a_i\big[(1-\beta)R_i + \beta\sum_j w_{ij}R_j\big] + \sum_j \eta_{ij}R_j, \quad a_i = 1 + 0.01\sin(2\pi\omega_i p). \tag{28}$$

For active rows, η_{ij} is the sum of the nonnegative golden-ratio and sum/difference match additions in lines 169–188. Those additions occur **after** the per-site sine multiplication; they are not multiplied by a_i there. H then multiplies R^{mix}, that precompression resonance controls band-local position fusion, and signed RPCO is last for R. Position noise is separate. This recovers the stage distinction used by accepted R0–R3 without rerunning them.

P18 p.5 Eq.(6) defines $\gamma_m = \zeta(2m+1)/(\pi e\varphi)$ for $m \geq 1$. Page 7 assigns $\omega_n = \omega_0\varphi^n$, $\omega_0 = 0.244$, then states

$$\pi e\varphi\,\gamma(\omega) = \zeta\Big(2\log_\varphi(\omega/\omega_0) + 3\Big). \tag{29}$$

At ω_n this samples $\zeta(2n+3)$, so matching the older odd-zeta indexing requires $m = n+1$, with $n \geq 0$. This is an index substitution, not a temporal recursion. The positive Dirichlet-series domain requires $\omega > \omega_0/\varphi$. At $\omega = \omega_0/\varphi$ the zeta argument is one and has a pole. At lower frequencies analytic continuation is a further prescription; it is not a positive coupling at every band. For instance $n = -2$ gives $-1/(12\pi e\varphi)$ if analytically continued.

The source's diagnostic explicitly requires zeta_arg > 1.01 and otherwise writes NaN (Tower II lines 293–298). No zeta evaluation occurs in the recovered update (28). The constant-frequency labels are assigned and are not generated by the mix. P18 itself calls the master equation static on p.7. Its limit $\gamma_m \to 1/(\pi e\varphi)$ follows from $\zeta(s) \to 1$ as $s \to +\infty$; neither that limit nor the CSV proves measured equilibrium couplings equal predictions.

9. Claim disposition and remaining definition

The source-grounded chain is:

1. **Recoverable mathematics:** circle projections; exact equilateral ratio; Fibonacci limit; exponential definition; iterated-integral identity; golden polylogarithm identity; cycle spectrum and all three observables.
2. **Recovered historical choices:** July lock formula, competing gamma meanings, threshold compression, prescribed H, January stencil/order/modes, March assigned frequencies and coupling coefficients.
3. **Later analytical choices:** scalar branch analysis with frozen feedback; normalized full-loop refinement and spectral powers; the prefactor importing the lock target; cap/floor and capture theorems.
4. **Incompatible descriptions:** P06's Euler-gamma glossary versus lock; P07's sign-retention claim versus positive exponential; P18's outside-sign equation versus reciprocal prose; Rainman's unchanged scaled integral and forward-image sign.
5. **Underdetermined:** original selector for $\pi e \varphi$; the missing GammaEstimate; an independently measured coupling record; the vesica-cycle-to-integration-chain map; temporal root selection; the exact informal "constant one" referent.

The first three categories support a careful mathematical reconstruction manuscript when attributed and qualified as above. Historical empirical/physical conclusions require evidence beyond these identities. The report proposes no replacement recursion or source repair. The most precise remaining mathematical question is whether an existing native state statistic and an independently specified selection rule choose (17)'s observable and scale. The bounded search found no such specification.

10. Reading the evidence responsibly

An equation can play several roles. It may define a chosen operation, express a theorem once its assumptions are fixed, describe a measured relationship, or propose a physical interpretation. The histories in this paper contain all four roles. The worked examples and proofs establish the stated algebra and limits; historical empirical descriptions remain attributed to their sources. No experiment, historical simulation or earlier verification suite was rerun for this edition.

The October lineage investigation received an internal GPT review, accepted with the explicit source-and-notation addendum incorporated here. This companion is a new presentation submitted for final review. Neither label means external peer review, journal acceptance or physical validation. Accepted SRG-I, source code and earlier research remain unchanged.

The proof chain is self-contained for the central identities: circle intersection, equilateral inradius, Fibonacci ratio, compounding limit, iterated integration, the cycle's finite inverse trace and continuum inverse powers, its negative finite part, the golden polylogarithm and finite-interval identities, and the source-specific compression/implicit branches. Standard trigonometric and polylogarithm definitions are linked in the source index.

The main unresolved question is precise: which independently specified native statistic and selection rule, if any, choose the historical coupling scale? A geometric interpretation of the factors exists. The bounded archive search did not recover the additional estimator needed to test that selection claim independently. Neither finding erases the other.

Assistance and attribution

Hilmir Frímann Halldórsson is the named human author. ChatGPT/GPT and Codex assisted with source recovery, mathematical checking, editorial organization and document preparation. Historical AI credits are preserved as source history, not adopted as institutional affiliation or endorsement. The classical identities are attributed to established mathematics; the present work claims a source-grounded exposition rather than their discovery.

Reuse

This edition adds no blanket reuse licence to the derived manuscript or its records. Linked historical works retain their source-specific notices, including the identified CC BY 4.0 deposits. Public visibility, internal acceptance and reuse permission are separate matters. The accompanying publication disposition records the proposed file set and review status.

11. Source keys and reading map

These keys are used throughout the paper. Page numbers are physical PDF pages. Exact file hashes and code access instructions are in the accompanying SOURCE_INDEX.md; full historical files are not republished.

P01 — Symbolic Recursion and SRG Fusion: Emergent Lattice Fields from Glyph Memory Dynamics

Hilmir Frímann Halldórsson; source AI credits retained in the original. 15 July 2025. Cited locations: pp.3–4, Eq.(1); Appendix A, pp.19–20. Read this source.

P03 — Symbolic Recursion and SRG Formation: Full Method and Emergence

Hilmir Frímann Halldórsson; source AI credits retained in the original. 14 July 2025. Cited locations: pp.2,5,7–8. Read this source.

P05 — Patch Research: Detailed Reconstruction of Missing Components in Symbolic Recursion and SRG Framework

Collaborative Reconstruction. 15 July 2025. Cited locations: pp.1–2. Read this source.

P06 — Recursive Dark Energy and Vacuum Feedback: A Vesica-Phase Cosmological Framework

Hilmir Frímann Halldórsson; source AI credits retained in the original. 13 July 2025. Cited locations: p.4 Eq.(2); p.6 glossary. Read this source.

P07 — Recursive Numerical Operators for Symbolic Dynamics

Hilmir Frímann Halldórsson; source AI credits retained in the original. 11 July 2025. Cited locations: pp.3–6. Read this source.

P08 — Recursive Symbolic Evolution and Feedback Dynamics in Python (three-page version)

Hilmir Frímann Halldórsson; source AI credits retained in the original. 29 June 2025. Cited locations: p.1 section 2.1. Read this source.

P09 — Recursive Symbolic Evolution and Feedback Dynamics in Python (seven-page version)

Hilmir Frímann Halldórsson; source AI credits retained in the original. 29 June 2025. Cited locations: pp.1–7, comparison. Read this source.

P11 — Why the Toroidal Laplacian Failed: A Structural Explanation

Author not printed on the inspected opening page; surviving project manuscript. Undated. Cited locations: pp.1–2. Read this source.

P13 — Unified Recursive Cosmological Model Memo

Author not printed on the inspected opening page; surviving project manuscript. Undated cover. Cited locations: p.5 master operator. Read this source.

P18 — Symbolic Resonant Geometry: Recursive Field Dynamics, the Golden Frequency Tower, and the Breathing Compression Field

pzychozen; source also credits Claude. 14 March 2026. Cited locations: p.4 Eq.(2); p.5 Eq.(6); p.7 Eqs.(10),(12). Read this source.

P19 — Unified Recursive Fields: Geometry, Recursion, and Consciousness

Hilmir Frímann Halldórsson; source AI credits retained in the original. 6 September 2025. Cited locations: pp.8,12–13. Read this source.

JANUARY — An Experimental Framework for Exploring Recursive Numerical Operators: TGMO, RPCO, and REFU as a Reproducible Computational Laboratory

Hilmir Frímann Halldórsson; source AI credits retained in the original. 25 January 2026. Cited locations: pp.2–3; exact companion source versions below. Read this source.

RAINMAN — Rainman

Hilmir Frímann Halldórsson; source AI credits retained in the original. 22 July 2025. Cited locations: pp.3–4,18,20–24. Author-supplied copy; no verified public URL. Relevant equations and counterexamples are stated in this paper.

Program families and accepted analyses

July reconstructed operators: TGMO.py samples a minor circle, REFU.py supplies H, RPCO.py samples a damped phase field, and recursive_field_evolve.py evolves positions. The exact standalone files are also linked individually in SOURCE_INDEX.md.

January fixed-grid sources: tgmo.py, refu.py and app.py, exact versions identified in SOURCE_INDEX.md. All January code locators refer to this version, not a later app revision.

March source and saved-output archive: simulations.zip, with Seed Explorer, Infinity, Condensate Breaking, Golden Tower, Tower II, resonance-lock analysis, its saved JSON and master-equation CSV. Cited line numbers use the exact members hashed in SOURCE_INDEX.md. No program was run for this edition.

DieGlocke_Sim_Core.py and DieGlocke_Run.py are separate author-supplied files with no verified public URL. SOURCE_INDEX.md records their exact identities; the relevant ratio and data-flow distinction are explicit in section 6.4.

SRG-I: Foundational Reconstruction, v0.1.1: internally accepted companion on the original implicit recursion and its unresolved definitions. Use section 3.2 (pp.4–6), section 5.3 (p.9) and section 6.4 (pp.12–13) for this paper's three precise cross-references.

F01 = Foundations-01; OC01 = Operator Composition 01; OC02 = Signed Plateaus and Capture Sets; R0–R3 = the closed Tower II audit. These internal records and the October Geometry–Zeta

note are background analyses identified by hash in SOURCE_INDEX.md. The current paper reproduces its central proofs instead of requiring access to those private records.

Classical references

NIST DLMF section 25.12, Polylogarithms; section 25.6, Integer Arguments of the Riemann Zeta Function.

Lubomir Markov, Hjortnaes Integrals Involving the Di- and Trilogarithms, International Journal of Applied Mathematics 37(3), 305–310 (2024), DOI 10.12732/ijam.v37i3.2; especially pp.305–308, Eqs.(2),(3),(8),(11).