Insert official SU logo


Salahaddin University-Erbil

School of Science | Department of Information Technology

First-year mathematics

COMPLEX NUMBERS

Properties of Complex Conjugate and Absolute Value



Prepared by

[STUDENT 1] [STUDENT 2] [STUDENT 3]

[STUDENT 4] [STUDENT 5] [STUDENT 6]

[STUDENT 7] [STUDENT 8] [STUDENT 9]

Supervised by [LECTURER NAME] **Academic year** [ACADEMIC YEAR]

Outline

1 **Introduction** — Slide 3
Why we need complex numbers — [STUDENT 1]

2 **The basics** — Slides 4–6
Complex numbers, real and imaginary parts, the complex plane — [STUDENT 2], [STUDENT 3]

3 **The complex conjugate** — Slides 7–10
Definition, picture on the graph, rules and examples — [STUDENT 3], [STUDENT 4], [STUDENT 5]

4 **Absolute value** — Slides 11–14
Meaning, picture on the graph, worked examples, properties — [STUDENT 5], [STUDENT 6], [STUDENT 7]

5 **Connecting the two ideas** — Slides 15–16
The identity $z\bar{z} = |z|^2$ and a side-by-side comparison — [STUDENT 7], [STUDENT 8]

6 **Applications, quiz and conclusion** — Slides 17–19
Where complex numbers are used, practice questions, summary — [STUDENT 8], [STUDENT 9]

Why Complex Numbers?

? Can we find the square root of a negative number?

With real numbers, no.

$2^2 = 4$

$(-2)^2 = 4$

$0^2 = 0$

- The square of a real number is never negative.
- So the equation $x^2 = -1$ has no real solution.
- We need a new kind of number to solve it.

The imaginary unit

Mathematicians defined a new number i so that:

$$i^2 = -1$$


$$i = \sqrt{(-1)}$$

Now $x^2 = -1$ has two answers:
x = i and x = –i

Source: Abramson (2021a)

Presenter: [STUDENT 1]

What Is a Complex Number?



- A complex number is a real number plus an imaginary number.
- a and b are real numbers.
- If b = 0, then z is just a real number.

Examples

$z = 3 + 4i$

real part 3, imaginary part 4

$z = 2 - 5i$

real part 2, imaginary part −5

Source: Abramson (2021a); Stewart, Redlin and Watson (2016)

Presenter: [STUDENT 2]

Understanding Real and Imaginary Parts

Complex number z	Real part Re(z)	Imaginary part Im(z)
$z = 5 + 2i$	5	2
$z = -3 + 7i$	−3	7


Be careful

- The imaginary part is only the number in front of i, without the i. For z = 5 + 2i it is 2, not 2i.
- Each part keeps its own sign, so for z = −3 + 7i the real part is −3.

Source: Abramson (2021a)

Presenter: [STUDENT 2]

The Complex Plane



Also called the Argand plane.

- Horizontal axis: the real axis, shows a.
- Vertical axis: the imaginary axis, shows b.
- Each complex number is a point: a + bi is drawn at (a, b).

To plot 3 + 4i: go 3 to the right, then 4 up.

Source: Abramson (2021b)

Presenter: [STUDENT 3]

What Is a Complex Conjugate?

Definition

$$If \quad z = a + bi$$

$$then \quad \bar{z} = a - bi$$

$\bar{z}$ is read "z bar".

How to remember it

- Keep the real part.
- Change the sign of the imaginary part.
- Nothing else changes.

Example

$$z = 3 + 4i$$




$$\bar{z} = 3 - 4i$$

The 3 stays. Only the sign of 4i changes.

Source: Abramson (2021a); Brown and Churchill (2014)

Presenter: [STUDENT 3]

Visualizing the Complex Conjugate



- Same real part: both points are 3 units to the right.
- Opposite imaginary part: 4 units up becomes 4 units down.
- The real axis is a mirror: the conjugate is the reflection of z across it.

If you fold the graph along the real axis, z lands on $\bar{z}$.

Source: Brown and Churchill (2014)

Presenter: [STUDENT 4]

Basic Properties of Complex Conjugates

1. Conjugate of the conjugate

$$\overline{\overline{z}} = z$$

Changing the sign twice gives back the original number.

Example:

$z = 2 + 3i$

$\overline{z} = 2 - 3i$

conjugate again: $2 + 3i$

2. Conjugate of a sum

$$\overline{(z_1 + z_2)} = \overline{z_1} + \overline{z_2}$$

Add first or conjugate first, the result is the same.

Example:

$z_1 = 1 + 2i,\ z_2 = 3 + i$

$z_1 + z_2 = 4 + 3i \rightarrow 4 - 3i$

$(1 - 2i) + (3 - i) = 4 - 3i$

3. Conjugate of a product

$$\overline{(z_1 z_2)} = \overline{z_1}\ \overline{z_2}$$

Multiply first or conjugate first, the result is the same.

Example:

$z_1 = 1 + i,\ z_2 = 1 + 2i$

$z_1 z_2 = -1 + 3i \rightarrow -1 - 3i$

$(1 - i)(1 - 2i) = -1 - 3i$

Source: Brown and Churchill (2014)

Presenter: [STUDENT 4]

Simple Examples of Complex Conjugates

Example 1

$$z = 6 + 2i$$



$$\bar{z} = 6 - 2i$$

The + in front of 2i becomes −.

Example 2

$$z = -4 - 3i$$



$$\bar{z} = -4 + 3i$$

The real part −4 does not change. Only −3i becomes +3i.

Example 3

$$z = 7$$



$$\bar{z} = 7$$

There is no imaginary part (b = 0), so nothing changes. A real number is its own conjugate.

Source: Abramson (2021a)

Presenter: [STUDENT 5]

Introduction to Absolute Value

For real numbers, absolute value is the distance from 0:




For a complex number, it is the distance from the origin:

If $z = a + bi$,

$$|z| = \sqrt{(a^2 + b^2)}$$

- |z| is also called the modulus of z.
- It is always a real number, zero or positive.

Where does the formula come from?




$$c^2 = a^2 + b^2$$

Pythagoras: |z| is the longest side of a right triangle.

Source: Abramson (2021b); Stewart, Redlin and Watson (2016)

Presenter: [STUDENT 5]

Visualizing Absolute Value



Reading the triangle

- horizontal side: real part = 3
- vertical side: imaginary part = 4
- long side: the distance |z|

$$|z| = \sqrt{(3^2 + 4^2)}$$
$$= \sqrt{(9 + 16)} = \sqrt{25}$$
$$= 5$$

Source: Abramson (2021b)

Presenter: [STUDENT 6]

Calculating Absolute Value: Easy Examples

Example 1

$$z = 3 + 4i$$

$\lvert z \rvert = \sqrt{(3^2 + 4^2)}$	square each part
$= \sqrt{(9 + 16)}$	add
$= \sqrt{25}$	square root

$= 5$

Example 2

$$z = 5 + 12i$$

$\lvert z \rvert = \sqrt{(5^2 + 12^2)}$	square each part
$= \sqrt{(25 + 144)}$	add
$= \sqrt{169}$	square root

$= 13$

Source: Abramson (2021b)

Presenter: [STUDENT 6]

Properties of Absolute Value

$|z| \geq 0$

A distance can never be negative.

Example: $|3 + 4i| = 5$

$|z| = 0$ only when $z = 0$

Only the origin is at distance zero.

Example: $|0 + 0i| = 0$

$|\bar{z}| = |z|$

A reflection is the same distance from the origin.

Example: $|3 - 4i| = 5 = |3 + 4i|$

$|z_1 z_2| = |z_1|\,|z_2|$

When we multiply numbers, their sizes multiply.

Example: $|(3 + 4i)\cdot 2| = |6 + 8i| = 10 = 5 \times 2$

Source: Brown and Churchill (2014)

Presenter: [STUDENT 7]

The Important Connection

$$z \cdot \bar{z} = |z|^2$$

Why is this useful?

- The result is always a real number (no i).
- It gives $|z|^2$ without a square root.
- We use it for division:

$1 \div (3 + 4i) = (3 - 4i) \div 25$

Check with z = 3 + 4i

$z \cdot \bar{z} = (3 + 4i)(3 - 4i)$

$= 9 - 12i + 12i - 16i^2$

$= 9 - 16i^2$ — middle terms cancel

$= 9 + 16$ — since $i^2 = -1$

$= 25$

and $|z|^2 = 5^2 = 25$. Same result.

Source: Brown and Churchill (2014); Stewart, Redlin and Watson (2016)

Presenter: [STUDENT 7]

Conjugate vs. Absolute Value


	Complex conjugate $\bar{z}$	Absolute value $\lvert z \rvert$
What it does	Changes the sign of the imaginary part	Gives the size of z
Result	Another complex number	A real number, zero or positive
Formula	$\bar{z} = a - bi$	$\lvert z \rvert = \sqrt{(a^2 + b^2)}$
On the graph	Reflection across the real axis	Distance from the origin
For z = 3 + 4i	$\bar{z} = 3 - 4i$	$\lvert z \rvert = 5$

Source: Abramson (2021a; 2021b)

Presenter: [STUDENT 8]

Real-Life Applications

- Electrical engineering: AC voltage and current are written as complex numbers.
- Digital signal processing: audio and image filters work with complex values.
- Communication systems: Wi-Fi and mobile signals carry data in magnitude and phase.
- Sound analysis: each frequency in a sound has a size and a timing.



Length of the arrow = magnitude, which is $|z|$
Angle of the arrow = phase

So one complex number can store two facts about a signal: its size and its timing.

Source: Alexander and Sadiku (2017); Stroud and Booth (2020)

Presenter: [STUDENT 8]

Practice and Quick Quiz

Try to answer each question first. The answer appears on the next click.

1. Find the conjugate of

$$z = 4 + 6i$$

$$\bar{z} = 4 - 6i$$

2. Find the absolute value of

$$z = 3 + 4i$$

$$|z| = 5$$

3. Find the conjugate of

$$z = -2 - 5i$$

$$\bar{z} = -2 + 5i$$

4. Calculate $|z|$ if

$$z = 5 + 12i$$

$$|z| = 13$$

Source: Questions written by the group, based on Abramson (2021a; 2021b)

Presenter: [STUDENT 9]

Conclusion

- A complex number has a real and an imaginary part: $z = a + bi$.
- The conjugate changes the sign of the imaginary part: $\bar{z} = a - bi$.
- The absolute value is the distance from the origin: $|z| = \sqrt{(a^2 + b^2)}$.
- The identity $z\bar{z} = |z|^2$ connects the two ideas.
- Complex numbers are used in mathematics, IT and engineering.

Thank you

Any questions?

References (Harvard style)

Abramson, J. (2021a) 'Section 3.1: Complex numbers', in *Precalculus 2e*. Houston, TX: OpenStax. Available at: https://openstax.org/books/precalculus-2e/pages/3-1-complex-numbers (Accessed: 3 November 2026).

Abramson, J. (2021b) 'Section 8.5: Polar form of complex numbers', in *Precalculus 2e*. Houston, TX: OpenStax. Available at: https://openstax.org/books/precalculus-2e/pages/8-5-polar-form-of-complex-numbers (Accessed: 3 November 2026).

Alexander, C.K. and Sadiku, M.N.O. (2017) *Fundamentals of electric circuits*. 6th edn. New York: McGraw-Hill Education.

Brown, J.W. and Churchill, R.V. (2014) *Complex variables and applications*. 9th edn. New York: McGraw-Hill Education.

Stewart, J., Redlin, L. and Watson, S. (2016) *Precalculus: mathematics for calculus*. 7th edn. Boston, MA: Cengage Learning.

Stroud, K.A. and Booth, D.J. (2020) *Engineering mathematics*. 8th edn. London: Red Globe Press.