

B.S., University of Toledo (2012) May 2019 May 1, 2019 Assistant Professor

ABSTRACT

Since the discovery of space plasmas, which are ionized gasses in outer space, the space science community have been deriving fluid models of plasmas to describe their properties and motion in various space environment and processes. The simplest case is the ideal magnetohydrodynamic (MHD) model which assumes a collisionless perfectly-conducting plasma under the influence of electromagnetic fields. Multi-fluid collisional MHD treats each fluid with its own set of MHD equations while using collisions to describe the mass, momentum, and energy interactions among them in processes such as friction and resistivity. One of the important examples of such multi-fluid plasma medium is the solar chromosphere. The high density and relatively cold plasma result in a partially ionized medium with strong collisions among different species so that the collision times are shorter than characteristic time scales of charged particles moving in the electromagnetic field. Multi-fluid collisional MHD has many cases from the Earth's ionosphere to stellar atmosphere and having a good grasp of the physics is important in studying these plasmas.

In this dissertation I go through the physics of collisional plasmas, designing the Collisional Multi-Fluid ion (CoMFi) code, with scientific computing principles, and its application on the solar chromosphere. With this simulation I found that it is possible for shocks and jets to form in the lower solar chromosphere through frictional heating alone and not with magnetic reconnection.

ACKNOWLEDGEMENTS

To my partner Alexis Ploss, who believed in me without judgment or apprehension. To my family, that supported me beyond the ocean. To my dog Ada Lovelace, that provided emotional support through the hardships.

Contents

I	INTRODUCTION	1
I.1	The Sun	2
I.1.1	The Chromosphere	3
I.2	Space Weather	3
I.2.1	Motivation: A storm is coming	4
I.2.2	Where does the chromosphere fit into this?	5
II	COLLISIONAL PLASMAS	7
II.1	Magnetohydrodynamics	7
II.1.1	Kinetic Theory	8
II.1.2	Moments of the Distribution	9

II.1.3	Multi-fluid MHD	11
II.1.4	Temperature Equation	13
II.1.5	Two-fluid MHD	13
II.2	Maxwell's Equations	16
II.2.1	Generalized Ohm's Law	17
II.2.2	Induction Equation	19
III	IMPLEMENTATION	21
III.1	Time Integration	21
III.1.1	Stability	23
III.2	Schemes	26
III.2.1	Flux Limiters	27
III.2.2	TVD-MUSCL	31
III.3	Maintaining a Divergence Free Field	32
III.3.1	Generalized Lagrange Multiplier	33
III.4	Architecture	35
III.4.1	Data Structure	36
III.5	Normalization	38
III.5.1	Constants	38
III.5.2	Normalized Equations	39
III.6	Validation	40
III.7	Visualization	42
III.7.1	Storage	42
III.7.2	Python Plot Scripts	42
IV	CHROMOSPHERE DYNAMICS	46
IV.1	Governing Equations	48

IV.2 Simulation Domain	51
IV.3 Results	54
IV.4 Discussion	58
V CONCLUSION	66
V.1 Future Work	67
LITERATURE CITED	76

List of Figures

I.1 A more complicated view of the sun's structure and layers. [Image from https://phys.org/]	2
I.2 A view of the upper, middle and lower chromosphere. The chromosphere is partially ionized and is much colder than the corona (above the transition region). [Song and Vasyliūnas, 2014, Avrett and Loeser, 2008a]	4
I.3 Filaments. These are images taken in H-alpha. [Image taken from https://solarscience.msfc.nasa.gov/]	6
III.1 A Sweby diagram of the current limiters in the CoMFi code. The shaded region is the required region for the limiter to be in in order to be second-order TVD.	30
III.2 Spurious oscillations that may form in the advection equation due to discretization errors. An explanation of this can be found in Section III.1.1. Imagen taken from [http://farside.ph.utexas.edu/].	31

III.3 Neutral fluid shock tube tests. The shocks are being captured appropriately due to the TVD-MUSCL scheme used.	44
III.4 The Soler et al. [2013] initial value problem tests. Velocity is taken at $z = 0$ of the standing Alfvén wave. The solid red is the ion velocity, the dashed blue line is neutral velocity. The top row is where the initial neutral velocity is the same as the ion velocity and the bottom row is where the initial neutral velocity is at zero. The results resemble reasonably well the tests in the Soler et al. [2013] paper.	45
III.5 Taken from Soler et al. [2013]. Compare with above (Figure III.4) for validation.	45
IV.1 Formation of a canopy seen after around 348ms (13000 time steps). The initial condition of a concentrated strong field region (1 kG) and weak field region (5G) creates vertical asymmetry as plasma beta drastically decreases. The left and right boundaries are mirrored so effects of other granules behaving the same way are considered. Because of the short time period this is <i>not</i> a relaxed system but simply a consequence of the pressure imbalance frequently seen in the chromosphere.	55
IV.2 A zoom into the shock wave front of the jet. The arrows represents the direction and magnitude of ion flow. The color bar represents the neutral number density in cm^{-3}	55
IV.3 Heating of the shock origin before the formation of the jet at $t=18.9\text{s}$. Top image is the frictional heating per neutral particle. Bottom image is the ohmic heating per neutral particle. The frictional heating is orders of magnitude larger than the heating due to currents.	56

IV.4	These figures show the jet in the middle of its journey to the corona at $t=26.8s$. From top to bottom, the temperature of the neutral plasma, the frictional heating per neutral particle, and the difference in speeds $\ \mathbf{V} - \mathbf{U}\ ^2$ of the charged and neutral plasma. The shock can be clearly seen in the bottom-left corner of the simulation where the magnetic flux concentration is strongest. Due to large plasma beta a ‘ring’ of slow moving plasma is formed and persists in the shock allowing the shock to start forming.	57
IV.5	Time series of the shock. From left to right and top to bottom the times are $t=26.751s$, $26.785s$, $26.831s$, $26.842s$, $26.844s$, $26.849s$. The white streamlines denote the magnetic field lines and the number density of the neutrals are in cm^{-3} . This shows them magnetic field geometry changing due to the jet and not the other way round. The jet is not generated because of the magnetic field.	58
IV.6	Time series of frictional heating in the jet. From left to right and top to bottom the times are $t=26.751s$, $26.785s$, $26.831s$, $26.842s$, $26.844s$, $26.849s$. The white streamlines denote flow.	59
IV.7	Ion number density snapshots of shock wave front at line of extraction $x=0.6Mm$. Ion number density is in units of $10^{11} cm^{-3}$. The wavefront can be clearly seen climbing the Chromosphere.	60
IV.8	Blue line is ion number density and the black and gray dashed lines are B_x and B_z respectively at $t=26.83s$ and vertical line at $x=0.6Mm$. Ion number density is in units of $10^{11} cm^{-3}$. The magnetic field is changed behind the shock wave front.	60

IV.9 Blue line is ion number density and red dashed line is frictional heating per neutral particle at $t=26.83s$ and vertical line $x=0.6Mm$. Ion num- ber density is in units of 10^{11} cm^{-3} . The frictional heating per neutral particle behind the wavefront is orders of magnitude higher than in front of it.	61
IV.10 Blue line is ion number density and black dashed line is vertical ion velocity at $t=26.83s$ and vertical line $x=0.6Mm$. Ion number density is in units of 10^{11} cm^{-3} . The jet speed is supersonic.	62

List of Tables

III.1 Normalization Constants	38
III.2 CoMFi.hdf5 data hierarchy	43

I. INTRODUCTION

The word ‘plasma’ was first used by Langmuir [1928] to describe oscillations in a gaseous electron medium [Tonks, 1967]. It is a state of matter that is can be described as an ionized gas. It is the most common form of baryonic matter in the universe. We default to plasma descriptions of the medium between stars and planets. The sun is, for example, made mostly of ionized hot plasma. It is made of plasma and shoots out supersonic plasma called the solar wind. This plasma does not entirely contact the Earth due to shielding resulting from the Earth’s magnetic field. Due to plasma’s ionization, it has a tendency to travel on magnetic field lines. This does not mean the Earth is completely shielded from the solar wind. Many space weather events are due to solar plasma’s effect on the Earth and its magnetic field. The sun is a very active star that has spontaneous flares, coronal mass ejections (CME) and active regions. Thus spontaneous storms occur on Earth and our understanding of the sun and space plasmas is essential to protect ourself against storms.

In this dissertation, an attempt will be made to show the benefits and utility of multi-fluid magnetohydrodynamic (MHD) models. We will show that the solar chromosphere is a good case for this. The implementation will also be studied with careful use of good, modern computer science practices.




Figure I.1: A more complicated view of the sun's structure and layers. [Image from <https://phys.org/>]

I.1 The Sun

Whenever I say ‘the sun’ I do not mean any star—I mean *our* star. The sun is a 4.6 billion year old yellow dwarf. It generates a lot of the energy we use for life on Earth. It creates this energy through nuclear fusion in its core, turning hydrogen into helium. The sun is mostly hydrogen with a small amount of helium and even smaller amounts of metal ions which require tremendous energy to fuse. The sun has six regions (from innermost to outermost): the core, the radiative zone, the convection zone, the photosphere, the chromosphere and the corona. The core is where the fusion happens. Here is where hydrogen is made into helium and other metals. The radiative zone and convection zone is where the energy travels radially upwards towards the surface, the photosphere. The photosphere is the surface of the Sun. Here photons are trapped but the ones that do escape mimic black-body radiation (not perfectly). It is optically thick and is the white light we see when looking at the Sun. Above the photosphere is the atmospheric layer, the chromosphere, which I will go in

much detail about in this dissertation. Above that, is the corona which is what gives the sun its tendrils and the hair we give our cartoon drawings of the sun.

I.1.1 The Chromosphere

The chromosphere is the interface between the surface, the photosphere, and the outer atmosphere, the corona. The plasma in it is partially ionized which means it consists of two kinds of plasma: ionized (charged) plasma, and neutral (uncharged) gas. This means the charged particles are affected by Lorentz forces while the uncharged particles are not. This is a result of the temperature of the chromosphere and the composition of the plasma convected in the convection zone. The plasma has a high temperature in Earth standards but to the sun it is relatively cold. The temperature of the corona, which keep in mind is the outermost layer, is in the million of degrees in Kelvin. While most of the chromosphere and the photosphere is 5000K-6500K. This discrepancy in the temperature of the lower layers to the higher layers is called the coronal heating problem. The coronal heating problem is an outstanding problem in physics which makes understanding the chromosphere of special importance. Why is the corona hotter than the chromosphere? This work aims to shed light on this unsolved problem in solar physics.

I.2 Space Weather

The term space weather refers to weather effects found above our immediate atmosphere which is conversely named terrestrial weather. It focuses on the conditions and changes found between the Sun, the Earth and even behind the Earth along its night time. The Earth's surface magnetic field strength is in the 0.5G range [Finlay et al., 2010] and that is strong enough to shield most of the solar wind headed our direction. However, magnetic reconnection events between the Earth's magnetosphere




Figure I.2: A view of the upper, middle and lower chromosphere. The chromosphere is partially ionized and is much colder than the corona (above the transition region). [Song and Vasyliūnas, 2014, Avrett and Loeser, 2008a]

(the sphere of Earth's magnetic influence) and the interstellar medium allows solar wind particles to enter the Earth's atmosphere. Massive eruptions from the sun can affect the Earth's magnetic fields. These solar events manifest itself in harmless weather, like auroras, or more harmful weather, like geomagnetic storms.

I.2.1 Motivation: A storm is coming

In 1859, a famous (or infamous) geomagnetic storm hit the Earth. Fortunately, we did not sprawl global electrical power grids at the time or it would have had devastating effects. It was caused by a CME observed by Richard C. Carrington and was thusly named the Carrington Event. Depending on the strength of the geomagnetic storm (which happens frequently at higher/lower latitudes due to night side reconnection) it may reach mid-latitude areas which we inhabit. Here geomagnetic

storms due to the quick changes in magnetic fields and the law of induction cause ground induced currents (GIC) that can have damaging effects to any ground based electrical devices.

The Carrington event was our last encounter with a storm of that scale, that does not mean we have not had any near misses. In 2012, a coronal mass ejection that has the strength and size comparable to the previously mentioned Carrington event almost hit the Earth. It missed the Earth by nine days [Baker et al., 2013]. Unlike in 1859, the world of 2012 was filled with electrical power grids. The internet is part of our daily lives and still relies on ground based electrical wiring.

A report by Predictive Science Inc. shows that the chance of another Carrington level storm striking the earth between 2012 and 2022 is 12% [Riley, 2012].

I.2.2 Where does the chromosphere fit into this?

Since the chromosphere is the intermediary layer between the surface and the coronal with entirely different physics, this means it is unfair to disentangle solar events from it.

An important solar event that deals with the chromosphere are solar flares. Solar flares are sometimes accompanied by CMEs. Solar flares are bright flashes that usually occur near sun-spots. They emit electromagnetic energy of the order of 10^{20} joules. Their x-rays can cause disruptions in communication devices on Earth like GPS satellites since it can have effects on the Earth's ionosphere.

A magnetic reconnection even close to the upper chromosphere may cause flares, and the physics of the upper chromosphere is very different than that of the corona. For example the former is occurring in partially ionized plasma and the latter in a fully ionized one.

Filaments are a feature found in chromosphere of dense cool plasma that are

associated by bright emissions in the chromosphere. These filaments can sometimes shoot off into the solar wind by a CME.




Figure I.3: Filaments. These are images taken in H-alpha. [Image taken from <https://solarscience.msfc.nasa.gov/>]

Since the chromosphere is a relatively understudied part of the chromosphere but still has an important role in space weather, this dissertation will focus on how to model, simulate, and study such a region while at the same time focussing on modern techniques of scientific computing.

II. COLLISIONAL PLASMAS

Although plasmas are classically defined as charged gasses, they can coexist with neutral plasma based on the temperature and pressure conditions surrounding them. This means I have two gas mediums, one with the ability to feel Lorentz forces and another that acts like a non-charged fluid. This will generate friction between the fluids when they are moving at different velocities. Due to collision rates being proportional to fluid density and temperature, there can exist a limit in which collision rates are larger than ionization/recombination rates of ions and therefore the fluid becomes collisional. In the reverse case a single-fluid is an apt description as the existence of two-fluids matters largely on the chemistry rather than the collisions. We will be focusing on hydrogen plasmas for our chromosphere case in Chapter IV and in that case two-fluids will be investigated.

II.1 Magnetohydrodynamics

Magnetohydrodynamics (MHD) is a fluid approximation of plasma popular amongst macroscopic studies of space plasmas. In order to arrive at the fluid approximation I must first look at the particles first and ensure the approximation is appropriate for our case. All of physics is a model and to go from the kinetic theory of plasmas by modeling particles to MHD plasmas by modeling fluids there must be a derivation since there are less assumptions in the kinetic theory than there are in

MHD.

II.1.1 Kinetic Theory

We start by assuming an ensemble phase space of particles $f(\mathbf{x}, \mathbf{v}, t)$ called the distribution function. \mathbf{x} is the location of the particle distribution in space, \mathbf{v} is the velocity distribution and t is the time. To study the distribution function's evolution through time I take the full derivative with respect to time $\frac{df}{dt}$ which by the product rule turns out to be:

$$\frac{\partial f}{\partial t} + \frac{\partial \mathbf{x}}{\partial t} \cdot \nabla_{\mathbf{x}} f + \frac{\partial \mathbf{v}}{\partial t} \cdot \nabla_{\mathbf{v}} f = \left(\frac{\delta f}{\delta t} \right)_{\text{collisions}} \quad (1)$$

This is called the Boltzmann equation [Lerner, 2005]. The term on the right hand side exists due to collisions changed the distribution function in its own way, this also include collisional ionization and other collisional effects that would change the distribution of particles. Simplifying the equation further by substituting how I expect the bulk particles to interact with fields I can get:

$$\frac{\partial f}{\partial t} + \mathbf{v} \cdot \nabla_{\mathbf{x}} f + \frac{q}{m} (\mathbf{E} + \mathbf{v} \times \mathbf{B}) \cdot \nabla_{\mathbf{v}} f = \left(\frac{\delta f}{\delta t} \right)_{\text{collisions}} \quad (2)$$

Where q is the charge of the particle, m is the mass of the particle, \mathbf{E} is the electric field and \mathbf{B} is the magnetic field.

This is the kinetic theory of plasmas. The assumptions made here are probability distributions of particles instead of singular particle effects. There are simulations and codes that model particles directly but because of the large memory size of such simulations the region of study must be very small.

II.1.2 Moments of the Distribution

In order to arrive at the macroscopic variables I must make an assumption about the distribution of the fluid. This is the most fundamental assumption of MHD is that the distribution of the particles are behaving like a Maxwellian velocity distribution [Mandl, 1988]. If it is not the case that the velocity distribution is Maxwellian then the approach of MHD is challenged and must be rewritten and integrated differently. The Maxwellian velocity distribution is common in space plasmas that are in thermal equilibrium with itself which lends to the popularity of MHD.

$$f(v) = n \left(\frac{m}{2\pi k_B T} \right)^{3/2} \exp \left(-\frac{mv^2}{2k_B T} \right) \quad (3)$$

Where $v = |\mathbf{v}|$ is the speed, k_B is the Boltzmann constant and T is temperature. This is assuming a Maxwellian velocity distribution over a thermal equilibrium and a thermal speed.

To arrive at the MHD equations I start to take the ‘velocity moments’ of the equations in order to have macroscopic variables that don’t depend on phase speed.

$$Moment_a(\mathbf{x}, t) = \int \mathbf{v}^a f(\mathbf{x}, \mathbf{v}, t) d\mathbf{v} \quad (4)$$

Here a is the order of the moment. The first few moments are of interest. The zeroth moment is the continuity equation.

$$\frac{\partial n_s}{\partial t} + \nabla \cdot (n_s \mathbf{v}_s) = \left(\frac{\delta n_s}{\delta t} \right)_{source} \quad (5)$$

Each species, s , of fluid has its own continuity equations but may be related by the source terms. The source terms come from integrating the collisional ionization, radiative recombination, etc.

The first order moment will give us the momentum equations:

$$\frac{\partial n_s \mathbf{v}_s}{\partial t} + \nabla \cdot (n_s \mathbf{v}_s \mathbf{v}_s + \frac{\mathbf{P}_s}{m_s}) - n_s \frac{q}{m_s} (\mathbf{E} + \mathbf{v}_s \times \mathbf{B}) = \left(\frac{\delta n_s \mathbf{v}_s}{\delta t} \right)_{source} \quad (6)$$

Where \mathbf{P}_s is the kinetic pressure tensor and can be simplified to the ideal gas law $P_s \mathbf{I} = n_s k_B T_s \mathbf{I}$ if applicable. The source terms due to the zeroth order also exists here as well as frictional terms that may appear in multi-fluid cases. It is apparent that this equation is fluid-like, a form of the Navier-Stokes equation and hence the naming of these equations as hydrodynamic.

To close the system equations I need one more for the pressure shown previously. So integrating for the second moment gives:

$$\frac{\partial E_s}{\partial t} + \nabla \cdot (\mathbf{v}_s E_s) + \nabla \cdot \mathbf{q}_s = \left(\frac{\delta E_s}{\delta t} \right)_{source} \quad (7)$$

Where q is the heat flow tensor which can be approached in multiple ways and E is the total energy:

$$E_s = e_s + \frac{1}{2} m_s v_s^2 + \frac{P_s}{\gamma_s - 1} \quad (8)$$

Here e is the internal energy and the potential energy is based on the momentum equation's extra acceleration terms like the fields or gravitational potential energy if included. γ is the ratio of specific heats typically related to the degrees of freedom the particles exhibit. For the monoatomic case I use in this dissertation this number becomes $\frac{5}{3}$.

In most cases, the second moment is enough to close the system of equations. Through the energy equation we may derive an 'equation of state' for the pressure

so we may use it in the momentum equation. The magnetic fields evolve due to Maxwell's equations which will be discussed in Section II.2.

II.1.3 Multi-fluid MHD

In order to make our code capable to solve the equations shown in the previous section (Section II.1.2) further approximations must be done. The three fluids of interest in a hydrogen plasma that includes neutrals are hydrogen ions, electrons and neutral hydrogen. Our plasma is collisional and in partially ionized plasmas I can use the Krook collision term [Bhatnagar et al., 1954].

$$\left(\frac{\delta f}{\delta t}\right)_{collisions} = \nu_n(f_n - f) \quad (9)$$

Where subscript n is the neutral species and ν is the collision rate. We can find the collision rate simply by using the collision cross-section of neutral particles and the average thermal speed. The average thermal speed is used because temperature in the case of MHD is based on the ‘peculiar’ speeds of the particles (i.e. particles with velocity other than that of the bulk plasma). This way peculiar velocities contribute to collisions because if both fluids are moving in the same direction.

$$\nu_n = n_n \sigma_n < v > \quad (10)$$

The thermal speed is a function of temperature and under the Maxwellian velocity distribution assumption, σ is the collisional cross-section which can be found in the literature through experiments. The collisional cross-section between ion and electrons include Coulomb effects as well.

The moment equations (Equation 5-7) for the three species will look like this:

$$\frac{\partial n_i}{\partial t} + \nabla \cdot (n_i \mathbf{v}_s) = S - L \quad (11)$$

$$\frac{\partial n_e}{\partial t} + \nabla \cdot (n_e \mathbf{v}_s) = S - L \quad (12)$$

$$\frac{\partial n_n}{\partial t} + \nabla \cdot (n_n \mathbf{v}_s) = L - S \quad (13)$$

$$\begin{aligned} & \frac{\partial n_i \mathbf{v}_i}{\partial t} + \nabla \cdot (n_i \mathbf{v}_i \mathbf{v}_i + \frac{\mathbf{P}_i}{m_i}) - n_i \frac{q_i}{m_i} (\mathbf{E} + \mathbf{v}_i \times \mathbf{B}) \\ &= - \sum_{i \neq s} [n_i \nu_{is} (\mathbf{v}_i - \mathbf{v}_s)] + \frac{1}{m_i} (S m_n \mathbf{v}_n - S m_e \mathbf{v}_e - L m_i \mathbf{v}_i) \end{aligned} \quad (14)$$

$$\begin{aligned} & \frac{\partial n_e \mathbf{v}_e}{\partial t} + \nabla \cdot (n_e \mathbf{v}_e \mathbf{v}_e + \frac{\mathbf{P}_e}{m_e}) - n_e \frac{q_e}{m_e} (\mathbf{E} + \mathbf{v}_e \times \mathbf{B}) \\ &= - \sum_{e \neq s} [n_e \nu_{es} (\mathbf{v}_e - \mathbf{v}_s)] + \frac{1}{m_e} (S m_n \mathbf{v}_n - S m_i \mathbf{v}_i - L m_e \mathbf{v}_e) \end{aligned} \quad (15)$$

$$\begin{aligned} & \frac{\partial n_n \mathbf{v}_n}{\partial t} + \nabla \cdot (n_n \mathbf{v}_n \mathbf{v}_n + \frac{\mathbf{P}_n}{m_n}) \\ &= - \sum_{n \neq s} [n_n \nu_{ns} (\mathbf{v}_n - \mathbf{v}_s)] + \frac{1}{m_n} (L (m_i \mathbf{v}_i + m_e \mathbf{v}_e) - S m_n \mathbf{v}_n) \end{aligned} \quad (16)$$

The first three equations are the continuity equations describing the number density evolution of the three kinds of species. The last three are the momentum equations describing the forces, pressures and convection for the three kinds of species.

S and L are source and loss terms respectively due to chemistry processes like ionization and recombination. These are three-fluid MHD equations if the only ion considered is the hydrogen ion. They appear in all the equations in some form.

II.1.4 Temperature Equation

We wish to study the thermal evolution of a plasma in Chapter IV. For this I must transform the energy equation (Equation 7) to a form that evolves thermal pressure. We can do this by taking the dot product of the momentum equation and subtracting it from the energy equation. This leaves the energy equation with the thermal energy equation:

$$\frac{\partial}{\partial t} \frac{P_s}{\gamma_s - 1} + \nabla \cdot \left(\frac{P_s}{\gamma_s - 1} \mathbf{v}_s \right) + \left(1 - \frac{P_s}{\gamma_s - 1} \right) \nabla \cdot \mathbf{v}_s = \left(\frac{\delta E_s}{\delta t} \right)_{source} - \mathbf{v}_s \cdot \left(\frac{\delta m_s n_s \mathbf{v}_s}{\delta t} \right) \quad (17)$$

Elastic collisions may take the form of [Banks and Kockarts, 1973]:

$$\left(\frac{\delta E_s}{\delta t} \right)_{collisions} = \mathbf{v}_s \cdot \left(\frac{\delta m_s n_s \mathbf{v}_s}{\delta t} \right)_{collisions} + m_{sn} \nu_{sn} n_s [(3k_B/m_s)(T_n - T_s) + |\mathbf{v}_s - \mathbf{v}_n|^2] \quad (18)$$

Where m_{sn} is the reduced mass of the species with the neutral species, T is temperature, k_B is the Boltzmann constant. The previous equation shows thermal exchange due to collision between the species as well as *frictional heating* term that is proportional to the difference in speeds between the two. For a conservative scheme it is more appropriate to use the total energy equation, but this depends on a case by case basis. The CoMFi code is capable of doing both approaches. For the chromosphere I will be concerned with the thermal evolution.

II.1.5 Two-fluid MHD

The ion and electron fluid can be simplified further to be one plasma that tracks the center of momentum between the two. Plasmas are typically quasi-neutral,

it has as many positive charges as negative ones. Any imbalance in the charge will create a strong electric field within the plasma and strongly attract outside sources of charges until it is quasi-neutral again. In single hydrogen ion case that means:

$$n_i = n_e = n \quad (19)$$

This can simplify the current density equation as well:

$$\mathbf{J} = e(n_i \mathbf{v}_i - n_e \mathbf{v}_e) = en(\mathbf{v}_i - \mathbf{v}_e) \quad (20)$$

Where e is the elementary charge. Subtracting the ion and electron continuity equations gives:

$$\frac{\partial n_i - n_e}{\partial t} + \nabla \cdot (n_i \mathbf{v}_i - n_e \mathbf{v}_e) = 0 \quad (21)$$

The charge conservation equation.

$$\nabla \cdot \mathbf{J} = 0 \quad (22)$$

To simplify the three MHD equations (Equations 14-16) I can take the multiple of their masses and add Equation 14 and 15.

$$\begin{aligned} & \frac{\partial \rho_i \mathbf{v}_i + \rho_e \mathbf{v}_e}{\partial t} + \nabla \cdot (\rho_e \mathbf{v}_e \mathbf{v}_e + \rho_i \mathbf{v}_i \mathbf{v}_i + \mathbf{P}_i + \mathbf{P}_e) - \mathbf{J} \times \mathbf{B} \\ & = -\rho_i \nu_{in}(\mathbf{v}_i - \mathbf{v}_n) - \rho_e \nu_{en}(\mathbf{v}_e - \mathbf{v}_n) + (Sm_n \mathbf{v}_n - Lm_i \mathbf{v}_i - Lm_e \mathbf{v}_e) \end{aligned} \quad (23)$$

The ρ here is the mass density of the fluid. Using the induction equation

Maxwell's equations I can simplify $\mathbf{J} \times \mathbf{B}$ further.

$$\mathbf{J} \times \mathbf{B} = \frac{1}{\mu_0} (\nabla \times \mathbf{B}) \times \mathbf{B} = -\nabla \cdot \left(\frac{B^2}{2\mu_0} \mathbf{I} - \mathbf{B}\mathbf{B} \right) \quad (24)$$

This way I can have a magnetic flux and make use of our conservative scheme in Section III.2.2. Finally by dividing by $m_i + m_e$ to Equation 23 I can find the center of momentum velocity and simplify the equation one last time:

$$\mathbf{V} = \frac{\rho_i \mathbf{v}_i + \rho_e \mathbf{v}_e}{\rho_i + \rho_e} \quad (25)$$

$$\mathbf{v}_i = \mathbf{V} + \frac{\rho_e}{\rho_i + \rho_e} \frac{1}{ne} \mathbf{J} \sim \mathbf{V} + \frac{m_e}{m_i} \frac{\mathbf{J}}{ne} \quad (26)$$

$$\mathbf{v}_e = \mathbf{V} - \frac{\rho_i}{\rho_i + \rho_e} \frac{1}{ne} \mathbf{J} \sim \mathbf{V} - \frac{\mathbf{J}}{ne} \quad (27)$$

$$\begin{aligned} & \frac{\partial \rho \mathbf{V}}{\partial t} + \nabla \cdot \left(\mathbf{K}_{i,e} + \frac{B^2}{2\mu_0} \mathbf{I} - \frac{\mathbf{B}\mathbf{B}}{\mu_0} \right) \\ &= -(\rho_i \nu_{in} + \rho_e \nu_{en}) (\mathbf{V} - \mathbf{U}) + \frac{m_e}{e} (\nu_{en} - \nu_{in}) \mathbf{J} \\ &+ [Sm_n \mathbf{U} - Lm_i (\mathbf{V} + \frac{m_e}{m_i} \frac{\mathbf{J}}{ne}) - Lm_e (\mathbf{V} - \frac{\mathbf{J}}{ne})] \end{aligned} \quad (28)$$

The dynamic pressure and thermal pressures have been combined into one kinetic tensor term \mathbf{K} . \mathbf{U} is the neutral velocity.

The neutral momentum equation is much easier since it is not affected by the

Lorentz force and the momentum exchange is reversed, so it becomes:

$$\begin{aligned}
& \frac{\partial \rho_n \mathbf{U}}{\partial t} + \nabla \cdot (\mathbf{K}_n) \\
&= (\rho_i \nu_{in} + \rho_e \nu_{en})(\mathbf{V} - \mathbf{U}) - \frac{m_e}{e}(\nu_{en} - \nu_{in})\mathbf{J} \\
&- [Sm_n \mathbf{U} - Lm_i(\mathbf{V} + \frac{m_e}{m_i} \frac{\mathbf{J}}{ne}) - Lm_e(\mathbf{V} - \frac{\mathbf{J}}{ne})]
\end{aligned} \tag{29}$$

Finally, following the ideal gas law $P_s = n_s k_B T_s$ and the monoatomic ratio of specific heats $\gamma = 5/3$, the last two equations needed are the temperature equations of the two fluids:

$$\begin{aligned}
& \frac{3}{2}n[\frac{\partial k_B T}{\partial t} + \nabla \cdot (k_B T \mathbf{V})] - \frac{1}{2}n k_B T (\nabla \cdot \mathbf{V}) + \nabla \cdot \mathbf{q} \\
&= nm_{in} \nu_{in} [3k_B (T_n - T) + m_n |\mathbf{U} - \mathbf{V}|^2] + Q
\end{aligned} \tag{30}$$

$$\begin{aligned}
& \frac{3}{2}n_n[\frac{\partial k_B T_n}{\partial t} + \nabla \cdot (k_B T_n \mathbf{U})] - \frac{1}{2}n_n k_B T_n (\nabla \cdot \mathbf{U}) + \nabla \cdot \mathbf{q} \\
&= n_n m_{ni} \nu_{ni} [3k_B (T - T_n) + m_i |\mathbf{V} - \mathbf{U}|^2] + Q_n
\end{aligned} \tag{31}$$

Where \mathbf{q} is the heat flux and Q are other heating and cooling terms such as radiative losses.

II.2 Maxwell's Equations

Since a plasma is a gas that reacts strongly to electro-magnetic fields. Maxwell's equations must be enforced in order to have a plasma consistent with reality. The Maxwell equations will also help us with the time evolution of the fields, as the fields are interconnected. The Maxwell's equations are:

$$\varepsilon_0 \nabla \cdot \mathbf{E} = nq \tag{32}$$

$$\nabla \cdot \mathbf{B} = 0 \quad (33)$$

$$\frac{\partial \mathbf{B}}{\partial t} = -\nabla \times \mathbf{E} \quad (34)$$

$$\mu_0 \varepsilon_0 \frac{\partial \mathbf{E}}{\partial t} + \mu_0 \mathbf{J} = \nabla \times \mathbf{B} \quad (35)$$

In a plasma, any evolution of the electric field is very quickly resolved in the timescales I am considering. Ions and electrons would rearrange quickly to cancel out the electric field so it can be assumed as a perfect electrical conductor. This means the evolution of the magnetic field is more useful than the evolution of the electric field in timescales of interest. This also leads to the approximation $\partial \mathbf{E} / \partial t \sim 0$:

$$\mu_0 \mathbf{J} = \nabla \times \mathbf{B} \quad (36)$$

The plasma is therefore in a static electric field. We will need a definition of the electric field in order to evolve our magnetic fields. This will be discussed in the next Section II.2.1.

The magnetic field has also never been observed to have monopoles which makes it a solenoidal vector field. This is called Gauss's law of magnetism. The magnetic field will need to keep this shape despite errors in the simulation. The implementation will be discussed in Section III.3.

II.2.1 Generalized Ohm's Law

When talking about the Generalized Ohm's Law (GOL) or the Ohm's Law we typically think about conductivity in metal. Or at least that is how it is taught in

college level physics. The Generalized Ohm's Law's purpose is to find the electric field within the plasma. This can be found by subtracting the ion and electron momentum equations 14-15 and multiplying by electron mass.

$$\begin{aligned}
& \frac{m_e}{e} \frac{\partial \mathbf{J}}{\partial t} + \nabla \cdot (m_e n [\mathbf{v}_i \mathbf{v}_i - \mathbf{v}_e \mathbf{v}_e] + m_e \frac{\mathbf{P}_i}{m_i} - \mathbf{P}_e) \\
& - ne \left(-\frac{m_e}{m_i} + 1 \right) \mathbf{E} - \frac{ne m_e}{m_i} \mathbf{v}_i \times \mathbf{B} - ne \mathbf{v}_e \times \mathbf{B} \\
& = -nm_e [\nu_{ie}(\mathbf{v}_i - \mathbf{v}_e) + \nu_{in}(\mathbf{v}_i - \mathbf{v}_n) - \nu_{ei}(\mathbf{v}_e - \mathbf{v}_i) - \nu_{en}(\mathbf{v}_e - \mathbf{v}_n)] \\
& + \left(\frac{m_e}{m_i} - 1 \right) S m_n \mathbf{v}_n - \frac{m_e}{m_i} S m_e \mathbf{v}_e - L m_e \mathbf{v}_i + S m_n \mathbf{v}_n - S m_i \mathbf{v}_i - L m_e \mathbf{v}_e
\end{aligned} \tag{37}$$

Simplifying further by taking the $m_e/m_i \sim 0$ approximation, current density 20, charge conservation 22, and the velocity definitions 26 and 27.

$$\begin{aligned}
& \frac{m_e}{ne^2} \frac{\partial \mathbf{J}}{\partial t} + \frac{m_e}{ne^2} \nabla \cdot [\mathbf{V} \mathbf{J} + \mathbf{J} \mathbf{V}] + \frac{-\nabla \cdot \mathbf{P}_e + \mathbf{J} \times \mathbf{B}}{ne} - \mathbf{E} - \mathbf{V} \times \mathbf{B} \\
& = -\frac{m_e}{e} [\nu_{in}(\mathbf{V} - \mathbf{v}_n) + \nu_{en}(\mathbf{V} - \mathbf{v}_n)] - \eta \mathbf{J}
\end{aligned} \tag{38}$$

$$\begin{aligned}
\mathbf{E} & = -\mathbf{V} \times \mathbf{B} + \frac{m_e}{ne^2} \frac{\partial \mathbf{J}}{\partial t} + \frac{m_e}{ne^2} \nabla \cdot [\mathbf{V} \mathbf{J} + \mathbf{J} \mathbf{V}] \\
& + \frac{-\nabla \cdot \mathbf{P}_e + \mathbf{J} \times \mathbf{B}}{ne} \\
& + \frac{m_e}{e} [\nu_{in}(\mathbf{V} - \mathbf{U}) - \nu_{en}(\mathbf{V} - \mathbf{U})] + \eta \mathbf{J}
\end{aligned} \tag{39}$$

Where $\eta = \frac{m_e \nu_e}{e^2 n}$ is the Ohmic resistivity. The relation between the electric field and the resistivity is what we think of when we invoke Ohm's law in college physics. Clearly the equation is a little more complicated in MHD. Note that collision with neutrals may change the electric field also. The collisions can cause a drift between the ions and electrons, separating them and causing an electric field.

We can take Equation 39 a step further and eliminate the need for an evolving

current and instead use the low-frequency approximation (i.e. the time scales is much smaller than the plasma frequency $\omega_p = \sqrt{\frac{n_e e^2}{\epsilon_0 m_e}}$). Removing the convective derivate of the current gives this approximation:

$$\begin{aligned} \mathbf{E} = & -\mathbf{V} \times \mathbf{B} \\ & + \frac{-\nabla \cdot \mathbf{P}_e + \mathbf{J} \times \mathbf{B}}{ne} \\ & + \frac{m_e}{e} [\nu_{in}(\mathbf{V} - \mathbf{U}) - \nu_{en}(\mathbf{V} - \mathbf{U})] + \eta \mathbf{J} \end{aligned} \quad (40)$$

And in a two-fluid approach I may define current along with the same assumptions:

$$\mathbf{J} = \frac{\nabla \times \mathbf{B}}{\mu_0} \quad (41)$$

The collisional terms may also be ignored under careful consideration of how small $\frac{\nu}{\omega_p^2}$ may be.

In Chapter IV I will simplify the above equation further to suit our needs. Typically in simulations you only want to include the terms that matter the most. With careful consideration I can come up with a GOL that is most appropriate for our application.

II.2.2 Induction Equation

To close our set of equations I need a way to evolve the magnetic fields. For this I can use the law of induction (Equation 34) with the GOL (Equation 40).

$$\begin{aligned} \frac{\partial \mathbf{B}}{\partial t} = & \nabla \times (\mathbf{V} \times \mathbf{B} - \frac{-\nabla \cdot \mathbf{P}_e + \mathbf{J} \times \mathbf{B}}{ne} \\ & - \frac{m_e}{e} [\nu_{in}(\mathbf{V} - \mathbf{U}) - \nu_{en}(\mathbf{V} - \mathbf{U})] - \eta \mathbf{J}) \end{aligned} \quad (42)$$

We may use vector identities to transform the advective terms.

$$\nabla \times (\mathbf{V} \times \mathbf{B}) = \nabla \cdot (\mathbf{B}\mathbf{V} - \mathbf{V}\mathbf{B}) \quad (43)$$

This also gives the opportunity to have as many terms in conservative forms as I can.

$$\begin{aligned} \frac{\partial \mathbf{B}}{\partial t} = & \\ & \nabla \cdot (\mathbf{B}\mathbf{V} - \mathbf{V}\mathbf{B} + \mathbf{B}\mathbf{V}_{hall} - \mathbf{V}_{hall}\mathbf{B}) \\ & - k_B \left(\frac{\nabla n \times \nabla T_e}{ne} \right) \\ & - \nabla \times \left(\frac{m_e}{e} [\nu_{in}(\mathbf{V} - \mathbf{U}) - \nu_{en}(\mathbf{V} - \mathbf{U})] \right) \\ & - \frac{1}{\mu_0} \nabla \eta \times \mathbf{J} + \eta \nabla^2 \mathbf{B} \end{aligned} \quad (44)$$

Where

$$\mathbf{V}_{hall} = \frac{\mathbf{J}}{ne} \quad (45)$$

Since it is associated with the Hall term in the induction equation.

In the coming chapters I will be using a simplified form of the GOL—dropping the terms that are less significant.

III. IMPLEMENTATION

Computational fluid dynamics is a well studied field in computer science, numerics and engineering. Many of its principles can be applied to MHD due to the fluid nature of its equations. We will be focusing on only the ones implemented in the code my for this dissertation, the Collisional Multi-Fluid ion (CoMFi) code [Al Shidi, 2019] was made for the chromosphere case [Al Shidi et al., 2019] which will be explored in more detail in the upcoming chapter (Chapter IV). The code is an ever-evolving code that uses Graphical Processing Units (GPUs) instead of the common CPUs typically seen for MHD codes. This is to implement MHD that makes use of the frontiers of computer technology. GPUs were made for graphical operations in video games. This means they are specifically designed to be very fast at matrix rotations, translations and other linear algebra. In this chapter I will explore proper general purpose GPU programming techniques for solving differential equations and the schemes used alongside it. CoMFi is written in C++ using ViennaCL [Karl Rupp and Selberherr, 2016] for GPU linear algebra template programming and Armadillo [Sanderson and Curtin, 2016] for setting up the matrices on the CPU.

III.1 Time Integration

Time integration is important to the time stepping nature of simulations. We can have both implicit time stepping and explicit time stepping. Explicit time

stepping for the Euler equation takes the form:

$$u^{n+1} = u^n + v \int_n^{n+1} \frac{d}{dx}(u^n) dt \quad (1)$$

This is the easiest approach to the solutions because you use values in the previous time step to solve for the next. This does not require a solution of a linear system which I will show later. Explicit schemes are limited to time steps that are smaller than the wave period, $\Delta x/v > \Delta t$, and depending on the scheme a multiple smaller but usually within the same order of magnitude. This is because the largest characteristic speed (the speed of information propagation) cannot be faster than the grid spacing and time steps. This will cause loss of information and instabilities as there will be an attempt to skip grids but due to discretization it will fail to resolve.

In implicit time steps

$$u^{n+1} - v \int_n^{n+1} \frac{d}{dx}(u^{n+1}) dt = u^n \quad (2)$$

This introduces a linear equation that must be solved, $Au^{n+1} = u^n$, and the solution is to invert the matrix A, $u^{n+1} = A^{-1}u^n$. This gets more complicated in non-linear systems. In this case I am not limited by that characteristic speed.

There is also a third option, semi-implicit schemes. This integrates some variables implicitly and others explicitly. This is helpful when you do not want to be limited by certain characteristic speeds. For the chromosphere example in Chapter IV a characteristic I try to overcome is the collision times since it is orders of magnitude larger than the Alfvén and sound times in the chromosphere.

III.1.1 Stability

In order for schemes to be stable, errors must not grow over time. We can make sure errors are not growing by employing the Von Neumann analysis [CHARNEY et al.]. Let's use a simple explicit forward difference scheme on the wave equation for example.

$$\frac{\partial u}{\partial t} = -v \frac{\partial u}{\partial x} \quad (3)$$

$$u_j^{n+1} = u_j^n - \frac{\Delta t}{\Delta x} v (u_{j+1}^n - u_j^n) \quad (4)$$

The above is the Euler equation for a wave propagating at constant speed v using a forward difference method, also called an upwind scheme. The Von Neumann method takes a Fourier analysis to the error waves generated due to discretization. We say that by using the Taylor series the discretization should have this *exact* solution.

$$u_{j+1} = u_j + \Delta x \frac{\partial u}{\partial x}|_j + \frac{\Delta x^2}{2} \frac{\partial^2 u}{\partial x^2}|_j + \frac{\Delta x^3}{3!} \frac{\partial^3 u}{\partial x^3}|_j + \dots \quad (5)$$

Where j is the location of the cell solution. Rearranging so that it looks like a forward difference scheme shows its relation to the exact solution of the partial derivative.

$$\epsilon = \frac{u_{j+1} - u_j}{\Delta x} - \frac{\partial u}{\partial x}|_j = \frac{\Delta x}{2} \frac{\partial^2 u}{\partial x^2}|_j + \frac{\Delta x^2}{3!} \frac{\partial^3 u}{\partial x^3}|_j + \dots = O(\Delta x) \quad (6)$$

This means the errors, ϵ , are of first order, proportional to Δx so the upwind scheme is a first order scheme in space. To see how the errors evolve in time I employ the Von Neumann analysis. Assuming the computed solution is a linear combination

of the real solution and the error.

$$u_{computed} = u_{exact} + \epsilon \quad (7)$$

We can subtract the exact solution and do a Fourier analysis of the errors found. Doing this on Equation 4 yields.

$$e^{a(t+\Delta t)+ikx} = e^{at+ikx} - \beta(e^{at+ik(x+\Delta x)} - e^{at+ikx}) \quad (8)$$

Where $\beta = v(\Delta t/\Delta x)$, a is the component of the errors in time, and k is the series of frequencies of the errors in space. To find the error growth due to time I must find the ratio $e^{at} = u^{n+1}/u^n$.

$$G = e^{a\Delta t} = 1 - \beta(e^{ik\Delta x} - 1) \quad (9)$$

G is the growth in error each time step also called the amplification factor. In order to keep the errors bounded I need to ensure that this ratio is less than or equal to 1.

$$|G| \leq 1 \quad (10)$$

For the upwind scheme example I used:

$$|1 - \beta(e^{ik\Delta x} - 1)| \leq 1 \quad (11)$$

This will lead to

$$\beta \leq 1 \quad (12)$$

Or the stability requirement, also known as the Courant-Friedrich-Lewy condition:

$$v \frac{\Delta t}{\Delta x} \leq 1 \quad (13)$$

An explicit central difference scheme behaves differently, however.

$$u_j^{n+1} = u_j^n - \beta \frac{u_{j+1}^n - u_{j-1}^n}{2} \quad (14)$$

Doing the Von Neumann analysis here using a more simplified notation:

$$u^{n+1} e^{ijk} = u^n e^{ijk} - \beta \frac{u^n e^{i(j+1)k} - u^n e^{i(j-1)k}}{2} \quad (15)$$

$$G = 1 - \beta \frac{e^{ik} - e^{-ik}}{2} = 1 - i\beta \sin(k) \quad (16)$$

$$|1 - i\beta \sin(k)| = 1 + \beta^2 \sin^2(k) \not\leq 1 \quad (17)$$

Since the amplification factor is always larger than 1, errors will always grow. This scheme is *unconditionally unstable*. Doing this scheme in an implicit fashion can fix this problem.

$$u_j^{n+1} = u_j^n - \beta \frac{u_{j+1}^{n+1} - u_{j-1}^{n+1}}{2} \quad (18)$$

Now doing the Fourier analysis,

$$u_j^{n+1} - \beta \frac{u_{j+1}^{n+1} - u_{j-1}^{n+1}}{2} = u_j^n \quad (19)$$

$$u^{n+1}e^{ijk} - \beta \frac{u^{n+1}e^{i(j+1)k} - u^{n+1}e^{i(j-1)k}}{2} = u^n \quad (20)$$

$$G(1 - \beta \frac{e^{ik} - e^{-ik}}{2}) = 1 \quad (21)$$

$$G = \frac{1}{1 - i\beta \sin(k)} \quad (22)$$

$$|G| = \frac{1}{1 + \beta^2 \sin^2(k)} \leq 1 \quad (23)$$

Since the amplification factor is always less than or equal to 1 this implicit central difference scheme is *unconditionally stable*.

III.2 Schemes

CoMFi uses the conservative MHD equations therefore conservative schemes are of interest. This makes use of the conservative variables unknowns as opposed to the primitive variables (nv as opposed to v). This way I can use a finite volume method [LeVeque et al., 2002] and conserve these quantities to the machine rounding error. Conservative schemes are also known to do well at capturing shocks and the chromosphere is a region where I expect such shocks to exist.

$$\frac{\partial \mathbf{u}}{\partial t} + \nabla \cdot \mathbf{F}(\mathbf{u}) = 0 \quad (24)$$

The above equation is a general Euler equation demonstrating a conservation law problem. The quantity to be conserved is \mathbf{u} and is a function the flux \mathbf{F} . We will be considering the linear case for simplicity however MHD has demonstrably non-

linear terms as the flux (e.g. Equation 14) but it can easily follow from this linear case. The point is to show the total volume is conserved:

$$\frac{d\mathbf{u}}{dt} + \frac{1}{V} \oint \mathbf{F} \cdot \mathbf{n} dS \quad (25)$$

In codes designed to solve differential equations numerically I must decide on the discretization. Since I am using a finite volume method I am discretizing the domain by cells with density quantities in them. The code is discretized in cells existing in Cartesian coordinates (rectangular cells) so I am tasked to find the values of the variables at the cell faces in order to conserve the volume.

We would also like a total-variation-diminishing (TVD) scheme [Harten, 1997], this is because spatial derivatives are approximated on a computer. If I take a Taylor series of a derivative from one cell to another and compare with an approximate derivative like $\frac{u(x+\Delta x) - u(x-\Delta x)}{2\Delta x}$ I see that there are many higher order terms missing. Taking the Fourier analysis of these terms show that any high-frequency errors may compound (sometimes referred to as spurious oscillations). This can be fixed by schemes that ensure the total variation TV is diminishing.

$$\frac{d(TV)}{dt} = \frac{d}{dt} \int \left| \frac{\partial u}{\partial x} \right| dx \leq 1 \quad (26)$$

Therefore, for the purpose of our code, I will use the Total-Variation-Diminishing Monotonic Upwind Scheme for Conservation Laws (TVD-MUSCL).

III.2.1 Flux Limiters

To conserve the cell's quantities I must be able to approximate the quantities on the cell edges. At the same time I must preserve monotonicity to avoid the oscillations. This can be achieved through flux limiters. The utility of the flux limiters

is to turn the scheme into a first-order upwind scheme in sharp discontinuities and a high-resolution scheme when the solution is smooth.

$$u_{j+1/2} = u_j + \frac{1}{2}\phi(r_j)(u_{j+1} - u_j) \quad (27)$$

$$r_j = \frac{u_j - u_{j-1}}{u_{j+1} - u_j} \quad (28)$$

Subscript j is the discretized cell location with quantity u . The ratio of successive gradients r determines how strongly the flux limiter acts to find the extrapolated cell edge variable. The flux limiter ϕ includes the cells around edge more readily the smoother the solution is ($r > 0$), or sharp it is ($r < 0$) in which case the flux limiter is zero ($\phi = 0$). These are the properties flux limiters must abide in order to turn the scheme upwind in sharp gradients or high-resolution in smoother gradients. The above equation is a linear reconstruction of a cell edge. Other reconstructions exist but are more diffusive and avoids the shock-capturing goal of the code.

We have some freedom in choosing how to set the flux limiter. Many flux limiters have been made and studied in the past. The simplest flux-limiter is the minmod.

$$\phi_{minmod}(r) = \max(0, \min(1, r)), \lim_{r \rightarrow \infty} \phi(r) = 1 \quad (29)$$

Implementing these on a computer however is not a trivial task. Many initial conditions require the entirety of the domain be the same number for instance. In that case I get an $r = \frac{0}{0}$ and this causes a computer to return NaN (not a number). Setting to $r = 0$ is appropriate but requires extra coding. A work around is to add a small number ϵ to the denominator known as the machine epsilon to avoid these cases.

The CoMFi code provides several flux limiters. The pre-made flux limiters tend to be symmetric:

$$\frac{\phi(r)}{r} = \phi\left(\frac{1}{r}\right) \quad (30)$$

This is to ensure that it is consistent in negative and positive fluxes.

The provided flux limiters are:

$$\phi_{ospre}(r) = 1.5 \frac{(r^2 + r)}{r^2 + r + 1} \quad (31)$$

This is the Ospre [Waterson and Deconinck, 2007] limiter, the default limiter in the code.

$$\phi_{vanAlbada}(r) = \frac{r^2 + r}{r^2 + 1} \quad (32)$$

By van Albada [1982].

$$\phi_{vanLeer}(r) = \frac{r + |r|}{1 + |r|} \quad (33)$$

By van Leer [1974]

These were chosen because of ViennaCL's limitation of not having element-wise min and max functions. ViennaCL does have the ability to compile custom kernels and examples of min-mod in that case does exist.

Only second-order TVD limiters were chosen because when the scheme is at its high-resolution it is a second-order scheme.

$$r_j = \frac{u_j - u_{j-1}}{u_{j+1} - u_j + \epsilon} \quad (34)$$

This machine epsilon should have no effect on the result so long as the variables




Figure III.1: A Sweby diagram of the current limiters in the CoMFi code. The shaded region is the required region for the limiter to be in in order to be second-order TVD.

have been normalized appropriately (see Section III.5). This means the valuable results are within an order of magnitude so the machine epsilon of a 64bit floating point number of $2^{-52} \sim 2.22 \times 10^{-16}$ would be negligible. If two adjacent cells have the same value, as is common with initial values, then dividing by zero is usefully avoided.




Figure III.2: Spurious oscillations that may form in the advection equation due to discretization errors. An explanation of this can be found in Section III.1.1. Imagen taken from [<http://farside.ph.utexas.edu/>].

III.2.2 TVD-MUSCL

The TVD-MUSCL scheme [van Leer, 1979] reconstructs ‘left’ (L) and ‘right’ (R) states of the cell edge variables.

$$u_{j-1/2}^L = u_{j-1} + \frac{1}{2}\phi(r_{j-1})(u_j - u_{j-1}) \quad (35)$$

$$u_{j+1/2}^L = u_j + \frac{1}{2}\phi(r_j)(u_j - u_{j-1}) \quad (36)$$

$$u_{j-1/2}^R = u_j + \frac{1}{2}\phi(r_j)(u_{j+1} - u_j) \quad (37)$$

$$u_{j+1/2}^R = u_{j+1} + \frac{1}{2}\phi(r_{j+1})(u_{j+1} - u_j) \quad (38)$$

With the left and right extrapolated variables I may construct a flux.

$$F_{j-1/2}^* = \frac{1}{2}([F(u_{j-1/2}^R) + F(u_{j-1/2}^L)] - \lambda_{j-1/2}[u_{j-1/2}^R - u_{j-1/2}^L]) \quad (39)$$

$$F_{j+1/2}^* = \frac{1}{2}([F(u_{j+1/2}^r) + F(u_{j+1/2}^l)] - \lambda_{j+1/2}[u_{j+1/2}^r - u_{j+1/2}^l]) \quad (40)$$

Here $Fu = \lambda u$ are the eigenvalues of the variables. This in physical terms means the largest wave speeds of the system. Typically for the ion plasma it is the fast-mode speed and for the neutrals it is the sound speed. It's important to note in conservative MHD this is only the wave speed (c_{fast}) and not the same as the fastest speed of information propagation ($v + c_{fast}$). The diffusive terms proportional to λ are meant to avoid the oscillations and if they are larger than they need to be the solution will be more diffusive.

III.3 Maintaining a Divergence Free Field

The Gauss's law for magnetic fields (Equation 33) needs to be taken care of because due to machine rounding error and other errors due to discretization, there will be evolution of the divergence. That is to say, although:

$$\frac{\partial \nabla \cdot \mathbf{B}}{\partial t} = -\nabla \cdot \nabla \times \mathbf{E} = 0 \quad (41)$$

The reality is, due to errors:

$$\frac{\partial \nabla \cdot \mathbf{B}}{\partial t} = \epsilon_{errors} \quad (42)$$

There are many methods to resolving this. A popular method is Yee grids, sometimes referred to as staggered grids. Placing the magnetic fields to the cell edges of the electric fields. By doing this, evolving the magnetic fields reduces errors to machine rounding errors since Stokes' theorem is imposed on the problem.

$$\iint_S \frac{\partial \mathbf{B}}{\partial t} \cdot d\mathbf{A} = \oint_{edges} \mathbf{E} \cdot d\mathbf{l} \quad (43)$$

This is not the method used for CoMFi. Staggering the grid makes using a finite volume method difficult especially when trying to find cell centers from one quantity to the next. Instead I employ the Generalized Lagrange Multiplier method introduced by Dedner et al. [2002].

III.3.1 Generalized Lagrange Multiplier

The Generalized Lagrange Multiplier (GLM) method adds another unknown to the system of equations to be solved. This has the drawback of using more memory. Its implementation is simple, however, which means adding it to an existing code is easy. The GLM Φ tries to minimize the monopoles created due to errors in the simulation. It does this by first creating some stress to the induction equation if there is some monopole.

$$\frac{\partial \mathbf{B}}{\partial t} = -\nabla \times \mathbf{E} + \nabla \Phi \quad (44)$$

And filter the monopoles into the GLM using this method of evolution:

$$\mathcal{D}(\Phi) + \nabla \cdot \mathbf{B} = 0 \quad (45)$$

\mathcal{D} is a differential operator and I can choose the GLM to evolve in any way I want. A hyperbolic evolution and/or a parabolic evolution may be used. Due to the non-linear and parabolic nature of our equations I opt to use both:

$$D(\Phi) = \frac{1}{c_h^2} \frac{\partial \Phi}{\partial t} + \frac{1}{c_p^2} \Phi \quad (46)$$

Here c_h and c_p are the hyperbolic and parabolic speeds of the divergence errors. That means the divergence errors will propagate at a speed of c_h and dissipate at rate of c_p . The new equation that must be solved is:

$$\frac{\partial \Phi}{\partial t} + c_h^2 \nabla \cdot \mathbf{B} = -\frac{c_h^2}{c_p^2} \Phi \quad (47)$$

The hyperbolic speed, c_h , can be chosen to be similar the wave speeds in the magnetic field (i.e. the fast-mode speed). We have more freedom to chose the ratio of dissipation to propagation, $\alpha = c_h/c_p^2$, also known as the damping parameter. Through testing 0.5 was find to be a good parameter [Mignone et al., 2010, p. 5899-5911].

We can separate Equation 47 into two parts two solve. We can solve the right hand side exactly, and use the conservative scheme to solve the second term on the left hand side. First solve the left hand side and setting the right hand side equal to zero. Then minimize it with the exact solution:

$$\Phi(t + \Delta t) = \Phi(t)e^{-c_h \alpha \Delta t} \quad (48)$$

The efficacy of this method is keeping the normalized divergence errors to lower than the normalized grid-spacing. This is useful so that the monopoles are not evident when drawing the magnetic field lines nor does it have a strong effect on the results.

III.4 Architecture

It is important in scientific computing not to reinvent the wheel. There are many libraries, particularly C++ libraries, that use template programming to make it easier to write linear algebra in code. It is always easier to add an entire vector with one $+$ sign than looping through each element of the two vectors and adding each. For this reason I can avoid low-level programming like Fortran for CPUs or OpenCL and CUDA for GPUs because the Basic Linear Algebra Subroutines (BLAS) have already been written and abstracted by someone else. C++ is a language that is all about abstraction and speed. There is a pervasive myth that if you want to do scientific computing you must use Fortran or CUDA but this is not true and is usually reinforced by scientists' lack of understanding of computer science and/or bad code.

The code uses to template expression libraries in C++. Template expression is how C++ abstracts operators such as $+$, $-$ and $/$ to deal with one variable and the next and call the low level BLAS functions. The libraries used are Armadillo for CPU. This is a code typically used for Machine Learning applications so speed and big data is its focus. For GPU programming ViennaCL is used, it's a library used by PETSc for its GPU portion. However, at the time of this writing, PETSc's GPU code is not fully developed so ViennaCL is used directly. Luckily, ViennaCL interfaces with Armadillo well and there exists copy constructors to go from shared memory to GPU memory.

Matrices *must* be built on the CPU. Accessing GPU elements individually is

a slow process that is limited by the PCI-e speed and when the machine code (running on CPU) is asked to go back and forth, I get slow code. Best practice is to first build a matrix on the CPU then copy to the GPU.

III.4.1 Data Structure

When using the semi-implicit method a linear system $Ab^{n+1} = b$ must be solved. We must think of an appropriate ordering of our vector in terms of the where the unknowns must be in the elements of the vector. For GPU programming in order to take advantage of its matrix operations speeds due to its many cores I must also be easily able to change back and forth between matrix and vector form between the implicit and explicit portions of the solution. A good ordering for a two-dimensional domain looks like this:

$$b = \begin{bmatrix} u_{0,00} \\ \vdots \\ u_{0,i0} \\ \vdots \\ u_{0,ij} \\ \vdots \\ u_{w,ij} \end{bmatrix} \quad (49)$$

The subscripts i and j describe the location of the cell in the x and y axis

respectively and w are the scalar unknowns.

$$u_w = \begin{bmatrix} B_x \\ B_\perp \\ B_z \\ n_i \\ n_n \\ n_i V_x \\ n_i V_\perp \\ n_i V_z \\ n_n U_x \\ n_n U_\perp \\ n_n U_z \\ T_i \\ T_n \\ \Phi \end{bmatrix} \quad (50)$$

This can be easily resized into a matrix of the form:

$$B = \begin{bmatrix} u_{0,00} & \cdots & u_{w,00} \\ \vdots & \ddots & \vdots \\ u_{0,i0} & \cdots & u_{w,i0} \\ \vdots & \ddots & \vdots \\ u_{0,ij} & \cdots & u_{w,ij} \end{bmatrix} \quad (51)$$

This has the benefits of making good use of GPU programming by using the parallelism found in matrix operations. This also means you can share boundary conditions by taking or providing rows for this matrix in chunks. You can do operations

on specific unknowns by taking the appropriate column.

III.5 Normalization

The unknowns in the simulations must be normalized. This is typical good practice in numerics when we're dealing with computers due to how computers deal with floating point number. Since computers deal with discrete numbers the floating point number line is discrete and also less regular than the real number line. There are more floating point numbers between -1 and 1 than there are between 10 and 11 for example. For this purpose, it is important to find a way in which the common numbers I will be dealing with be normalized to 1. We can do this by having, usually, the largest numbers found in the simulation domain be the normalizing constants.

III.5.1 Constants

The normalization constants were chosen for the chromospheric case I study in Chapter IV. For a more in-depth discussion on why these were chosen see Section IV.2.

Table III.1: Normalization constants chosen for the chromospheric simulation

Magnetic Field	B_0	0.1 T
Length	L_0	2.1×10^6 m
Number Density	n_0	1.0×10^{20} m ⁻³
Mass	$m_0 = m_i$	$1.672621898 \times 10^{-27}$ kg
Vacuum Permeability	μ_0	$1.2566370614 \times 10^{-6}$ Vs/A/m
Charge	e_0	$1.6021766208 \times 10^{-19}$ C
Boltzmann Constant	k_B	1.38065×10^{-23} J/K
Velocity	$V_0 = \frac{B_0}{\sqrt{\mu_0 m_i n_0}}$	2.18×10^8 m/s
Pressure	$p_0 = \frac{B_0^2}{\mu_0}$	8.0×10^3 Pa
Temperature	$T_0 = \frac{p_0}{n_0 k_B}$	5.79438×10^6 K
Current Density	$j_0 = e_0 n_0 V_0$	3.493×10^9 A/m/m
Heat Flow	$q_0 = p_0 V_0$	1.744×10^{12} Pa m/s

In bold are the free parameters you may choose for the CoMFi code. The con-

stants must be chosen carefully so that the largest numbers of interest are normalized to 1. The reason for choosing the quantities above is discussed in Chapter IV.

III.5.2 Normalized Equations

Once I have the normalization constants I may normalize the equations. Replacing each term by its normalized version reduces the equations further. There is less need to include physical constants like μ_0 and k_B as they are implicit in the quantities. Hence, I get normalized MHD equations of the form:

$$\frac{\partial \bar{n}_s}{\partial \bar{t}} + \nabla \cdot (\bar{n}_s \bar{\mathbf{v}}_s) = \bar{S} - \bar{L} \quad (52)$$

$$\begin{aligned} & \frac{\partial \bar{n} \bar{\mathbf{V}}}{\partial \bar{t}} + \nabla \cdot (\bar{\mathbf{K}}_{i,e} + \frac{\bar{B}^2}{2} \mathbf{I} - \bar{\mathbf{B}} \bar{\mathbf{B}}) \\ &= -(\bar{n} \bar{\nu}_{in} + \bar{m}_e \bar{n} \bar{\nu}_{en})(\bar{\mathbf{V}} - \bar{\mathbf{U}}) + \frac{\bar{m}_e}{\bar{e}} (\bar{\nu}_{en} - \bar{\nu}_{in}) \bar{\mathbf{J}} \\ &+ (\bar{S} \bar{m}_n \bar{\mathbf{U}} - \bar{L} (\bar{\mathbf{V}} + \frac{\bar{m}_e}{\bar{n} \bar{e}} \bar{\mathbf{J}}) - \bar{L} \bar{m}_e (\bar{\mathbf{V}} - \frac{\bar{\mathbf{J}}}{\bar{n} \bar{e}})) \end{aligned} \quad (53)$$

$$\begin{aligned} & \frac{\partial \bar{n}_n \bar{\mathbf{U}}}{\partial \bar{t}} + \nabla \cdot (\bar{\mathbf{K}}_n) \\ &= +(\bar{n} \bar{\nu}_{in} + \bar{m}_e \bar{n} \bar{\nu}_{en})(\bar{\mathbf{V}} - \bar{\mathbf{U}}) - \frac{\bar{m}_e}{\bar{e}} (\bar{\nu}_{en} - \bar{\nu}_{in}) \bar{\mathbf{J}} \\ &- (\bar{S} \bar{m}_n \bar{\mathbf{U}} - \bar{L} (\bar{\mathbf{V}} + \frac{\bar{m}_e}{\bar{n} \bar{e}} \bar{\mathbf{J}}) - \bar{L} \bar{m}_e (\bar{\mathbf{V}} - \frac{\bar{\mathbf{J}}}{\bar{n} \bar{e}})) \end{aligned} \quad (54)$$

$$\begin{aligned} & [\frac{\partial \bar{T}}{\partial \bar{t}} + \nabla \cdot (\bar{T} \bar{\mathbf{V}})] - \frac{1}{3} \bar{T} (\nabla \cdot \bar{\mathbf{V}}) + \frac{1}{\bar{n}} \nabla \cdot \bar{\mathbf{q}} \\ &= \bar{m}_{in} \bar{\nu}_{in} [2(\bar{T}_n - \bar{T}) + \frac{1}{3} \bar{m}_n |\bar{\mathbf{U}} - \bar{\mathbf{V}}|^2] + \frac{2}{3} \bar{Q} \end{aligned} \quad (55)$$

$$\begin{aligned} & [\frac{\partial \bar{T}_n}{\partial \bar{t}} + \nabla \cdot (\bar{T}_n \bar{\mathbf{U}})] - \frac{1}{3} \bar{T}_n (\nabla \cdot \bar{\mathbf{U}}) + \frac{1}{\bar{n}_n} \nabla \cdot \bar{\mathbf{q}}_n \\ &= \bar{m}_{ni} \bar{\nu}_{ni} [2(\bar{T} - \bar{T}_n) + \frac{1}{3} \bar{m}_i |\bar{\mathbf{V}} - \bar{\mathbf{U}}|^2] + \frac{2}{3} \bar{Q}_n \end{aligned} \quad (56)$$

$$\begin{aligned}
\frac{\partial \bar{\mathbf{B}}}{\partial t} = & \\
& \nabla \cdot (\bar{\mathbf{B}}\bar{\mathbf{V}} - \bar{\mathbf{V}}\bar{\mathbf{B}} + \bar{\mathbf{B}}\bar{\mathbf{V}}_{hall} - \bar{\mathbf{V}}_{hall}\bar{\mathbf{B}}) \\
& - \frac{\nabla \bar{n} \times \nabla \bar{T}_e}{\bar{n}\bar{e}} \\
& - \nabla \times \left(\frac{\bar{m}_e}{\bar{e}} [\bar{\nu}_{in}(\bar{\mathbf{V}} - \bar{\mathbf{U}}) - \bar{\nu}_{en}(\bar{\mathbf{V}} - \bar{\mathbf{U}})] \right) \\
& - \nabla \bar{\eta} \times \bar{\mathbf{J}} + \eta \nabla^2 \bar{\mathbf{B}}
\end{aligned} \tag{57}$$

III.6 Validation

The simulation now needs to be validated against well known tests. Our goals are to have a scheme that is shock-capturing, and collisional. A classic test fluid codes undergo is the Sod shock tube test [Sod, 1978]. It is inspired by real life shock tubes in which a discontinuity is forced between the left and right halves of the tube. In a real shock tube the diaphragm between the two states is ruptured and the shocks are then observed. They're mostly used to observe material responses to shocks, however, we do have an analytical expectation of how the shock tube evolves through time.

The initial conditions for our left and right states in the CoMFi code are:

$$\begin{pmatrix} n_l \\ P_l \\ V_l \end{pmatrix} = \begin{pmatrix} 1.0 \\ 1.0 \\ 0.0 \end{pmatrix} \tag{58}$$

$$\begin{pmatrix} n_r \\ P_r \\ V_r \end{pmatrix} = \begin{pmatrix} 0.125 \\ 0.1 \\ 0.0 \end{pmatrix} \tag{59}$$

Figure III.3 shows the tube after time $t = 0.1$ in arbitrary units. In the density

graph it can be clearly seen the rarefaction wave, the contact discontinuity and the shock discontinuity. The exact sharpness is difficult to maintain in these codes but what is important is that it has not been too diffusive and the shocks are clear and maintaining their correct values. For the exact, analytical, solution see Sod [1978]. A resolution of 1001 grid points were used for this simulation. The boundary conditions on the right and left side are set equal to those initial values at all times. The energy equation was changed to the conservative internal energy equation to match Sod's equation set and what is used is the neutral component since it is acting more like a classical fluid and the Lorentz forces are irrelevant.

Soler et al. [2013] devised an initial value problem to test Alfvén waves in partially ionized plasmas. In the paper two-fluid plasma dynamics were explored in many ways. Analytical derivations and wave mode derivations were made. The initial value problem posed by Soler et al. [2013] is based on their fundamental standing mode of an Alfvén wave in the fully ionized case of $\omega L/V_A = \pi/20$. This means in the simulation domain of $z = (-10L, 10L)$ an initial value for the velocity perturbation is:

$$V_{\perp}(t = 0, z) = \cos\left(\frac{\pi}{20L}z\right) \quad (60)$$

The left and right boundary conditions in this case are set at $V_{\perp} = 0$ for both ions and neutrals. Constant and arbitrary for all other values. Three different orders of magnitude of the collision rate was used for two different cases. The first with a neutral initial value the same as the ions $V_{\perp} = U_{\perp}$ and the other where $U = 0$. The results in Figure III.4 show the results at $z = 0$ and the Alfvén wave can be seen clearly damped in the cases. Note how at $\nu_{ni} = 1$ the medium is moving in unison but still being damped. It is also important to note the equations used to solve this are fundamentally different than the ones in Soler et al. [2013]. They solve for the

primitive variables while CoMFi solves the conservative form. Still the graphs look similar (Figure III.5) which helps with the case of the validation of the code.

III.7 Visualization

CoMFi comes with its own visualization suite written in Python using Matplotlib [Hunter, 2007]. The data is saved in the HDF5 file format [The HDF Group, 2000-2010] which is compatible with many other visualization software, however the bundled `plot-scripts` folder contains the file `comfi.py` when imported gives tools to make plots and even animations.

III.7.1 Storage

CoMFi uses the HDF5 file format to store the data. The data hierarchy is shown in Table III.7.1. Other software like Origin, IDL or Tecplot may be easily used with this format. The data found in each variable is a matrix of the form, $u[x, nz - z]$, where nz is the number of grid cells in height. This was chosen so the matrix is displayed properly if shown in matrix form. Some software will require the matrix to be flipped when plotted.

III.7.2 Python Plot Scripts

In folder, `plot-scripts`, the python scripts required to make several plots are shown as examples. As well as examples required to make animations into `mp4` file format. This is done using Matplotlib and `h5py` for reading the HDF5 files. The Python header to import is `plot-scripts/comfi.py` and the functions are well documented using the PEP8 standard.

Table III.2: The HDF5 data hierarchy of the CoMFi code, this may be used in any visualization software with HDF5 support like IDL. All data is normalized. The **time-step** parent data name is referring to the time step of interest in the data. The **constant** name refers to the constant of interest with proper capitalization.

Time Vector	/t
Normalization Constants	/norm/[constant]
Ion Density	/[time-step]/Np
Neutral Density	/[time-step]/Nn
Ion Momentum (Horizontal)	/[time-step]/NVx
Ion Momentum (Vertical)	/[time-step]/NVz
Ion Momentum (Perpendicular)	/[time-step]/NVp
Neutral Momentum (Horizontal)	/[time-step]/NUx
Neutral Momentum (Vertical)	/[time-step]/NUz
Neutral Momentum (Perpendicular)	/[time-step]/NUp
Ion Temperature	/[time-step]/Tp
Neutral Temperature	/[time-step]/Tn
Magnetic Field (Horizontal)	/[time-step]/Bx
Magnetic Field (Vertical)	/[time-step]/Bz
Magnetic Field (Perpendicular)	/[time-step]/Bp
Electron Temperature (Optional)	/[time-step]/Te
Electron Pressure (Optional)	/[time-step]/Pe
Ion Energy (Optional)	/[time-step]/Ep
Neutral Energy (Optional)	/[time-step]/En




Figure III.3: Neutral fluid shock tube tests. The shocks are being captured appropriately due to the TVD-MUSCL scheme used.




Figure III.4: The Soler et al. [2013] initial value problem tests. Velocity is taken at $z = 0$ of the standing Alfvén wave. The solid red is the ion velocity, the dashed blue line is neutral velocity. The top row is where the initial neutral velocity is the same as the ion velocity and the bottom row is where the initial neutral velocity is at zero. The results resemble reasonably well the tests in the Soler et al. [2013] paper.




Figure III.5: Taken from Soler et al. [2013]. Compare with above (Figure III.4) for validation.

IV. CHROMOSPHERE DYNAMICS

The base of the solar atmosphere (right above the photosphere), referred to as the chromosphere, is highly dynamic and is strongly influenced by magnetic field activities [Hasan, 2008]. It is partially ionized, with as much as a thousand times more neutral Hydrogen than ions [Alissandrakis et al., 2018]. Song and Vasyliūnas [2011] and Leake et al. [2014] discussed in detail the effects of ion-neutral coupling in the chromosphere by considering neutrals and collisions into the momentum and Maxwell equations. A local-scale configuration commonly used in chromosphere models are so-called supergranules, convection cells of 30 Mm in the horizontal directions. Supergranules are separated by strong magnetic field regions, referred to as networks. It is believed that the network regions correspond to strong magnetic field at the photosphere altitude. Because the thermal pressure decreases with height while the total magnetic flux is conserved, the magnetic field tends to expand in higher altitudes to form a canopy or ‘wine glass’ shaped magnetic geometry so that total pressure, the magnetic pressure, p_B , plus thermal pressure, p_T , i.e., $p_T + p_B$, is balanced at each height [Gabriel, 1976]. This geometry is reflected in a significant altitudinal change in the plasma beta, which is the ratio of the thermal to magnetic pressures, $\beta = p_T/p_B$. The chromosphere has much lower temperature than that of the hot corona. The average temperature of the Chromosphere is around 6000 K while the coronal temperature is over million degrees [Avrett and Loeser, 2008b]. This dramatic change in

the temperature, taking place in a very narrow region of a few hundred kilometers referred to as the transition region, known as the coronal heating problem and it is a long-standing problem in solar physics.

In the chromosphere shocks often observed and jets are commonplace in the form of spicules. Spicules are local, hot upward jets of plasma observed in the corona. The origination and mechanism behind spicule formation is not well known. Spicules are categorized into two types [de Pontieu et al., 2007]. Type I spicules have longer lifetimes and are typically larger (5 min and ~ 15 Mm). Type II spicules have shorter lifetimes and are smaller (≤ 2 min and ~ 6 Mm). De Pontieu et al. [2004] suggested that photospheric oscillations give birth to spicules. It was also suggested that magnetic reconnection may be the cause of Type II spicules [de Pontieu et al., 2007].

There is strong observational evidence for the ubiquity of Alfvén waves in the sun’s atmosphere [Mathioudakis et al., 2013, McIntosh et al., 2011, Jess et al., 2009, Ofman and Wang, 2008, Aschwanden et al., 1999, Nakariakov et al., 1999]. These observations of the solar limbs show Alfvén waves with amplitudes in the range of 5-25 km/s and phase speeds 250-500 km/s. de Pontieu et al. [2007] have shown that Alfvén waves are carried by spicules with significant amplitudes of order 20 km s^{-1} . It has been suggested that Alfvén waves could be damped in the chromosphere due to collisions between plasma ions and neutrals and causes sufficient heating [Song and Vasyliūnas, 2011] and [Song and Vasyliūnas, 2014]. Song [2017] further proposed that the nonuniform heating associated with the Alfvén waves may explain the formation of the spicules. Khomenko [2017] looks at other waves as well as Alfvén waves and their effects in weakly ionized solar plasma.

Recent simulations and models of the chromosphere have adopted a single-fluid approach [Kuzma et al., 2017, Ni et al., 2016, Brady and Arber, 2016, Gudiksen et al., 2011]. In order to account for neutrals’ effect on the plasma and electric field

an ‘ambipolar diffusion’ term is added to the Generalized Ohm’s Law to account for collisions [Martinez-Sykora et al., 2012, 2017]. This is useful to study the highly magnetized, low β , environment of the upper chromosphere. A two-fluid approach takes into account separate fluid motions of plasma and neutrals especially in highly collisional environments like the low solar atmosphere in regions of high magnetic gradients. Soler et al. [2013] studied Alfvén wave propagation in partially ionized plasma while taking a generalized approach with free parameters with the lower solar atmosphere case as an example. Maneva et al. [2017] and Wójcik et al. [2018] studied acoustic wave propagation using a collisional two-fluid approach as well. Three-fluid simulations, with electrons being the third fluid, were made to study reconnection events in the chromosphere [Leake et al., 2013]. In these highly localized small regions electron motion is separate from ion and neutral motion and its effects cannot be neglected. The previously mentioned work uses the *Bifrost* [Gudiksen et al., 2011] code which employs a very large artificial viscosity for stability. By self-consistently solving plasma, neutral fluids, and Maxwell’s equations, Tu and Song [2013b] have shown that due to Joule dissipation Alfvén waves can provide strong enough chromospheric heating and that the heating rate is consistent with the theoretical model of Song and Vasyliūnas [2011].

IV.1 Governing Equations

In the model presented here, a two-fluid, collisional approach is used to account for the friction between ions and neutrals. The first fluid, the ion fluid, is similar to the traditional, single-fluid plasma, and assumes charge quasi-neutrality. With the Maxwell’s equations, this fluid is used to calculate self-consistently the evolution of the plasma motion, magnetic fields and currents in the system. The second, neutral fluid, accounts for the neutral particles in the system. The two fluids interact by

collisions. The friction between ions and neutrals affects the neutral fluid motion so that the neutrals experience the electromagnetic effects.

This is similar to the three-fluid HiFi code used for studying reconnection in the chromosphere [Leake et al., 2012]. Except electrons are not treated as a separated fluid in this model because the electron's mass is more than three orders of magnitude smaller than ion or neutral particle mass, so its contribution to heating and the equations considering the scales is negligible. The effects of its motion are included in the current terms. We use empirical number density model that comes from observations assuming steady-states in the chromosphere [Avrett and Loeser, 2008b]. Here, I do not account for ionization and recombination which I plan to implement photo-chemistry in future work.

Our model uses Cartesian, 2.5 dimensionality, which means that all the quantities are functions of horizontal (x) and vertical (z) directions but uniform in the y -direction, while vector quantities have all three components including the third component that is perpendicular (\perp) to the $x - z$ plane. The perpendicular components are used to generate Alfvén waves at the photospheric boundary.

The governing equations are adapted from Tu and Song [2016] for the two-fluid (plasma and neutral) system as follows:

$$\frac{\partial \rho_i}{\partial t} + \nabla \cdot \rho_i \mathbf{V} = 0 \quad (1)$$

$$\frac{\partial \rho_n}{\partial t} + \nabla \cdot \rho_n \mathbf{U} = 0 \quad (2)$$

$$\frac{\partial \rho_i \mathbf{V}}{\partial t} = -\nabla \cdot [\rho_i \mathbf{V} \mathbf{V} + n_i k_b T_i + \frac{B^2}{2\mu_0} \mathbf{I} - \frac{\mathbf{B} \mathbf{B}}{\mu_0}] + \rho_i \mathbf{g} - \rho_i \nu_{in} (\mathbf{V} - \mathbf{U}) \quad (3)$$

$$\frac{\partial \rho_n \mathbf{U}}{\partial t} = -\nabla \cdot [\rho_n \mathbf{U} \mathbf{U} + n_n k_b T_n] + \rho_n \mathbf{g} + \rho_i \nu_{in} (\mathbf{V} - \mathbf{U}) \quad (4)$$

$$\frac{\partial \mathbf{B}}{\partial t} = -\nabla \cdot (\mathbf{V} \mathbf{B} - \mathbf{B} \mathbf{V}) - \nabla \times (\eta \mathbf{J} + \frac{1}{n_e e} \mathbf{J} \times \mathbf{B}) \quad (5)$$

$$\begin{aligned} \frac{\partial k_b T_i}{\partial t} = & -\nabla \cdot (k_b T_i \mathbf{V}) + \frac{1}{3} k_b T_i \nabla \cdot \mathbf{V} - \frac{2}{3} \frac{\nabla \cdot (\mathbf{q}_i + \eta \mathbf{B} \times \mathbf{J})}{n_i} \\ & + \frac{2}{3} \frac{m_i}{m_i + m_n} \nu_{in} [3k_b (T_n - T_i) + m_n |\mathbf{V} - \mathbf{U}|^2] \end{aligned} \quad (6)$$

$$\begin{aligned} \frac{\partial k_b T_n}{\partial t} = & -\nabla \cdot (k_b T_n \mathbf{U}) + \frac{1}{3} k_b T_n \nabla \cdot \mathbf{U} - \frac{2}{3} \frac{\nabla \cdot \mathbf{q}_n}{n_n} \\ & + \frac{2}{3} \frac{m_n}{m_i + m_n} \nu_{in} [3k_b (T_i - T_n) + m_i |\mathbf{V} - \mathbf{U}|^2] \end{aligned} \quad (7)$$

$$\nabla \times \mathbf{B} = \mu_0 \mathbf{J} \quad (8)$$

$$\mathbf{q} = -\kappa \nabla T \quad (9)$$

Here i , e and n subscripts denote proton, electron and neutral, respectively. The primitive scalars m , ρ , T are particle mass, mass density and temperature, respectively. \mathbf{V} , \mathbf{U} , and \mathbf{B} are ion velocity, neutral velocity, and the magnetic field, respectively, with \mathbf{g} denoting the solar gravitational acceleration near the solar surface. κ is the Spitzer conductivity. ν_{ij} denotes collision rate of species i to species j , and $\eta = m_e(\nu_{en} + \nu_{ei})/e^2 n_e$ is the ohmic resistivity. We use the following collision

rates [see Pontieu et al., 2001, Appendix A]

$$\begin{aligned}
 \nu_{in} &= 7.4 \times 10^{-11} n_n T^{1/2} \\
 \nu_{en} &= 1.95 \times 10^{-10} n_n T^{1/2} \\
 \nu_{ei} &= 3.759 \times 10^{-6} n_e T_e^{-3/2} \ln \Lambda
 \end{aligned} \tag{10}$$

where $\ln \Lambda$ is the Coulomb logarithm taken as 10 for simplicity and n is in cm^{-3} and $T = \frac{T_i + T_n}{2}$ in K.

Comparing this with the two-fluid equations seen in Section II.1.5 we see that we've taken the more extreme case of $m_e \sim 0$. This is for the first-order approximation that electrons contribute less to heating than hydrogen ions and neutrals. The CoMFi code is capable of reverting back to the approximation, $\frac{m_e}{m_i} \sim 0$, for cases where the current density terms matter like in mid-chromosphere reconnection.

IV.2 Simulation Domain

In the horizontal dimension, I simulate a supergranule similar to that described in Song [2017]. Because of symmetry, our simulation domain contains half of a supergranule, from the center of a network with strong field to the center of internetwork at the photospheric boundary. If I further assume that the neighboring supergranules have the same polarity as the one in the simulation domain, there is no reconnection between the supergranules of interest. In this case the normal component of the field, B_x , is zero at the left and right boundaries. The right and left boundaries act as mirrors to reflect any normal component of the field. Similarly, the normal component of the velocity is also set to zero at the right and left boundaries, i.e. there is no horizontal flow from one supergranule to another at the chromospheric altitude although inter-supergranule flow is possible in the corona height and in or

below the photosphere.

Since I only simulate the chromosphere and treat the transition region above and photosphere below as upper and lower boundaries, respectively. The photosphere below is made up as a Dirichlet boundary conditions below the surface of the photosphere, the sun is highly dynamic. The assumption made in this mode is the Alfvén waves are generated at the photosphere and propagate while undergoing damping towards the transition region. For this reason to imitate the energy flux and activity of the photosphere, a driver at the bottom of the simulation domain is made as a Dirichlet boundary condition. Using the broadband spectrum driving method of Tu and Song [2013b], the perpendicular velocity is a continuous oscillation with a broadband time series of the form:

$$V_{\perp}(t, z = 0) = V_b \left[\sum_{k=0}^s (\omega_k / \omega_p)^{5/6} \cos(\omega_k t + \Phi_k) + \sum_{j=s+1}^m (\omega_p / \omega_j)^{5/6} \cos(\omega_j t + \Phi_j) \right] \quad (11)$$

where V_{\perp} will be the imposed speed on the lower boundary in the direction normal to the $x - z$ plane. V_b is the amplitude of the broadband spectrum, 2 km/s, to ensure an appropriate photospheric flux of $2 \times 10^7 \text{ erg cm}^{-2} \text{ s}^{-1}$ [see Tu and Song, 2013b, section 3.2], ω_j and ω_k are angular frequencies filled logarithmically, ω_p is the peak angular frequency at 1.65 mHz. The driver also has random phases Φ_j, Φ_k which is a multiple of $2\pi \times$ by random number between 0 and $s+m$. For the case of this simulation $s=100$ and $m=500$.

The number density of the ions and neutrals are specified at the boundary as the photospheric values from Avrett and Loeser [2008b] of 10^{20} m^{-3} and 10^{23} m^{-3} , respectively. As an initial condition, a stratified hydrostatic profile is made for the ion densities $n(z) = n(z = 0)e^{-m_i g z / k T_0}$ and a more realistic neutral number

density profile is filled based on the above mentioned model. Here $T_0 = 7000$ K which is the temperature imposed on the photosphere and also the initial condition for both ionized plasma and neutrals in the entire domain. An isothermal approach was chosen initially because the chromosphere's temperature does not change much, between 6000K to 8000 K, between the photosphere and the transition region. At later times, the temperatures are determined from the energy equations 6 and 7.

A requirement in some MHD codes with zero magnetic field divergence constraints is that the initial condition must also be divergence free. For this reason $B_x(t = 0)$ and $B_\perp(t = 0)$ is chosen to be zero while $B_z(x, t = 0)$ varies in the horizontal x direction. It consists of regions of strong field, $\sim 1\text{kG}$, for the network region and weak field, ~ 50 G, for the internetwork region with a sharp linear drop in between. The 1 kG region only takes up 0.05 Mm in the horizontal direction and is in the left part of the domain. The weak region part of the rest of the domain. The goal is to allow this initial condition to quickly relax to a canopy because of the large β change due to height without solving a force-free field. At the bottom boundary, a Dirichlet condition is applied to $B_z(x)$ but for the other two components the field is set Neumann conditions $\frac{\partial B_{z,\perp}}{\partial z}\big|_{z=0\text{km}} = 0$. This configuration resembles the stem of the canopy and the magnitude of the supergranule strengths at the photosphere. The bottom boundary's $B_z(x)$ is an unchanging Dirichlet condition to mimic the convection cell underneath.

At the top boundary which represents the transition region and the corona above it, the model uses Neumann conditions to allow free-flow of the plasma through it (i.e. $\frac{\partial \mathbf{u}}{\partial z}\big|_{z=2100\text{km}} = 0$).

In the simulations and results shown in this paper the domain is 15 Mm horizontally (half the size of the magnetic internetwork) and 2.1 Mm vertically (typical thickness of the chromosphere). It is a uniform Cartesian grid of 256×128 where

$\Delta x = 58.6$ km and $\Delta z = 16.4$ km. This is the largest resolution chosen for speed and memory considering the usage of a dedicated scientific computing GPU (see Chapter III). This is larger than the length scale of the collisionally reduced Alfvén wavelength ($\frac{V_A}{\nu_{in}}$) which means for this simulation the plasma and neutrals are coupled. Since this is a 2.5D simulation all spatial derivatives in the perpendicular direction are taken as zero ($\frac{\partial}{\partial \perp} = 0$).

All momentum components are set initially as zero.

IV.3 Results

Figure IV.1 shows a snapshot on the chromospheric solution after 348 *ms*. It shows that our simulation produces magnetic field topology that is characterized by canopy or ‘wine glass’ shape [Rutten, 2007], with a strong field of about 1 *kG* at the stem on the left, and a weaker field of about 50 *G* at the near the upper boundary of the simulation domain, near the transition region. This topology is obtained fairly quickly, in around ~ 350 ms, because of the magnetic pressure imbalance imposed in the initial conditions. This is not a relaxed system as the canopy holds for a few seconds before an eruption starts forming.

Figure IV.2 shows a formation of what resembles a jet — a parcel of plasma quickly moving upward towards the transition region. The shock spontaneously forms after around 20s and the jet speeds up and shoots up outside of the chromosphere into the corona in about 5s.

In Figure IV.3, I show the distribution of the two heating terms—the ohmic heating, ηJ^2 , and the frictional heating, $\nu_{in} \rho_i |\mathbf{V} - \mathbf{U}|^2$, in the simulation domain, right before the eruption. The heating rates are normalized per neutral particle. It can be seen that the frictional heating dominates the ohmic heating by orders of magnitude, especially in the strong field region where the shock forms.




Figure IV.1: Formation of a canopy seen after around 348ms (13000 time steps). The initial condition of a concentrated strong field region (1 kG) and weak field region (5G) creates vertical asymmetry as plasma beta drastically decreases. The left and right boundaries are mirrored so effects of other granules behaving the same way are considered. Because of the short time period this is *not* a relaxed system but simply a consequence of the pressure imbalance frequently seen in the chromosphere.




Figure IV.2: A zoom into the shock wave front of the jet. The arrows represents the direction and magnitude of ion flow. The color bar represents the neutral number density in cm^{-3}

Figure IV.4 shows the neutral temperature, the frictional heating, and the velocity difference, $|\mathbf{V} - \mathbf{U}|^2$ during the emergence of the jet to the corona (at $t =$




Figure IV.3: Heating of the shock origin before the formation of the jet at $t=18.9s$. Top image is the frictional heating per neutral particle. Bottom image is the ohmic heating per neutral particle. The frictional heating is orders of magnitude larger than the heating due to currents.

26.8 s). The plots show the formation of an appreciable shock which separates a low number density, high temperature and strong heating region with a high number density, lower temperature and weaker heating with the rest of the medium. The figure also shows a very clear difference in heating rates on both sides of the shock, where the front of the shock travels at super sonic speeds. The shock propagates supersonically until the plasma is ejected when the chromosphere is depleted and starts to fill up again with flows going back down. Downstream of the shock, the plasma flow is returning due to a cavity left behind, where upstream of the shock, the chromospheric plasma is still unchanged.

Figure IV.5 shows a time series of the evolution of the shock number density and magnetic field lines. It can be seen that the field lines follow the shock evolution, which means magnetic field topology (i.e., reconnection region) is not a driver of the jet. Figure IV.6 shows the same time series but with contours of frictional heating




Figure IV.4: These figures show the jet in the middle of its journey to the corona at $t=26.8s$. From top to bottom, the temperature of the neutral plasma, the frictional heating per neutral particle, and the difference in speeds $\|\mathbf{V} - \mathbf{U}\|^2$ of the charged and neutral plasma. The shock can be clearly seen in the bottom-left corner of the simulation where the magnetic flux concentration is strongest. Due to large plasma beta a ‘ring’ of slow moving plasma is formed and persists in the shock allowing the shock to start forming.

and flow streamlines. It clearly shows that the jet is driven by frictional joule heating.

Figures IV.7, IV.8, IV.9 and IV.10 show one-dimensional extractions of the solution during the shock emergence along a vertical line at $x = 0.6 \text{ Mm}$. These plots exemplify the cause and effect of the shock. The first plot shows the time series of the wavefront of the shock climbing in the chromosphere. The second and third show that the frictional heating jump follows the shock, and that the magnetic field




Figure IV.5: Time series of the shock. From left to right and top to bottom the times are $t=26.751\text{s}$, 26.785s , 26.831s , 26.842s , 26.844s , 26.849s . The white streamlines denote the magnetic field lines and the number density of the neutrals are in cm^{-3} . This shows them magnetic field geometry changing due to the jet and not the other way round. The jet is not generated because of the magnetic field.

changes correspond to the jet behind the shock. The last plot shows a velocity profile of the jet.

IV.4 Discussion

A magnetic canopy forms due to the magnetic and thermal pressure imbalance of the strong field regions from the photosphere and the weak field but high temper-




Figure IV.6: Time series of frictional heating in the jet. From left to right and top to bottom the times are $t=26.751\text{s}$, 26.785s , 26.831s , 26.842s , 26.844s , 26.849s . The white streamlines denote flow.

ature regions scaling up to the transition region. This is consistent with a two-fluid analytic model that was presented in Song [2017]. It shows that the model can provide numerical solutions for chromospheric conditions. However, the magnetic canopy does not relax or reach a steady-state due to the nature of the collisional model and not in the timescales of the current simulation. The super granule canopy may not last because the photospheric power flux, generated at the lower boundary, continuously adds energy into the system. Small scale processes or highly dynamic events like jets override the canopy magnetic field because of large plasma speeds. Our results show




Figure IV.7: Ion number density snapshots of shock wave front at line of extraction $x=0.6\text{Mm}$. Ion number density is in units of 10^{11} cm^{-3} . The wavefront can be clearly seen climbing the Chromosphere.




Figure IV.8: Blue line is ion number density and the black and gray dashed lines are B_x and B_z respectively at $t=26.83\text{s}$ and vertical line at $x=0.6\text{Mm}$. Ion number density is in units of 10^{11} cm^{-3} . The magnetic field is changed behind the shock wave front.




Figure IV.9: Blue line is ion number density and red dashed line is frictional heating per neutral particle at $t=26.83s$ and vertical line $x=0.6Mm$. Ion number density is in units of 10^{11} cm^{-3} . The frictional heating per neutral particle behind the wavefront is orders of magnitude higher than in front of it.

that the jet has immediate effects on the magnetic field due to induction and not vice versa.

Largest Alfvén speeds seen in the simulation is 11 Mm/s which means that in the 30s of simulation, magnetic effects can travel through the domain at least 3 times with constant density and even less in a hydrostatic density. Alfvén waves generated as the drivers for frictional heating at the bottom boundary do not reflect at the top boundary enough times to warrant a time-averaged study of the entire heating of the chromosphere nor does it have enough energy to drive the jet of the same kind as presented. The presence of the shock also inhibits any long-term ideal studies of the atmosphere since it depletes and cavities form with the jet. In a future study, I will see how Alfvén waves affect the chromosphere with force-free initial conditions so as




Figure IV.10: Blue line is ion number density and black dashed line is vertical ion velocity at $t=26.83s$ and vertical line $x=0.6Mm$. Ion number density is in units of 10^{11} cm^{-3} . The jet speed is supersonic.

to not form any jets that can destroy a canopy configuration.

Although the frictional heating is large in the entire magnetic network column, an instance of no frictional heating creates a separation of upstream and downstream fluids preceding the shock. This can be seen in Figure IV.3 and IV.4 and is due to a high temperatures and density which causes the two fluids to move in unison and therefore $\rho_{in}\nu_{in}\|V - U\| = 0$. The cause of the frictional heating is not Alfvén wave damping but the imbalanced initial conditions and the horizontal $\mathbf{J} \times \mathbf{B}$ force. The lower medium heats faster than above it and this creates the shock in the lower atmosphere that then propagates due to thermal pressure.

It has been suggested that the driver of these kinds of spicule-like events is magnetic reconnection which originates from the vicinity of a large magnetic flux concentration in the network [de Pontieu et al., 2007]. In our simulation, the frictional

heating heats a fluid downstream of the unison speed ring larger than that of upstream causing a shock wave as expected from a Type I spicule. However the simulation does not show reconnection effects with enough energy to drive a spicule. Although MHD simulations do not mimic reconnection directly, currents present in reconnection events would accelerate electrons into a collision due to the dense and hot medium surrounding it. Therefore, magnetic reconnection is directly related to ohmic heating in the chromosphere. If comparing the dominating heating terms, frictional heating dominates ohmic heating which means *collisions are causing these jets to form and not reconnection-like currents*. The size and extreme temperatures of the jet in the simulation closely resemble chromospheric jets [Shimojo and Shibata, 2000, Nishizuka et al., 2011].

In Figure IV.7 the jet can be seen accumulating a shock of number density 10^{11} cm^{-3} . This density is consistent with IRIS observations of spicules in the corona [Alexandakis et al., 2018]. The simulation does show that changes in the magnetic field happen behind the shock wave front's direction of motion. Figure IV.8 implies the change in the magnetic field is a consequence of the jet and the rising magnetic field density in the plane of the simulation is due to compression from the shock itself. The shock in the frictional heating term, $\frac{\rho_i \nu_{in} \|\mathbf{V} - \mathbf{U}\|^2}{n_n}$, seen in Figure IV.9 coincides exactly at the shock showing enhanced heating behind the shock front. The cause of the shock climb is the thermal pressure behind and in front of the shock.

The jet's temperatures and speed are much larger than observed. This is mostly due to the fact that in my model I have not included the radiative loss effect in the energy equations. The larger than reality shock heating because of the greater flow speed may also further amplify the temperature. The over-estimated temperature of the downstream medium may also lead to estimate of the collision frequencies. This growing downstream temperature has a compounding effect on the thermal

pressure gradient downstream and upstream and causes the speeds to go supersonic. Further studies will incorporate the photo-chemistry and also radiative losses. These implementations will help to see how they affect shock formation and the energy balance inside the chromosphere. It is also important to note that this shock travels at super-Alfvénic speeds so it is a super-Alfvénic jet.

Kuźma et al. [2017] made a numerical simulation of a solar spicule by imposing a vertical pulse in the solar limb. In contrast, the shock shown in our simulation self-consistently evolves without forcing a push on the plasma and the jet is a consequence of thermal effects. Thus, it provides a complete picture of the chromospheric heating conditions for shock formation. The spicule speeds in Kuźma et al. [2017] match more closely to the observed 25 km s^{-1} than our simulation, but that is expected in the case where the pulse is controlled and the simulation is single-fluid. Kuźma et al. [2017] also investigated adiabatic and non-adiabatic studies of spicules and the wave structures found in the plasma. The jets in our simulation are different than the imposed pulses of Kuźma et al. [2017] since that is a cold dense plasma that has been imposed, while in our simulation, it is a hot large plasma jet that evolved self-consistently.

Song and Vasyliūnas [2014] found that by subtracting the two fluids' momentum equations the evolution of the speed differences, $d/dt|\mathbf{V} - \mathbf{U}|$, which leads to frictional heating is caused and maintained by magnetic pressure and thermal pressure differences between the fluids. Tu and Song [2013b], who studied a time-averaged heating in the chromosphere, found that ohmic heating dominated the lower chromosphere and frictional heating dominated the upper chromosphere which adequately explained the heating of the chromosphere. This picture seems to be different from the results of our simulation as frictional heating dominated ohmic heating in the lower chromosphere but this is most likely due to the field strength used in the two

studies. Their study limited the field in less than 100 G, but our model includes region of 1 kG field. The one order of magnitude higher in the field decreases the ohmic-to-frictional heating ratio by two orders of magnitude.

We use our model to demonstrate the importance of collisions in the chromosphere. A model without collisions is incomplete as neutral particles dominates the charged particles in this small layer of the sun. By simply adding collision terms I show that the simulation replicates chromospheric conditions well. Such a model can be used as an interface between the (fully) neutral photosphere and the (fully) ionized corona. Hillier, A. et al. investigated 1D slow-mode shocks in chromospheric conditions and how magnetic energy can be converted to frictional heating in these cases.

Finally, this work serves as a step towards a better understanding of the solar chromosphere. The *Parker Solar Probe* mission [Fox et al., 2016] is expected to shed light on the dominant processes in the solar atmosphere, including the chromosphere, and these processes will be implemented in the model.

V. CONCLUSION

MHD collisional modeling, chromospheric modeling, scientific computing, and introducing the CoMFi code were the focus of this dissertation. Typically in the world of physics, the uptake of new ideas in science is not without inertia. This is understandable as I spend our time focusing on the physics above all else. If I take Moore's Law as a metaphor, I should understand that eventually computers will be able to do whatever I please it to and relying on legacy codes will be a thing of the past. An example of this is the Python programming language. A couple decades ago I could not prototype the kind of software I am able to nowadays by the invention of this programming language. Analogously, decades from now I should be able to solve differential equations by taking new computer science inventions and hopefully with less resistance.

We present a self-consistent, two-fluid MHD model to study the chromosphere. The model includes ion-neutral interaction, which allows us to investigate the role of frictional heating in the dynamics of the chromosphere. The very high collision rates in the partially ionized chromosphere stabilizes the code so that no numerical viscosity is needed.

The model obtains preliminary results which require further investigation. Our simulation presents that, between network and internetwork regions, a jet could potentially be driven purely by thermal processes, especially the frictional heating,

and not by magnetic reconnection. Thus, it is crucial to include the neutrals in any attempt to model the lower chromosphere.

The discrepancy between observational speeds and speeds in the simulation could be explained by the need to include the effects of photo chemical reactions in the chromosphere and the radiative losses. Further refinement of the model with photo-chemical effects, radiation and new observations from the upcoming *Parker Solar Probe* mission or the Daniel K. Inouye Solar Telescope (DKIST) [Tritschler et al., 2015] will lead to an improved model of the chromosphere.

It is my hope that the CoMFi code can continue to provide good physics in the future and hopefully adapted to suit the needs of the user. GPU clusters are an attractive new frontier in scientific computing and its thousands of cores per unit is exactly suited to solve our differential equations. As GPUs become cheaper our codes must be sure to utilize them.

V.1 Future Work

Extending the CoMFi code further to be used on GPU clusters is a high priority. Many studies can be made if higher resolutions are resolved. Typically memory becomes an issue in GPUs and it can only compute as much as it has. Chromosphere physics are very similar to ionosphere physics in that they are both multi-fluid. This means adding a fluid generator into the code can help with multi-fluid cases which increases the utility of the code.

Different chromosphere events would be interesting to simulate. Although the focus of the previous study was the jet formed due to frictional heating, the code is capable of simulating magnetic reconnection events in the chromosphere. A follow-up approach for the jet that formed due to frictional heating in the previous study would be to increase the resolution and only simulate the lower chromosphere and be sure

to include force-free initial conditions (except with magnetic pressure) to study in more detail why the jet is forming. In addition, a parametrized study can be made to see how much magnetic energy is required to convert into the jet by changing the magnetic field gradients.

LITERATURE CITED

- Preface to the second edition. In Charles Hirsch, editor, *Numerical Computation of Internal and External Flows (Second Edition)*, pages xv – xvii. Butterworth-Heinemann, Oxford, second edition edition, 2007. ISBN 978-0-7506-6594-0. doi: <https://doi.org/10.1016/B978-075066594-0/50037-0>. URL <http://www.sciencedirect.com/science/article/pii/B9780750665940500370>.
- Qusai Al Shidi. CoMFi - Collisional Multi-Fluid ion code. <https://github.com/qalshidi/comfi/>, 2019.
- Qusai Al Shidi, Ofer Cohen, Paul Song, and Jiannan Tu. Time-Dependent Two-Fluid Magnetohydrodynamic Model and Simulation of the Chromosphere. *arXiv e-prints*, art. arXiv:1904.01572, Apr 2019.
- C. E. Alissandrakis, J. C. Vial, A. Koukras, E. Buchlin, and M. Chane-Yook. IRIS Observations of Spicules and Structures Near the Solar Limb. , 293:20, February 2018. doi: 10.1007/s11207-018-1242-4.
- M. J. Aschwanden, L. Fletcher, C. J. Schrijver, and D. Alexander. Coronal Loop Oscillations Observed with the Transition Region and Coronal Explorer. , 520: 880–894, August 1999. doi: 10.1086/307502.
- Eugene H. Avrett and Rudolf Loeser. Models of the solar chromosphere and transition region from sumer and hrts observations: Formation of the extreme-ultraviolet spectrum of hydrogen, carbon, and oxygen. *The Astrophysical Journal Supplement Series*, 175(1):229, 2008a. URL <http://stacks.iop.org/0067-0049/175/i=1/a=229>.
- Eugene H. Avrett and Rudolf Loeser. Models of the solar chromosphere and transition region from sumer and hrts observations: Formation of the extreme-ultraviolet spectrum of hydrogen, carbon, and oxygen. *The Astrophysical Journal Supplement Series*, 175(1):229, 2008b. URL <http://stacks.iop.org/0067-0049/175/i=1/a=229>.

- D. N. Baker, X. Li, A. Pulkkinen, C. M. Ngwira, M. L. Mays, A. B. Galvin, and K. D. C. Simunac. A major solar eruptive event in july 2012: Defining extreme space weather scenarios. *Space Weather*, 11(10):585–591, 2013. doi: 10.1002/swe.20097. URL <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/swe.20097>.
- P.M. Banks and G. Kockarts. Chapter 22 - thermal processes of the ionosphere. In P.M. Banks and G. Kockarts, editors, *Aeronomy*, pages 238 – 286. Academic Press, 1973. ISBN 978-0-12-077802-7. doi: <https://doi.org/10.1016/B978-0-12-077802-7.50014-2>. URL <http://www.sciencedirect.com/science/article/pii/B9780120778027500142>.
- W. Baumjohann and R.A. Treumann. *Basic Space Plasma Physics*. Imperial College Press, 1997. ISBN 9781860940798. URL <https://books.google.com/books?id=bbVACbm9FjYC>.
- P. L. Bhatnagar, E. P. Gross, and M. Krook. A model for collision processes in gases. i. small amplitude processes in charged and neutral one-component systems. *Phys. Rev.*, 94:511–525, May 1954. doi: 10.1103/PhysRev.94.511. URL <https://link.aps.org/doi/10.1103/PhysRev.94.511>.
- C. S. Brady and T. D. Arber. Simulations of alfvén and kink wave driving of the solar chromosphere: Efficient heating and spicule launching. *The Astrophysical Journal*, 829(2):80, 2016. URL <http://stacks.iop.org/0004-637X/829/i=2/a=80>.
- J. G. CHARNEY, R. FJÖRTOFT, and J. Von NEUMANN. Numerical integration of the barotropic vorticity equation. *Tellus*, 2(4):237–254. doi: 10.1111/j.2153-3490.1950.tb00336.x. URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.2153-3490.1950.tb00336.x>.
- B. De Pontieu, R. Erdelyi, I. De Moortel, and T. Metcalf. Photospheric Oscillations in the Solar Atmosphere: Driving Chromospheric Spicules and Coronal Waves. *AGU Fall Meeting Abstracts*, art. SH13A-1142, December 2004.
- B. de Pontieu, S. McIntosh, V. H. Hansteen, M. Carlsson, C. J. Schrijver, T. D. Tarbell, A. M. Title, R. A. Shine, Y. Suematsu, S. Tsuneta, Y. Katsukawa, K. Ichimoto, T. Shimizu, and S. Nagata. A Tale of Two Spicules: The Impact of Spicules on the Magnetic Chromosphere. , 59:S655–S662, November 2007. doi: 10.1093/pasj/59.sp3.S655.
- A. Dedner, F. Kemm, D. Kröner, C.-D. Munz, T. Schnitzer, and M. Wesenberg. Hyperbolic Divergence Cleaning for the MHD Equations. *Journal of Computational Physics*, 175:645–673, January 2002. doi: 10.1006/jcph.2001.6961.

- C. C. Finlay, B. Langlais, F. J. Lowes, M. Manda, M. Menvielle, L. Tøffner-Clausen, N. Olsen, A. Tangborn, Z. Wei, C. Manoj, S. Maus, S. McLean, A. W. P. Thomson, B. Hamilton, C. D. Beggan, S. Macmillan, T. A. Chernova, T. I. Zvereva, T. N. Bondar, V. P. Golovkov, A. Chambodut, A. Chulliat, E. Thébaud, G. Hulot, H. Lühr, I. Michaelis, I. Wardinski, J. Rauberg, M. Hamoudi, M. Rother, V. Lesur, R. Holme, T. J. Sabaka, and W. Kuang. International Geomagnetic Reference Field: the eleventh generation. *Geophysical Journal International*, 183(3):1216–1230, 12 2010. ISSN 0956-540X. doi: 10.1111/j.1365-246X.2010.04804.x. URL <https://dx.doi.org/10.1111/j.1365-246X.2010.04804.x>.
- N. J. Fox, M. C. Velli, S. D. Bale, R. Decker, A. Driesman, R. A. Howard, J. C. Kasper, J. Kinnison, M. Kusterer, D. Lario, M. K. Lockwood, D. J. McComas, N. E. Raouafi, and A. Szabo. The Solar Probe Plus Mission: Humanity’s First Visit to Our Star. , 204:7–48, December 2016. doi: 10.1007/s11214-015-0211-6.
- A. H. Gabriel. A magnetic model of the solar transition region. *Philosophical Transactions of the Royal Society of London Series A*, 281:339–352, May 1976. doi: 10.1098/rsta.1976.0031.
- B. V. Gudiksen, M. Carlsson, V. H. Hansteen, W. Hayek, J. Leenaarts, and J. Martínez-Sykora. The stellar atmosphere simulation code Bifrost. Code description and validation. , 531:A154, July 2011. doi: 10.1051/0004-6361/201116520.
- Ami Harten. High resolution schemes for hyperbolic conservation laws. *Journal of Computational Physics*, 135(2):260 – 278, 1997. ISSN 0021-9991. doi: <https://doi.org/10.1006/jcph.1997.5713>. URL <http://www.sciencedirect.com/science/article/pii/S0021999197957132>.
- S. S. Hasan. Chromospheric dynamics. *Advances in Space Research*, 42:86–95, July 2008. doi: 10.1016/j.asr.2007.08.019.
- Hillier, A., Takasao, S., and Nakamura, N. The formation and evolution of reconnection-driven, slow-mode shocks in a partially ionised plasma. doi: 10.1051/0004-6361/201628215.
- J. D. Hunter. Matplotlib: A 2d graphics environment. *Computing In Science & Engineering*, 9(3):90–95, 2007. doi: 10.1109/MCSE.2007.55.
- D. B. Jess, M. Mathioudakis, R. Erdélyi, P. J. Crockett, F. P. Keenan, and D. J. Christian. Alfvén Waves in the Lower Solar Atmosphere. *Science*, 323:1582, March 2009. doi: 10.1126/science.1168680.
- Florian Rudolf Josef Weinbub Andreas Morhammer Tibor Grasser Ansgar Jüngel Karl Rupp, Philippe Tillet and Siegfried Selberherr. Viennacl - linear algebra library for multi- and many-core architectures, 2016. URL <http://viennacl.sourceforge.net/>.

- Elena Khomenko. On the effects of ion-neutral interactions in solar plasmas. *Plasma Physics and Controlled Fusion*, 59(1):014038, 2017. URL <http://stacks.iop.org/0741-3335/59/i=1/a=014038>.
- Delores J. Knipp. Fall 2018 agu editors' highlights: Living within the sun's stormy atmosphere. *Space Weather*, 17(1):3–5, 2019. doi: 10.1029/2019SW002154. URL <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019SW002154>.
- Delores J. Knipp, Brian J. Fraser, M. A. Shea, and D. F. Smart. On the little-known consequences of the 4 august 1972 ultra-fast coronal mass ejecta: Facts, commentary, and call to action. *Space Weather*, 16(11):1635–1643, 2018. doi: 10.1029/2018SW002024. URL <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018SW002024>.
- Alexander Kurganov and Eitan Tadmor. New high-resolution central schemes for nonlinear conservation laws and convection–diffusion equations. *Journal of Computational Physics*, 160(1):241 – 282, 2000. ISSN 0021-9991. doi: <https://doi.org/10.1006/jcph.2000.6459>. URL <http://www.sciencedirect.com/science/article/pii/S0021999100964593>.
- B. Kuźma, K. Murawski, T. V. Zaqarashvili, P. Konkol, and A. Mignone. Numerical simulations of solar spicules: Adiabatic and non-adiabatic studies. , 597:A133, January 2017. doi: 10.1051/0004-6361/201628747.
- Błażej Kuźma, Kris Murawski, Pradeep Kayshap, Darek Wójcik, Abhishek Kumar Srivastava, and Bhola N. Dwivedi. Two-fluid numerical simulations of solar spicules. *The Astrophysical Journal*, 849(2):78, 2017. URL <http://stacks.iop.org/0004-637X/849/i=2/a=78>.
- I. Langmuir. Oscillations in Ionized Gases. *Proceedings of the National Academy of Science*, 14:627–637, August 1928. doi: 10.1073/pnas.14.8.627.
- J. E. Leake, C. R. DeVore, J. P. Thayer, A. G. Burns, G. Crowley, H. R. Gilbert, J. D. Huba, J. Krall, M. G. Linton, V. S. Lukin, and W. Wang. Ionized plasma and neutral gas coupling in the sun's chromosphere and earth's ionosphere/thermosphere. *Space Science Reviews*, 184(1):107–172, Nov 2014. ISSN 1572-9672. doi: 10.1007/s11214-014-0103-1. URL <https://doi.org/10.1007/s11214-014-0103-1>.
- James E. Leake, Vyacheslav S. Lukin, Mark G. Linton, and Eric T. Meier. Multi-fluid simulations of chromospheric magnetic reconnection in a weakly ionized reacting plasma. *The Astrophysical Journal*, 760(2):109, 2012. URL <http://stacks.iop.org/0004-637X/760/i=2/a=109>.

- James E. Leake, Vyacheslav S. Lukin, and Mark G. Linton. Magnetic reconnection in a weakly ionized plasma. *Physics of Plasmas*, 20(6):061202, 2013. doi: 10.1063/1.4811140. URL <https://doi.org/10.1063/1.4811140>.
- Rita Lerner. *Encyclopedia of physics*. Wiley-VCH, Weinheim, 2005. ISBN 978-3527405541.
- R.J. LeVeque, L.R. J, and D.G. Crighton. *Finite Volume Methods for Hyperbolic Problems*. Cambridge Texts in Applied Mathematics. Cambridge University Press, 2002. ISBN 9780521009249. URL <https://books.google.com/books?id=QazcnD7GUoUC>.
- F Mandl. *Statistical physics*. Wiley, Chichester West Sussex New York, 1988. ISBN 9780471915331.
- Yana G. Maneva, Alejandro Alvarez Laguna, Andrea Lani, and Stefaan Poedts. Multi-fluid modeling of magnetosonic wave propagation in the solar chromosphere: Effects of impact ionization and radiative recombination. *The Astrophysical Journal*, 836(2):197, 2017. URL <http://stacks.iop.org/0004-637X/836/i=2/a=197>.
- Juan Martinez-Sykora, Bart De Pontieu, and Viggo Hansteen. Two-dimensional radiative magnetohydrodynamic simulations of the importance of partial ionization in the chromosphere. *The Astrophysical Journal*, 753(2):161, 2012. URL <http://stacks.iop.org/0004-637X/753/i=2/a=161>.
- Juan Martinez-Sykora, Bart De Pontieu, Mats Carlsson, Viggo H. Hansteen, Daniel Nóbrega-Siverio, and Boris V. Gudiksen. Two-dimensional radiative magnetohydrodynamic simulations of partial ionization in the chromosphere. ii. dynamics and energetics of the low solar atmosphere. *The Astrophysical Journal*, 847(1):36, 2017. URL <http://stacks.iop.org/0004-637X/847/i=1/a=36>.
- M. Mathioudakis, D. B. Jess, and R. Erdélyi. Alfvén Waves in the Solar Atmosphere. From Theory to Observations. , 175:1–27, June 2013. doi: 10.1007/s11214-012-9944-7.
- S. W. McIntosh, B. de Pontieu, M. Carlsson, V. Hansteen, P. Boerner, and M. Goossens. Alfvénic waves with sufficient energy to power the quiet solar corona and fast solar wind. , 475:477–480, July 2011. doi: 10.1038/nature10235.
- A. Mignone, P. Tzeferacos, and G. Bodo. High-order conservative finite difference GLM-MHD schemes for cell-centered MHD. *Journal of Computational Physics*, 229:5896–5920, August 2010. doi: 10.1016/j.jcp.2010.04.013.
- V. M. Nakariakov, L. Ofman, E. E. Deluca, B. Roberts, and J. M. Davila. TRACE observation of damped coronal loop oscillations: Implications for coronal heating. *Science*, 285:862–864, August 1999. doi: 10.1126/science.285.5429.862.

- L. Ni, J. Lin, I. I. Roussev, and B. Schmieder. Heating Mechanisms in the Low Solar Atmosphere through Magnetic Reconnection in Current Sheets. , 832:195, December 2016. doi: 10.3847/0004-637X/832/2/195.
- N. Nishizuka, T. Nakamura, T. Kawate, K. A. P. Singh, and K. Shibata. Statistical study of chromospheric anemone jets observed with hinode/sot. *The Astrophysical Journal*, 731(1):43, 2011. URL <http://stacks.iop.org/0004-637X/731/i=1/a=43>.
- L. Ofman and T. J. Wang. Hinode observations of transverse waves with flows in coronal loops. , 482:L9–L12, May 2008. doi: 10.1051/0004-6361:20079340.
- R.H. Pletcher, J.C. Tannehill, and D. Anderson. *Computational Fluid Mechanics and Heat Transfer, Second Edition*. Series in Computational and Physical Processes in Mechanics and Thermal Sciences. Taylor & Francis, 1997. ISBN 9781560320463. URL <https://books.google.com/books?id=ZJPbtHeilCgC>.
- B. De Pontieu, P. C. H. Martens, and H. S. Hudson. Chromospheric damping of alfvén waves. *The Astrophysical Journal*, 558(2):859, 2001. URL <http://stacks.iop.org/0004-637X/558/i=2/a=859>.
- Pete Riley. On the probability of occurrence of extreme space weather events. *Space Weather*, 10(2), 2012. doi: 10.1029/2011SW000734. URL <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2011SW000734>.
- P L Roe. Characteristic-based schemes for the euler equations. *Annual Review of Fluid Mechanics*, 18(1):337–365, 1986. doi: 10.1146/annurev.fl.18.010186.002005. URL <https://doi.org/10.1146/annurev.fl.18.010186.002005>.
- R. J. Rutten. Observing the Solar Chromosphere. In P. Heinzel, I. Dorotović, and R. J. Rutten, editors, *The Physics of Chromospheric Plasmas*, volume 368 of *Astronomical Society of the Pacific Conference Series*, page Heinzel, May 2007.
- Conrad Sanderson and Ryan Curtin. Armadillo: a template-based c++ library for linear algebra, 2016. URL <http://arma.sourceforge.net/>.
- Masumi Shimojo and Kazunari Shibata. Physical parameters of solar x-ray jets. *The Astrophysical Journal*, 542(2):1100, 2000. URL <http://stacks.iop.org/0004-637X/542/i=2/a=1100>.
- Gary A Sod. A survey of several finite difference methods for systems of nonlinear hyperbolic conservation laws. *Journal of Computational Physics*, 27(1):1 – 31, 1978. ISSN 0021-9991. doi: [https://doi.org/10.1016/0021-9991\(78\)90023-2](https://doi.org/10.1016/0021-9991(78)90023-2). URL <http://www.sciencedirect.com/science/article/pii/0021999178900232>.

- R. Soler, M. Carbonell, J. L. Ballester, and J. Terradas. Alfvén Waves in a Partially Ionized Two-fluid Plasma. , 767:171, April 2013. doi: 10.1088/0004-637X/767/2/171.
- P. Song. A model of the solar chromosphere: Structure and internal circulation. *Astrophysical Journal*, 846(2), 9 2017. doi: 10.3847/1538-4357/AA85E1.
- P. Song and V. M. Vasyliūnas. Heating of the solar atmosphere by strong damping of Alfvén waves. *Journal of Geophysical Research (Space Physics)*, 116:A09104, September 2011. doi: 10.1029/2011JA016679.
- P. Song and V. M. Vasyliūnas. Effect of horizontally inhomogeneous heating on flow and magnetic field in the chromosphere of the sun. *The Astrophysical Journal Letters*, 796(2):L23, 2014. URL <http://stacks.iop.org/2041-8205/796/i=2/a=L23>.
- The HDF Group. Hierarchical data format version 5, 2000-2010. URL <http://www.hdfgroup.org/HDF5>.
- Lewi Tonks. The birth of plasma. *American Journal of Physics*, 35(9):857–858, 1967. doi: 10.1119/1.1974266. URL <https://doi.org/10.1119/1.1974266>.
- A Tritschler, TR Rimmele, S Berukoff, R Casini, SC Craig, DF Elmore, RP Hubbard, JR Kuhn, H Lin, JP McMullin, et al. Dkist: observing the sun at high resolution. In *18th Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun*, volume 18, pages 933–944, 2015.
- J. Tu and P. Song. A two-dimensional global simulation study of inductive-dynamic magnetosphere-ionosphere coupling. *Journal of Geophysical Research (Space Physics)*, 121:11, December 2016. doi: 10.1002/2016JA023393.
- Jiannan Tu and Paul Song. A study of alfvén wave propagation and heating the chromosphere. *The Astrophysical Journal*, 777(1):53, 2013a. URL <http://stacks.iop.org/0004-637X/777/i=1/a=53>.
- Jiannan Tu and Paul Song. A study of alfvén wave propagation and heating the chromosphere. *The Astrophysical Journal*, 777(1):53, 2013b. URL <http://stacks.iop.org/0004-637X/777/i=1/a=53>.
- T. S. van Albada. Dissipationless galaxy formation and the $r^{1/4}$ law. *Monthly Notices of the Royal Astronomical Society*, 201(4):939–955, 12 1982. ISSN 0035-8711. doi: 10.1093/mnras/201.4.939. URL <https://doi.org/10.1093/mnras/201.4.939>.
- Bram van Leer. Towards the ultimate conservative difference scheme. ii. monotonicity and conservation combined in a second-order scheme. *Journal of Computational Physics*, 14(4):361 – 370, 1974. ISSN 0021-9991. doi: <https://doi.org/>

- 10.1016/0021-9991(74)90019-9. URL <http://www.sciencedirect.com/science/article/pii/0021999174900199>.
- Bram van Leer. Towards the ultimate conservative difference scheme. v. a second-order sequel to godunov's method. *Journal of Computational Physics*, 32(1):101 – 136, 1979. ISSN 0021-9991. doi: [https://doi.org/10.1016/0021-9991\(79\)90145-1](https://doi.org/10.1016/0021-9991(79)90145-1). URL <http://www.sciencedirect.com/science/article/pii/0021999179901451>.
- N. P. Waterson and H. Deconinck. Design principles for bounded higher-order convection schemes - a unified approach. *J. Comput. Phys.*, 224(1):182–207, May 2007. ISSN 0021-9991. doi: [10.1016/j.jcp.2007.01.021](https://doi.org/10.1016/j.jcp.2007.01.021). URL <http://dx.doi.org/10.1016/j.jcp.2007.01.021>.
- D Wójcik, K Murawski, and Z E Musielak. Acoustic waves in two-fluid solar atmosphere model: cut-off periods, chromospheric cavity, and wave tunnelling. *Monthly Notices of the Royal Astronomical Society*, 481(1):262–267, 2018. doi: [10.1093/mnras/sty2306](https://doi.org/10.1093/mnras/sty2306). URL <http://dx.doi.org/10.1093/mnras/sty2306>.

BIOGRAPHICAL SKETCH OF THE AUTHOR

I moved to the United States in September 2008 to attend the University of Toledo in Ohio. I majored in Physics and graduated with a B.S. in 2012. As an undergraduate researcher in University of Toledo I worked on condensed matter physics with Dr. Jacques Amar. Shortly after I moved to Massachusetts to start my PhD in Physics at the University of Massachusetts—Lowell. After a few years I landed with the Space Science Lab and Lowell Center for Space Science and Technology (LoCSST) to start my research on the solar chromosphere and in turn write MHD simulations.