

因此要使  $af(h)+bf(2h)-f(0)$  为  $h$  高阶无穷小量必须有

$$\begin{cases} a+b-1=0, \\ a+2b=0, \end{cases} \quad \text{即} \quad \begin{cases} a=2, \\ b=-1. \end{cases}$$

【例 4】 设  $f(x)$  在点  $x=0$  的某个邻域内二阶可导, 且  $\lim_{x \rightarrow 0} \frac{\sin x + xf(x)}{x^3} = \frac{1}{2}$ , 求  $f(0)$ ,  $f'(0)$  及  $f''(0)$  的值.

解 因为  $\sin x = x - \frac{1}{3!}x^3 + o(x^3)$ ,  $f(x) = f(0) + f'(0)x + \frac{1}{2!}f''(0)x^2 + o(x^2)$ .

$$\begin{aligned} \text{所以} \quad \frac{1}{2} &= \lim_{x \rightarrow 0} \frac{\sin x + xf(x)}{x^3} \\ &= \lim_{x \rightarrow 0} \frac{1}{x^3} \left[ x - \frac{1}{3!}x^3 + o(x^3) + f(0)x + f'(0)x^2 + \frac{1}{2!}f''(0)x^2 + o(x^3) \right] \\ &= \lim_{x \rightarrow 0} \frac{1}{x^3} \left[ (1+f(0))x + f'(0)x^2 + \left( \frac{1}{2}f''(0) - \frac{1}{6} \right)x^3 + o(x^3) \right], \end{aligned}$$

$$\text{从而知} \quad 1+f(0)=0, \quad f'(0)=0, \quad \frac{1}{2}f''(0)-\frac{1}{6}=\frac{1}{2},$$

$$\text{故} \quad f(0)=-1, \quad f'(0)=0, \quad f''(0)=\frac{4}{3}.$$

【例 5】 (2021 数学一, 5 分) 设函数  $f(x) = \frac{\sin x}{1+x^2}$  在  $x=0$  处的三次泰勒多项式为  $ax+bx^2+cx^3$ , 则\_\_\_\_\_.

$$(A) a=1, b=0, c=-\frac{7}{6}$$

$$(B) a=1, b=0, c=\frac{7}{6}$$

$$(C) a=-1, b=-1, c=-\frac{7}{6}$$

$$(D) a=-1, b=-1, c=\frac{7}{6}$$

解 因为  $\sin x = x - \frac{x^3}{3!} + o(x^3)$ ,  $\frac{1}{1+x^2} = 1 - x^2 + o(x^2)$ ,  $|x| < 1$ .

$$\text{故} \quad \frac{\sin x}{1+x^2} = x - \frac{7}{6}x^3 + o(x^3), \text{ 即 } a=1, b=0, c=-\frac{7}{6}.$$

故应选(A).

### 习题 3-3 解答

1. 按  $x-4$  的幂展开多项式  $f(x) = x^4 - 5x^3 + x^2 - 3x + 4$ .

解 因为  $f'(x) = 4x^3 - 15x^2 + 2x - 3$ ,

$$f''(x) = 12x^2 - 30x + 2,$$

$$f'''(x) = 24x - 30,$$

$$f^{(4)}(x) = 24,$$

$$f^{(n)}(x)=0 \quad (n \geq 5).$$

$$f(4)=-56, \quad f'(4)=21, \quad f''(4)=74, \quad f'''(4)=66, \quad f^{(4)}(4)=24.$$

$$\text{故 } x^4-5x^3+x^2-3x+4$$

$$\begin{aligned} &= f(4) + f'(4)(x-4) + \frac{f''(4)}{2!}(x-4)^2 + \frac{f'''(4)}{3!}(x-4)^3 + \frac{f^{(4)}(4)}{4!}(x-4)^4 \\ &= -56 + 21(x-4) + 37(x-4)^2 + 11(x-4)^3 + (x-4)^4. \end{aligned}$$

2. 应用麦克劳林公式,按  $x$  的幂展开函数  $f(x)=(x^2-3x+1)^3$ .

$$\begin{aligned} \text{解 } f(x) &= x^6 - 9x^5 + 30x^4 - 45x^3 + 30x^2 - 9x + 1, & f(0) &= 1, \\ f'(x) &= 6x^5 - 45x^4 + 120x^3 - 135x^2 + 60x - 9, & f'(0) &= -9, \\ f''(x) &= 30x^4 - 180x^3 + 360x^2 - 270x + 60, & f''(0) &= 60, \\ f'''(x) &= 120x^3 - 540x^2 + 720x - 270, & f'''(0) &= -270, \\ f^{(4)}(x) &= 360x^2 - 1080x + 720, & f^{(4)}(0) &= 720, \\ f^{(5)}(x) &= 720x - 1080, & f^{(5)}(0) &= -1080, \\ f^{(6)}(x) &= 720, & f^{(6)}(0) &= 720, \\ f^{(n)}(x) &= 0 \quad (n \geq 7), \end{aligned}$$

$$\text{故 } (x^2-3x+1)^3$$

$$\begin{aligned} &= f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \frac{f^{(5)}(0)}{5!}x^5 + \frac{f^{(6)}(0)}{6!}x^6 \\ &= 1 - 9x + 30x^2 - 45x^3 + 30x^4 - 9x^5 + x^6. \end{aligned}$$

3. 求函数  $f(x)=\sqrt{x}$  按  $x-4$  的幂展开的带有拉格朗日余项的 3 阶泰勒公式.

$$\text{解 } \text{因为 } f(x)=\sqrt{x}, \quad f'(x)=\frac{1}{2}x^{-\frac{1}{2}}, \quad f''(x)=-\frac{1}{4}x^{-\frac{3}{2}}, \quad f'''(x)=\frac{3}{8}x^{-\frac{5}{2}},$$

$$f^{(4)}(x)=-\frac{15}{16}x^{-\frac{7}{2}}, \quad f(4)=2, \quad f'(4)=\frac{1}{4}, \quad f''(4)=-\frac{1}{32}, \quad f'''(4)=\frac{3}{256}.$$

$$\begin{aligned} \text{故 } \sqrt{x} &= f(4) + f'(4)(x-4) + \frac{f''(4)}{2!}(x-4)^2 + \frac{f'''(4)}{3!}(x-4)^3 + \frac{f^{(4)}(\xi)}{4!}(x-4)^4 \\ &= 2 + \frac{1}{4}(x-4) - \frac{1}{64}(x-4)^2 + \frac{1}{512}(x-4)^3 - \frac{15}{384\xi^{\frac{7}{2}}}(x-4)^4, \end{aligned}$$

其中  $\xi$  介于  $x$  与 4 之间.

4. 求函数  $f(x)=\ln x$  按  $x-2$  的幂展开的带有佩亚诺型余项的  $n$  阶泰勒公式.

$$\text{解 } \text{因为 } f^{(n)}(x)=\frac{(-1)^{n-1}(n-1)!}{x^n}, \quad f^{(n)}(2)=\frac{(-1)^{n-1}(n-1)!}{2^n}, \text{ 故}$$

$$\begin{aligned} \ln x &= f(2) + f'(2)(x-2) + \frac{f''(2)}{2!}(x-2)^2 + \frac{f'''(2)}{3!}(x-2)^3 + \cdots \\ &\quad + \frac{f^{(n)}(2)}{n!}(x-2)^n + o[(x-2)^n] \\ &= \ln 2 + \frac{1}{2}(x-2) - \frac{1}{2^3}(x-2)^2 + \frac{1}{3 \times 2^3}(x-2)^3 + \cdots \\ &\quad + (-1)^{n-1} \frac{1}{n \times 2^n}(x-2)^n + o[(x-2)^n]. \end{aligned}$$

5. 求函数  $f(x)=\frac{1}{x}$  按  $x+1$  的幂展开的带有拉格朗日型余项的  $n$  阶泰勒公式.

$$\text{解 } \text{因为 } f^{(n)}(x)=\frac{(-1)^n n!}{x^{n+1}}, \quad f^{(n)}(-1)=-n!, \text{ 故}$$

$$\frac{1}{x} = f(-1) + f'(-1)(x+1) + \frac{f''(-1)}{2!}(x+1)^2 + \frac{f'''(-1)}{3!}(x+1)^3 + \cdots$$



5 题视频解析

$$\begin{aligned}
 & + \frac{f^{(n)}(-1)}{n!} (x+1)^n + \frac{f^{(n+1)}(\xi)}{(n+1)!} (x+1)^{n+1} \\
 & = -[1 + (x+1) + (x+1)^2 + \cdots + (x+1)^n] + (-1)^{n+1} \xi^{-(n+2)} (x+1)^{n+1},
 \end{aligned}$$

其中  $\xi$  介于  $x$  与  $-1$  之间.

6. 求函数  $f(x) = \tan x$  的带有佩亚诺余项的 3 阶麦克劳林公式.

**解** 因为  $f(x) = \tan x$ ,  $f'(x) = \sec^2 x$ ,  $f''(x) = 2\sec^2 x \tan x$ ,  $f'''(x) = 4\sec^2 x \tan^2 x + 2\sec^4 x$ ,  
 $f(0) = 0$ ,  $f'(0) = 1$ ,  $f''(0) = 0$ ,  $f'''(0) = 2$ ,

因此  $f(x) = x + \frac{x^3}{3} + o(x^3)$ .

7. 求函数  $f(x) = xe^x$  的带有佩亚诺余项的  $n$  阶麦克劳林公式.

**解** 因为  $f(x) = xe^x$ ,  $f^{(n)}(x) = (n+x)e^x$  (见习题 2-3, 10(4)),  $f^{(n)}(0) = n$ , 故

$$\begin{aligned}
 xe^x &= f(0) + f'(0)x + \frac{1}{2!}f''(0)x^2 + \cdots + \frac{1}{n!}f^{(n)}(0)x^n + o(x^n) \\
 &= x + x^2 + \frac{x^3}{2!} + \cdots + \frac{x^n}{(n-1)!} + o(x^n).
 \end{aligned}$$

8. 验证当  $0 < x \leq \frac{1}{2}$  时, 按公式  $e^x \approx 1 + x + \frac{x^2}{2} + \frac{x^3}{6}$  计算  $e^x$  的近似值时, 所产生的误差小于 0.01, 并求  $\sqrt{e}$  的近似值, 使误差小于 0.01.

**证** 设  $f(x) = e^x$ , 则  $f^{(n)}(0) = 1$ , 故  $f(x) = e^x$  的三阶麦克劳林公式为  $e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{e^\xi}{4!}x^4$ ,

其中  $\xi$  介于  $0, x$  之间. 按  $e^x \approx 1 + x + \frac{x^2}{2} + \frac{x^3}{6}$  计算  $e^x$  的近似值时, 其误差为  $|R_3(x)| = \frac{e^\xi}{4!}x^4$

当  $0 < x \leq \frac{1}{2}$  时,  $0 < \xi \leq \frac{1}{2}$ ,  $|R_3(x)| \leq \frac{3^{\frac{1}{2}}}{4!} \left(\frac{1}{2}\right)^4 \approx 0.0045 < 0.01$ ,

$$\sqrt{e} \approx 1 + \frac{1}{2} + \frac{1}{2} \left(\frac{1}{2}\right)^2 + \frac{1}{6} \left(\frac{1}{2}\right)^3 \approx 1.645.$$

9. 应用 3 阶泰勒公式求下列各数的近似值, 并估计误差:

(1)  $\sqrt[3]{30}$ ; (2)  $\sin 18^\circ$ .

**解** (1) 因为  $f(x) = \sqrt[3]{1+x} = (1+x)^{\frac{1}{3}} \approx 1 + \frac{1}{3}x + \frac{\frac{1}{3}(\frac{1}{3}-1)}{2!}x^2 + \frac{\frac{1}{3}(\frac{1}{3}-1)(\frac{1}{3}-2)}{3!}x^3$

$$= 1 + \frac{1}{3}x - \frac{1}{9}x^2 + \frac{5}{81}x^3,$$

$$R_3(x) = \frac{\frac{1}{3}(\frac{1}{3}-1)(\frac{1}{3}-2)(\frac{1}{3}-3)}{4!}(1+\xi)^{\frac{1}{3}-4}x^4,$$

其中  $\xi$  介于  $0$  与  $x$  之间, 故

$$\sqrt[3]{30} = \sqrt[3]{27+3} = 3\sqrt[3]{1+\frac{1}{9}} \approx 3\left[1 + \frac{1}{3} \times \frac{1}{9} - \frac{1}{9} \left(\frac{1}{9}\right)^2 + \frac{5}{81} \left(\frac{1}{9}\right)^3\right] \approx 3.10724.$$

$$\text{误差 } |R_3| = 3 \times \left| \frac{\frac{1}{3}(\frac{1}{3}-1)(\frac{1}{3}-2)(\frac{1}{3}-3)}{4!}(1+\xi)^{\frac{1}{3}-4} \left(\frac{1}{9}\right)^4 \right|,$$

$\xi$  介于  $0$  与  $\frac{1}{9}$  之间, 即  $0 < \xi < \frac{1}{9}$ , 因此  $|R_3| = \left| \frac{80}{4! \times 3^{11}} \right| \approx 1.88 \times 10^{-5}$ .

(2) 已知  $\sin x \approx x - \frac{x^3}{3!}$ ,  $R_4(x) = \frac{\sin(\xi + \frac{5}{2}\pi)}{5!}x^5$ ,  $\xi$  介于  $0$  与  $\frac{\pi}{10}$  之间, 故

$$\sin 18^\circ = \sin \frac{\pi}{10} \approx \frac{\pi}{10} - \frac{1}{3!} \left(\frac{\pi}{10}\right)^3 \approx 0.3090,$$



$$|R_4| \leq \frac{1}{5!} \left(\frac{\pi}{10}\right)^5 \approx 2.55 \times 10^{-5}.$$

评注: 利用  $R_3(x) = \frac{\sin(\xi + \frac{4}{2}\pi)}{4!} x^4$ ,  $\xi \in (0, \frac{\pi}{10})$ , 可得误差  $|R_3| \leq \frac{1}{4!} \left(\frac{\pi}{10}\right)^4 \approx 1.3 \times 10^{-4}$ .

• 10. 利用泰勒公式求下列极限:

$$(1) \lim_{x \rightarrow +\infty} (\sqrt[3]{x^3+3x^2} - \sqrt{x^4-2x^3});$$

$$(2) \lim_{x \rightarrow 0} \frac{\cos x - e^{-\frac{x^2}{2}}}{x^2 [x + \ln(1-x)]};$$

$$(3) \lim_{x \rightarrow 0} \frac{1 + \frac{1}{2}x^2 - \sqrt{1+x^2}}{(\cos x - e^{x^2}) \sin x^2};$$

$$(4) \lim_{x \rightarrow \infty} \left[ x - x^2 \ln \left( 1 + \frac{1}{x} \right) \right].$$



10 题视频解析

解 (1)  $\lim_{x \rightarrow +\infty} (\sqrt[3]{x^3+3x^2} - \sqrt{x^4-2x^3}) = \lim_{x \rightarrow +\infty} x \left[ \left(1 + \frac{3}{x}\right)^{\frac{1}{3}} - \left(1 - \frac{2}{x}\right)^{\frac{1}{2}} \right]$

$$= \lim_{x \rightarrow +\infty} x \left[ 1 + \frac{1}{3} \times \frac{3}{x} + o\left(\frac{1}{x}\right) - 1 + \frac{1}{4} \times \frac{2}{x} + o\left(\frac{1}{x}\right) \right] = \lim_{x \rightarrow +\infty} \left[ \frac{3}{2} + \frac{o\left(\frac{1}{x}\right)}{\frac{1}{x}} \right] = \frac{3}{2}.$$

(2)  $\lim_{x \rightarrow 0} \frac{\cos x - e^{-\frac{x^2}{2}}}{x^2 [x + \ln(1-x)]} = \lim_{x \rightarrow 0} \frac{1 - \frac{x^2}{2} + \frac{x^4}{4!} + o(x^4) - 1 - \left(-\frac{x^2}{2}\right) - \frac{1}{2} \left(-\frac{x^2}{2}\right)^2 + o(x^4)}{x^2 \left[ x + \left(-x - \frac{1}{2}x^2 + o(x^2)\right) \right]}$

$$= \lim_{x \rightarrow 0} \frac{\left(\frac{1}{4!} - \frac{1}{8}\right)x^4 + o(x^4)}{-\frac{1}{2}x^4 + o(x^4)} = \lim_{x \rightarrow 0} \frac{-\frac{1}{12} + \frac{o(x^4)}{x^4}}{-\frac{1}{2} + \frac{o(x^4)}{x^4}} = \frac{-\frac{1}{12}}{-\frac{1}{2}} = \frac{1}{6}.$$

(3)  $\lim_{x \rightarrow 0} \frac{1 + \frac{1}{2}x^2 - \sqrt{1+x^2}}{(\cos x - e^{x^2}) \sin x^2} = \lim_{x \rightarrow 0} \frac{1 + \frac{1}{2}x^2 - \left(1 + \frac{1}{2}x^2 - \frac{1}{8}x^4 + o(x^4)\right)}{\left[1 - \frac{1}{2}x^2 + o(x^2) - 1 - x^2 + o(x^2)\right] [x^2 + o(x^2)]}$

$$= \lim_{x \rightarrow 0} \frac{\frac{1}{8}x^4 + o(x^4)}{-\frac{3}{2}x^4 + o(x^4)} = \lim_{x \rightarrow 0} \frac{\frac{1}{8} + \frac{o(x^4)}{x^4}}{-\frac{3}{2} + \frac{o(x^4)}{x^4}} = \frac{\frac{1}{8}}{-\frac{3}{2}} = -\frac{1}{12}.$$

(4)  $\lim_{x \rightarrow \infty} \left[ x - x^2 \ln \left( 1 + \frac{1}{x} \right) \right] = \lim_{x \rightarrow \infty} \left[ x - x^2 \left( \frac{1}{x} - \frac{1}{2} \cdot \frac{1}{x^2} + o\left(\frac{1}{x^2}\right) \right) \right]$

$$= \lim_{x \rightarrow \infty} \left[ x - x + \frac{1}{2} + \frac{o\left(\frac{1}{x^2}\right)}{\frac{1}{x^2}} \right] = \frac{1}{2}.$$

11. 若函数  $f(x)$  在  $(-\infty, +\infty)$  内具有二阶导数, 且  $f''(x) > 0$ , 又  $\lim_{x \rightarrow 0} \frac{f(x)}{x} = 1$ , 证明:  $f(x) \geq x, x \in (-\infty, +\infty)$ .



11 题视频解析

证 易知  $f(x)$  在  $x=0$  处连续, 由  $\lim_{x \rightarrow 0} \frac{f(x)}{x} = 1$  知  $f(0) = 0$  且  $f'(0) = 1$ ,

将  $f(x)$  在  $x=0$  处展开, 可得

$$f(x) = f(0) + f'(0)x + \frac{f''(\xi)}{2!}x^2 = x + \frac{f''(\xi)}{2!}x^2 \geq x \quad (\xi \text{ 介于 } 0 \text{ 与 } x \text{ 之间}).$$