

A Guide to the Depression, Anxiety and Stress Scale

(DASS 21)

By Fernando Gomez- Consultant Clinical Psychologist

Introduction

The DASS 21 is a 21 item self report questionnaire designed to measure the severity of a range of symptoms common to both Depression and Anxiety. In completing the DASS, the individual is required to indicate the presence of a symptom over the previous week. Each item is scored from 0 (did not apply to me at all over the last week) to 3 (applied to me very much or most of the time over the past week).

The essential function of the DASS is to assess the severity of the core symptoms of Depression, Anxiety and Stress. Accordingly, the DASS allows not only a way to measure the severity of a patient's symptoms but a means by which a patient's response to treatment can also be measured.

The DASS and Diagnosis

Although the DASS may contribute to the diagnosis of Anxiety or Depression, it is not designed as a diagnostic tool. Indeed, a number of symptoms typical of Depression such as sleep, appetite and sexual disturbances, are not covered by the DASS and will need to be assessed independently. The DASS is not meant to replace a comprehensive clinical interview.

Suicide

Suicidality is not assessed by the DASS. Accordingly, the clinician will need to address directly this important symptom of Depression in their clinical interview.

How often to use the DASS?

Although the DASS can provide a comparison of symptoms from week to week, it is best given on first presentation and again after a period of time has lapsed long enough for the chosen treatment to have effect. In the case of antidepressant medication, the second administration should be between the 2-4 week period after the individual has commenced taking the medication. This period is long enough for most antidepressants to be expected to show some change in the patient.

Scoring the DASS

The scale to which each item belongs is indicated by the letters D (Depression), A (Anxiety) and S (Stress). For each scale (D, A & S) sum the scores for identified items. Because the DASS 21 is a short form version of the DASS (the Long Form has 42 items), the final score of each item groups (Depression, Anxiety and Stress) needs to be multiplied by two (x2).

Interpreting the DASS

Once multiplied by 2, each score can now be transferred to the DASS profile sheet, enabling comparisons to be made between the three scales and also giving percentile rankings and severity labels.

DASS Severity Ratings

(Don't forget to multiply summed scores by x 2)

Severity	Depression	Anxiety	Stress
Normal	0-9	0-7	0-14
Mild	10-13	8-9	15-18
Moderate	14-20	10-14	19-25
Severe	21-27	15-19	26-33
Extremely Severe	28+	20+	34+

As previously mentioned, the DASS should not be used on its own to assess the presence or absence of Depression or Anxiety. High scores on the DASS would certainly alert the clinician to a high level of distress in the patient and this would need to be explored further within the interview process. Similarly, low scores on the DASS should not be a substitute for a comprehensive clinical interview.

High DASS scores which are not changing, may prompt the clinician to look for explanations and perhaps augment dosages or change medication. Here again, the DASS should be interpreted alongside the clinical interview.

Changes in scores in one scale (EG: Depression), with consistently high and unchanging scores in another scale (Anxiety) may alert the clinician to pay particular attention to the presence of a co-existing anxiety disorder which may need specific treatment in its own right. Similarly, decreasing Depression scores alongside unchanging Stress scores may alert the clinician to the presence of some life event or problem, which may need to be addressed directly.



DASS 21

NAME _____ DATE _____

BLACK DOG INSTITUTE

Please read each statement and circle a number 0, 1, 2 or 3 which indicates how much the statement applied to you over the past week. There are no right or wrong answers. Do not spend too much time on any statement.

The rating scale is as follows:

- 0 Did not apply to me at all - NEVER
- 1 Applied to me to some degree, or some of the time - SOMETIMES
- 2 Applied to me to a considerable degree, or a good part of time - OFTEN
- 3 Applied to me very much, or most of the time - ALMOST ALWAYS

FOR OFFICE USE

		N	S	O	AA	D	A	S
1	I found it hard to wind down	0	1	2	3			
2	I was aware of dryness of my mouth	0	1	2	3			
3	I couldn't seem to experience any positive feeling at all	0	1	2	3			
4	I experienced breathing difficulty (eg, excessively rapid breathing, breathlessness in the absence of physical exertion)	0	1	2	3			
5	I found it difficult to work up the initiative to do things	0	1	2	3			
6	I tended to over-react to situations	0	1	2	3			
7	I experienced trembling (eg, in the hands)	0	1	2	3			
8	I felt that I was using a lot of nervous energy	0	1	2	3			
9	I was worried about situations in which I might panic and make a fool of myself	0	1	2	3			
10	I felt that I had nothing to look forward to	0	1	2	3			
11	I found myself getting agitated	0	1	2	3			
12	I found it difficult to relax	0	1	2	3			
13	I felt down-hearted and blue	0	1	2	3			
14	I was intolerant of anything that kept me from getting on with what I was doing	0	1	2	3			
15	I felt I was close to panic	0	1	2	3			
16	I was unable to become enthusiastic about anything	0	1	2	3			
17	I felt I wasn't worth much as a person	0	1	2	3			
18	I felt that I was rather touchy	0	1	2	3			
19	I was aware of the action of my heart in the absence of physical exertion (eg, sense of heart rate increase, heart missing a beat)	0	1	2	3			
20	I felt scared without any good reason	0	1	2	3			
21	I felt that life was meaningless	0	1	2	3			
TOTALS								

DASS Severity Ratings

The DASS is a **quantitative** measure of distress along the 3 axes of depression, anxiety¹ and stress². It is not a categorical measure of clinical diagnoses.

Emotional syndromes like depression and anxiety are intrinsically dimensional - they vary along a continuum of severity (independent of the specific diagnosis). Hence the selection of a single cut-off score to represent clinical severity is necessarily arbitrary. A scale such as the DASS can lead to a useful assessment of **disturbance**, for example individuals who may fall short of a clinical cut-off for a specific diagnosis can be correctly recognised as experiencing considerable symptoms and as being at high risk of further problems.

However for clinical purposes it can be helpful to have 'labels' to characterise degree of severity relative to the population. Thus the following cut-off scores have been developed for defining mild/moderate/severe/extremely severe scores for each DASS scale.

Note: the severity labels are used to describe the full range of scores in the population, so 'mild' for example means that the person is above the population mean but probably still way below the typical severity of someone seeking help (ie it does not mean a mild level of disorder).

The individual DASS scores do not define appropriate interventions. They should be used in conjunction with all clinical information available to you in determining appropriate treatment for any individual.

¹Symptoms of psychological arousal

²The more cognitive, subjective symptoms of anxiety

DASS 21 SCORE

DEPRESSION SCORE	ANXIETY SCORE	STRESS SCORE

	Depression	Anxiety	Stress
Normal	0 - 4	0 - 3	0 - 7
Mild	5 - 6	4 - 5	8 - 9
Moderate	7 - 10	6 - 7	10 - 12
Severe	11 - 13	8 - 9	13 - 16
Extremely Severe	14 +	10 +	17 +