

Verifying Tartaglia's Theory of Cannonball Motion

Pranav Arunkumar,^{*} Dylan Arellano, Lukas Fendrick, Carter Krawiec, and Sophia Maselli[†]
Science & Engineering Magnet Program, Manalapan High School, Englishtown, NJ 07726 USA
 (Received 13 May 2026; accepted 13 May 2026; published 13 May 2026)

Abstract goes here

I. INTRODUCTION

Nicolo Tartaglia was an Italian Renaissance mathematician, who was tasked with modeling the motion of cannonballs by the Doge of Venice in order to confer a military advantage. His predicted trajectory of a cannonball is shown in Figure 1.

He also predicted the horizontal ranges of cannonballs shot at various angles, concluding that they are symmetric at 45° and travel furthest at 45°. In this experiment, we aim to test his theories with more established mathematical models of projectile motion.

Consider the horizontal and vertical components of projectile motion as given by Equations 1 and 2, where v_0 represents the initial velocity of the projectile, θ represents the initial angle, and x_0 and y_0 represent the initial positions on their respective axis [1].

$$x(t) = v_0 \cos \theta t + x_0 \quad (1)$$

$$y(t) = v_0 \sin \theta t - \frac{1}{2}gt^2 - y_0 \quad (2)$$

In order to calculate the horizontal range, we need to compute when $y(t) = 0$ via quadratic formula, and substitute the resulting t value into $x(t)$.

Doing so, we get our final equation for which we fitted the data to.

$$\text{displacement} = v_0 \cos \theta * \frac{-v_0 \sin \theta - \sqrt{v^2 \sin^2 \theta - 4(y_0)(-\frac{1}{2}g)}}{-g} \quad (3)$$

II. METHODS AND MATERIALS

We conducted this experiment using a PASCO ME-6825 Mini Launcher (PASCO Scientific, Roseville, California) alongside a 16mm stainless steel marble with an average mass of 16.7g across 7 marbles, weighed using a Digital Pro Pocket Scale (Smart Weigh, New York, NY). We used these materials to repeat our methods across three trials. We used a C-clamp to hold the launcher against the table to ensure it did not move much over the course of the experiment. Setting this up, we used two metersticks, one at the standard length of 100cm, and another snapped in half, measuring around 55cm. We lined these up next to each other to measure the distance the marble was shot out of our launcher.

We began each of our three trials by setting our mini launcher to be parallel to the ground, at an angle of 0°. We then inserted the marble into the launcher until it clicked once. We recorded each trial with the webcam of the Lenovo Ideapad 5i (Lenovo, Morrisville, North Carolina) at 30 FPS. We would then shoot the marble, having a group member estimate the distance of the shot by recording the length at which the marble fell below the table. Then, using a protractor, we increased the angle by 7.5° and repeated the process again, eventually stopping the trial after 90°.

^{*} Manalapan High School

[†] Contact author: 428parunkumar@frhsd.com

Published by the Journal of Science & Engineering under the terms of the [Creative Commons Attribution-NonCommercial 4.0 International](https://creativecommons.org/licenses/by-nc/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

At a later point in time, we recorded a brief trajectory shot using an Arducam OV9281 1MP USB Camera (Arducam, Nanjing, China) set to 60FPS. We then extracted the frames via ffmpeg(v8.1.2) and superimposed the images via AstroImageJ[2].

To perform statistical analysis, we plotted our data and created a regression model with the use of the NumPy, SciPy, and Matplotlib libraries[3–5].

All code and figures used in this paper can be found at <https://github.com/qazniak-bl/marble-lab>.

III. RESULTS

The data in Table I show our measurements across three trials at different angles and a calculated average horizontal range.

Using this data we used python to plot and create a regression function according to the model in Equation (3) via Scipy and Matplotlib[4, 5]. Our regression is visible in Fig. 2.

We used this same regression to generate a plot of the residuals, visible in Fig. 3.

To verify that the regression equation was a good fit for the data, we conducted a Shapiro-Wilk test on the residuals to assess their normality, with the null hypothesis being that if the regression is a good fit for the data, the residuals would be normally distributed. In doing so, we got a test statistic of 0.9473 and a p-value of 0.06682.

The super-imposed trajectory of the marble can be found in Fig. 4.

IV. DISCUSSION

Our polynomial fit produces an R^2 value of 0.8971, which means that our model accounted for 89.71% of the variation when we model horizontal range as a function of angle. This indicates our model fit the data fairly well. The results of our Shapiro test also corroborate this: our p-value of 0.06682 is greater than our significance level of $\alpha = 0.5$, so we have no reason to reject our null hypothesis that the residuals are normally distributed.

Furthermore, our trajectory found in Fig. 4 is nothing like what Tartaglia predicted in Fig. 1. As such, we can discredit his second theory.

Tartaglia also predicted that horizontal range would be maximized at 45° , but our data in Table I shows that horizontal range actually reached its maximum at 37.5° . Additionally, there is no obvious symmetry around the angle of maximum range, although this could be due to our lack of proper recording equipment. As such, we conclude that Tartaglia's initial claim about the angle of maximum range was wrong, but we can not reliably deny his claim regarding the symmetry of the range.

We encountered a couple human errors as we did our tests and experiments. A major experience with these errors was a mistake we encountered while angling the cannon. These measurements caused our own data tables to have a large margin of error as we progressed in our trials. Our cannon was supposed to start at 0° , incrementing by 7.5° for each shot until 90° is reached; however we ended up shortening the increment, and so, when we were intending to be at 90° when we were actually at around 80° . To remedy this, we re-did our original trial. This took more time, and in hindsight we should have been more precise with our measurements. We also did not have the greatest tools to record the distance at which the marble fell below the table.

Additionally, the camera we used to map the trajectory of the marble (Arducam) had some lens distortion, resulting in a parabolic view of the shot. While we could have accounted for this, we had no need to, as this video was only used to map out the trajectory and not used to extract measurements.

V. ACKNOWLEDGEMENTS

SM tracked the motion of the marble and recorded the distance it traveled. LF recorded the videos and wrote part of the discussion. PA and CK took turns launching the marble at various degrees in different trials. DA ensured the marble did not stray too far from the site after each launch.

[1] P. A. Tipler and G. Mosca, *Physics for Scientists and Engineers*, 5th ed. (W H Freeman and Company, New York, 2004).

[2] e. a. Collins, Kielkopf, Astroimagej: Image processing and photometric extraction for ultra-precise astronomical light curves, *The Astronomical Journal* **151**, 10.3847/1538-3881/153/2/77 (2017).

- [3] C. R. Harris, K. J. Millman, S. J. van der Walt, R. Gommers, P. Virtanen, D. Cournapeau, E. Wieser, J. Taylor, S. Berg, N. J. Smith, R. Kern, M. Picus, S. Hoyer, M. H. van Kerkwijk, M. Brett, A. Haldane, J. F. del Río, M. Wiebe, P. Peterson, P. Gérard-Marchant, K. Sheppard, T. Reddy, W. Weckesser, H. Abbasi, C. Gohlke, and T. E. Oliphant, Array programming with *numpy*, *Nature* **585**, 357 (2020).
- [4] P. Virtanen, R. Gommers, T. E. Oliphant, M. Haberland, T. Reddy, D. Cournapeau, E. Burovski, P. Peterson, W. Weckesser, J. Bright, S. J. van der Walt, M. Brett, J. Wilson, K. J. Millman, N. Mayorov, A. R. J. Nelson, E. Jones, R. Kern, E. Larson, C. J. Carey, Í. Polat, Y. Feng, E. W. Moore, J. VanderPlas, D. Laxalde, J. Perktold, R. Cimrman, I. Henriksen, E. A. Quintero, C. R. Harris, A. M. Archibald, A. H. Ribeiro, F. Pedregosa, P. van Mulbregt, and SciPy 1.0 Contributors, *scipy* 1.0: fundamental algorithms for scientific computing in Python, *Nature Methods* **17**, 261 (2020).
- [5] J. D. Hunter, *matplotlib*: a 2D graphics environment, *Computing in Science & Engineering* **9**, 90 (2007).

FIGURES




FIG. 1. Tartaglia's prediction of cannonball flight




FIG. 2. Plot of horizontal range vs. degrees




FIG. 3. Plot of residuals




FIG. 4. Trajectory of the marble

TABLES

TABLE I. Marble Horizontal Range at Different Launch Angles

Angle (°)	Trial 1 (cm)	Trial 2 (cm)	Trial 3 (cm)	Average (cm)
0.0	20	25	21	22.00
7.5	51	50	52	51.00
15.0	75	65	60	66.67
22.5	99	96	87	94.00
30.0	103	97	97	99.00
37.5	102	103	98	101.00
45.0	99	96	91	95.33
52.5	90	90	85	88.33
60.0	80	77	76	77.67
67.5	45	46	47	46.00
75.0	30	32	31	31.00
82.5	2	2	10	4.67
90.0	0	0	0	0.00