0.1 Syntax

$$\begin{array}{lll} Ar & ::= s_k \mid (x :_U M) \rightarrow Ar \\ Pos & ::= X \overrightarrow{m} \mid (x :_s M) \rightarrow Pos \mid (x :_s M) \multimap Pos \\ Co_U & ::= X \overrightarrow{m} \mid (_:_U P) \rightarrow Co'_U \mid (x :_U M) \rightarrow Co'_U \\ Co'_U & ::= X \overrightarrow{m} \mid (_:_U P) \multimap Co'_U \mid (x :_U M) \multimap Co'_U \\ Co_L & ::= X \overrightarrow{m} \mid (_:_s P) \rightarrow Co'_L \mid (x :_U M) \rightarrow Co'_L \mid (_:_L M) \rightarrow Co'_L \\ Co'_L & ::= X \overrightarrow{m} \mid (_:_s P) \multimap Co'_L \mid (x :_U M) \multimap Co'_L \mid (_:_L M) \multimap Co'_L \\ \end{array}$$

0.2 Introduction Rules

$$\frac{(\forall i=1...n) \quad arity(A,s_k) \quad constructor_s(C_i,X)}{|\Gamma| \quad \Gamma \vdash A: U_{k+1} \quad \Gamma, X:_U A \vdash C_i: U_k} \\ \frac{|\Gamma| \quad \Gamma \vdash Ind_s(X:A)\{C_1|...|C_n\}: A}{\Gamma \vdash Ind_s(X:A)\{C_1|...|C_n\}: A} \qquad \frac{|\Gamma| \quad \Gamma \vdash Ind_s(X:A)\{C_1|...|C_n\}: A}{\Gamma \vdash Constr(i,I_s): C_i[I_s/X]}$$

0.3 Non-Dependent Elimination

$$\begin{array}{ll} ((x:_{s'}M) \rightarrow P)\{s\} & = U \\ ((x:_{s'}M) \multimap P)\{s\} & = L \\ (X \overrightarrow{m})\{s\} & = s \end{array}$$

$$\begin{array}{lll} ((_:_UP)\to C)\{X,Q,s\} & = (_:_UP)\to (_:_{s'}P[Q/X])\multimap C\{X,Q,s\} & (where \ s'=P\{s\}) \\ ((_:_UP)\multimap C)\{X,Q,s\} & = (_:_UP)\multimap (_:_{s'}P[Q/X])\multimap C\{X,Q,s\} & (where \ s'=P\{s\}) \\ ((_:_LP)\to C)\{X,Q,s\} & = (_:_{s'}P[Q/X])\to C\{X,Q,s\} & (where \ s'=P\{s\}) \\ ((_:_LP)\multimap C)\{X,Q,s\} & = (_:_{s'}P[Q/X])\multimap C\{X,Q,s\} & (where \ s'=P\{s\}) \\ ((x:_sM)\to C)\{X,Q,s\} & = (x:_sM)\to C\{X,Q,s\} \\ ((x:_sM)\multimap C)\{X,Q,s\} & = (x:_sM)\multimap C\{X,Q,s\} \\ (X\overrightarrow{m})\{X,Q,s\} & = Q\overrightarrow{m} \end{array}$$

$$\begin{array}{ll} ((_:_s P) \to C)\{X,Q\} & = (_:_s P) \multimap C\{X,Q\} \\ ((_:_s P) \multimap C)\{X,Q\} & = (_:_s P) \multimap C\{X,Q\} \\ ((x:_s M) \to C)\{X,Q\} & = (x:_s M) \multimap C\{X,Q\} \\ ((x:_s M) \multimap C)\{X,Q\} & = (x:_s M) \multimap C\{X,Q\} \\ (X \overrightarrow{m})\{X,Q\} & = Q \overrightarrow{m} \end{array}$$

$$\frac{\Gamma_1 \vdash c : (I_s \overrightarrow{d}) \quad \begin{array}{c|c} (\forall i = \ 1...n) & |\Gamma_2| & \Gamma_1 \ddagger \Gamma_2 \ddagger \Gamma \\ \hline \Gamma_1 \vdash c : (I_s \overrightarrow{d}) & \Gamma_2 \vdash Q : (\overrightarrow{x} :_U \overrightarrow{A}) \rightarrow s' & \Gamma_2 \vdash f_i : C_i \{I_s, Q, s'\} \\ \hline \Gamma \vdash Rec(c, Q) \{f_1|...|f_n\} : (Q\overrightarrow{d}) \end{array}$$

$$\frac{(\forall i = 1...n)}{\Gamma_1 \vdash c : (I_s \overrightarrow{d})} \frac{(\forall i = 1...n)}{\overline{\Gamma_2} \vdash Q : (\overrightarrow{x} :_U \overrightarrow{A}) \rightarrow s'} \frac{\Gamma_1 \ddagger \Gamma_2 \ddagger \Gamma}{\Gamma_2 \vdash f_i : C_i \{I_s, Q\}} \frac{\Gamma_1 \vdash c : (I_s \overrightarrow{d})}{\Gamma \vdash Elim(c, Q) \{f_1 | ... | f_n\} : (Q \overrightarrow{d})}$$

0.4 Dependent Elimination

$$\begin{array}{ll} ((x:_sM)\to P)\{Q,p\} & = (x:_sM)\to P\{Q,(p\ x)\} \\ ((x:_sM)\multimap P)\{Q,p\} & = (x:_sM)\multimap P\{Q,(p\ x)\} \\ (X\ \overrightarrow{m})\{Q,p\} & = Q\ \overrightarrow{m}\ p \end{array}$$

$$\begin{array}{lll} ((_:_UP)\to C)\{X,Q,s,c\} & = (p:_UP)\to (_:_{s'}P\{Q,p\}) \multimap C\{X,Q,s,(c\;p)\} & (where\;s'=P\{s\}) \\ ((_:_UP)\multimap C)\{X,Q,s,c\} & = (p:_UP)\multimap (_:_{s'}P\{Q,p\}) \multimap C\{X,Q,s,(c\;p)\} & (where\;s'=P\{s\}) \\ ((x:_UM)\to C)\{X,Q,s,c\} & = (x:_UM)\to C\{X,Q,s,(c\;x)\} \\ ((x:_UM)\multimap C)\{X,Q,s,c\} & = (x:_UM)\multimap C\{X,Q,s,(c\;x)\} \\ (X\;\overrightarrow{m})\{X,Q,s,c\} & = (Q\;\overrightarrow{m}\;c) \end{array}$$

$$\begin{array}{ll} ((_:_UP)\to C)\{X,Q,c\} & = (p:_UP)\multimap C\{X,Q,(c\;p)\} \\ ((_:_UP)\multimap C)\{X,Q,c\} & = (p:_UP)\multimap C\{X,Q,(c\;p)\} \\ ((x:_UP)\to C)\{X,Q,c\} & = (x:_UM)\multimap C\{X,Q,(c\;x)\} \\ ((x:_UP)\multimap C)\{X,Q,c\} & = (x:_UM)\multimap C\{X,Q,(c\;x)\} \\ (X\;\overrightarrow{m})\{X,Q,c\} & = (Q\;\overrightarrow{m}\;c) \end{array}$$

$$\frac{(\forall i = 1...n) \qquad |\Gamma_2| \qquad \Gamma_1 \ddagger \Gamma_2 \ddagger \Gamma}{\Gamma_1 \vdash c : (I_U \overrightarrow{a}) \qquad \Gamma_2 \vdash Q : (\overrightarrow{x} :_U \overrightarrow{A}) \rightarrow (_:_U I_U \overrightarrow{x}) \rightarrow s' \qquad \Gamma_2 \vdash f_i : C_i \{I_U, Q, s', Constr(i, I_U)\}}{\Gamma \vdash Rec(c, Q) \{f_1|...|f_n\} : (Q \overrightarrow{a} c)}$$

$$\frac{(\forall i = 1...n) \qquad \Gamma_1 \ddagger \Gamma_2 \ddagger \Gamma}{\Gamma_1 \vdash c : (I_U \overrightarrow{a}) \qquad \overline{\Gamma_2} \vdash Q : (\overrightarrow{x} :_U \overrightarrow{A}) \rightarrow (_:_U I_U \overrightarrow{x}) \rightarrow s' \qquad \Gamma_2 \vdash f_i : C_i \{I_U, Q, Constr(i, I_U)\}\}}{\Gamma \vdash Elim(c, Q) \{f_1 | ... | f_n\} : (Q \overrightarrow{a} c)}$$

0.5 Conversion

$$\begin{array}{ll} ((\cdot :_{U}P) \rightarrow C)[X,F,f] & = \lambda p.C[X,F,(f\ p\ \lambda\overrightarrow{x}.(F\ \overrightarrow{m}\ (p\ \overrightarrow{x})))] \\ ((\cdot :_{U}P) \rightarrow C)[X,F,f] & = \lambda p.C[X,F,(f\ p\ \lambda\overrightarrow{x}.(F\ \overrightarrow{m}\ (p\ \overrightarrow{x})))] \\ ((\cdot :_{L}P) \rightarrow C)[X,F,f] & = \lambda p.C[X,F,(f\ \lambda\overrightarrow{x}.(F\ \overrightarrow{m}\ (p\ \overrightarrow{x})))] \\ ((\cdot :_{L}P) \rightarrow C)[X,F,f] & = \lambda p.C[X,F,(f\ \lambda\overrightarrow{x}.(F\ \overrightarrow{m}\ (p\ \overrightarrow{x})))] \\ ((x:_{S}M) \rightarrow C)[X,F,f] & = \lambda x.C[X,F,(f\ x)] \\ ((x:_{S}M) \rightarrow C)[X,F,f] & = \lambda x.C[X,F,(f\ x)] \\ (X\ \overrightarrow{m})[X,F,f] & = f \end{array}$$

$$\begin{array}{ll} ((_:_s P) \to C)[X,f] & = \lambda p.C[X,(f \ p)] \\ ((_:_s P) \multimap C)[X,f] & = \lambda p.C[X,(f \ p)] \\ ((x:_s M) \to C)[X,f] & = \lambda x.C[X,(f \ x)] \\ ((x:_s M) \multimap C)[X,f] & = \lambda x.C[X,(f \ x)] \\ (X \overrightarrow{m})[X,f] & = f \end{array}$$

$$Fun_Rec(I_s, Q, \overrightarrow{f}) = \lambda \overrightarrow{x}.\lambda c.Rec(c, Q)\{\overrightarrow{f}\}$$

 $Rec((Constr(i, I_s) \overrightarrow{m}), Q)\{\overrightarrow{f}\} \qquad \rightarrow_{\iota} \qquad (C_i[I_s, Fun_Rec(I_s, Q, \overrightarrow{f}), f_i] \overrightarrow{m})$ $Elim((Constr(i, I_s) \overrightarrow{m}), Q)\{\overrightarrow{f}\} \qquad \rightarrow_{\iota} \qquad (C_i[I_s, f_i] \overrightarrow{m})$