

# Propagators: An Introduction

George Wilson

Data61/CSIRO


[george.wilson@data61.csiro.au](mailto:george.wilson@data61.csiro.au)

November 3, 2017





What?




Why?

Roots as early as the 1970's at MIT

- Guy L. Steele Jr.
- Gerald J. Sussman
- Richard Stallman

More recently:


- Alexey Radul



```
(define (map f xs)
  (cond ((null? xs) '())
        (else (cons (f (car xs))
                      (map f (cdr xs)))))))
```

And then

- Edward Kmett



$$y \implies f(x) \leq f(y)$$

# Propagators


The *propagator model* is a model of computation


We model computations as *propagator networks*


A propagator network comprises


- cells
- propagators
- connections between cells and propagators











































$$z \leftarrow x + y$$

$$z = x + y$$




$$7 = x + 4$$


$$z = x + y$$


$$z \leftarrow x + y$$


$$x \leftarrow z - y$$

$$y \leftarrow z - x$$











Propagators let us express multidirectional relationships!




$$^{\circ}F = ^{\circ}C \times \frac{9}{5} + 32$$




$$^{\circ}F = ^{\circ}C \times \frac{9}{5} + 32$$



$$^{\circ}F = ^{\circ}C \times \frac{9}{5} + 32$$




$$^{\circ}F = ^{\circ}C \times \frac{9}{5} + 32$$




$$^{\circ}F = ^{\circ}C \times \frac{9}{5} + 32$$

$$^{\circ}C = (^{\circ}F - 32) \div \frac{9}{5}$$




$$^{\circ}F = ^{\circ}C \times \frac{9}{5} + 32$$

$$^{\circ}C = (^{\circ}F - 32) \div \frac{9}{5}$$




$$^{\circ}F = ^{\circ}C \times \frac{9}{5} + 32$$

$$^{\circ}C = (^{\circ}F - 32) \div \frac{9}{5}$$



$$^{\circ}F = ^{\circ}C \times \frac{9}{5} + 32$$


$$^{\circ}C = (^{\circ}F - 32) \div \frac{9}{5}$$






$$^{\circ}F = ^{\circ}C \times \frac{9}{5} + 32$$

$$^{\circ}C = (^{\circ}F - 32) \div \frac{9}{5}$$




$$^{\circ}F = ^{\circ}C \times \frac{9}{5} + 32$$


$$^{\circ}C = (^{\circ}F - 32) \div \frac{9}{5}$$





$$^{\circ}F = ^{\circ}C \times \frac{9}{5} + 32$$


$$^{\circ}C = (^{\circ}F - 32) \div \frac{9}{5}$$


















What types are the values of the cells?

'c'





```
data Maybe a = Nothing | Just a
```

Partial information!

?

$$[1, 5]$$



TODO interval arithmetic examples

$\{True, False\}$

TODO set intersection examples