

AN EXAMPLE ARTICLE*

DIANNE DOE[†], PAUL T. FRANK[‡], AND JANE E. SMITH[‡]

Abstract. This is an example SIAM L^AT_EX article. This can be used as a template for new articles. Abstracts must be able to stand alone and so cannot contain citations to the paper's references, equations, etc. An abstract must consist of a single paragraph and be concise. Because of online formatting, abstracts must appear as plain as possible. Any equations should be inline.

Key words. example, L^AT_EX

MSC codes. 68Q25, 68R10, 68U05

1. Introduction. The introduction introduces the context and summarizes the manuscript. It is importantly to clearly state the contributions of this piece of work.

For $\Omega = (0, 1)$, $1 < \alpha < 2$, suppose $f \in C^2(\Omega)$

$$(1.1) \quad \begin{cases} (-\Delta)^{\frac{\alpha}{2}} u(x) = f(x), & x \in \Omega \\ u(x) = 0, & x \in \mathbb{R} \setminus \Omega \end{cases}$$

where

$$(1.2) \quad (-\Delta)^{\frac{\alpha}{2}} u(x) = -\frac{\partial^\alpha u}{\partial |x|^\alpha} = -\kappa_\alpha \frac{d^2}{dx^2} \int_\Omega \frac{u(y)}{|x-y|^{\alpha-1}} dy$$

$$(1.3) \quad \kappa_\alpha = -\frac{1}{2 \cos(\alpha\pi/2) \Gamma(2-\alpha)} > 0$$

The paper is organized as follows. Our main results are in section 3, experimental results are in section 5, and the conclusions follow in section 7.

2. Numeric Format.

$$(2.1) \quad x_i = \begin{cases} \frac{1}{2} \left(\frac{i}{N} \right)^r, & 0 \leq i \leq N \\ 1 - \frac{1}{2} \left(\frac{2N-i}{N} \right)^r, & N \leq i \leq 2N \end{cases}$$

where $r \geq 1$. And let

$$(2.2) \quad h_j = x_j - x_{j-1}, \quad 1 \leq j \leq 2N$$

Let $\{\phi_j(x)\}_{j=1}^{2N-1}$ be standard hat functions, which are basis of the piecewise linear function space.

$$(2.3) \quad \phi_j(x) = \begin{cases} \frac{1}{h_j}(x - x_{j-1}), & x_{j-1} \leq x \leq x_j \\ \frac{1}{h_{j+1}}(x_{j+1} - x), & x_j \leq x \leq x_{j+1} \\ 0, & \text{otherwise} \end{cases}$$

*Submitted to the editors DATE.

Funding: This work was funded by the Fog Research Institute under contract no. FRI-454.

[†]Imagination Corp., Chicago, IL (ddoe@imag.com, <http://www.imag.com/~ddoe/>).

[‡]Department of Applied Mathematics, Fictional University, Boise, ID (ptfrank@fictional.edu, jesmith@fictional.edu).

28 And then, we can approximate $u(x)$ with

$$29 \quad (2.4) \quad u_h(x) := \sum_{j=1}^{2N-1} u(x_j) \phi_j(x)$$

30

31 For convience, we denote

$$32 \quad (2.5) \quad I_h(x) := \int_{\Omega} |x - y|^{1-\alpha} u_h(y) dy$$

33 And now, we can approximate the operator (1.2) at x_i with
(2.6)

$$34 \quad D_h^\alpha u_h(x_i) := -\kappa_\alpha \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} I_h(x_{i-1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) I_h(x_i) + \frac{1}{h_{i+1}} I_h(x_{i+1}) \right)$$

35 Finally, we approximate the equation (1.1) with

$$36 \quad (2.7) \quad D_h^\alpha u_h(x_i) = f(x_i), \quad 1 \leq i \leq 2N - 1$$

37

38 The discrete equation (2.7) can be written in matrix form

$$39 \quad (2.8) \quad AU = F$$

40 where U is unknown, $F = (f(x_1), \dots, f(x_{2N-1}))$. The matrix A is constructed as
41 follows: Since

$$\begin{aligned} (2.9) \quad I_h(x_i) &= \int_{\Omega} |x_i - y|^{1-\alpha} u_h(y) dy = \sum_{j=1}^{2N-1} \int_{\Omega} |x_i - y|^{1-\alpha} u(x_j) \phi_j(y) dy \\ &= \sum_{j=1}^{2N-1} u(x_j) \int_{x_{j-1}}^{x_{j+1}} |x_i - y|^{1-\alpha} \phi_j(y) dy \\ 42 \quad &= \sum_{j=1}^{2N-1} \frac{u(x_j)}{(2-\alpha)(3-\alpha)} \left(\frac{|x_i - x_{j-1}|^{3-\alpha}}{h_j} - \frac{h_j + h_{j+1}}{h_j h_{j+1}} |x_i - x_j|^{3-\alpha} + \frac{|x_i - x_{j+1}|^{3-\alpha}}{h_{j+1}} \right) \\ &=: \sum_{j=1}^{2N-1} \tilde{a}_{ij} u(x_j), \quad 0 \leq i \leq 2N \end{aligned}$$

43 Then, substitute in (2.6), we have

$$44 \quad (2.10) \quad D_h^\alpha u_h(x_i) = \sum_{j=1}^{2N-1} a_{ij} u(x_j)$$

45 where

$$46 \quad (2.11) \quad a_{ij} = -\kappa_\alpha \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} \tilde{a}_{i-1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) \tilde{a}_{i,j} + \frac{1}{h_{i+1}} \tilde{a}_{i+1,j} \right)$$

47

3. Main results. Here we state our main results; the proof is deferred to ??.

Let's denote $h = \frac{1}{N}$, we have

THEOREM 3.1 (Truncation Error). *If $f \in C^2(\Omega)$ and $\alpha \in (1, 2)$, and $u(x)$ is a solution of the equation (1.1), then there exists a constant $C = C(\alpha, r, \|f\|_{C^2(\Omega)})$, such that the truncation error of the discrete format satisfies*

$$\begin{aligned} |D_h^\alpha u_h(x_i) - f(x_i)| \leq & C(h^{r\alpha/2+r}(x_i^{-1-\alpha} + (1-x_i)^{-1-\alpha})) \\ & + h^2(x_i^{-\alpha/2-2/r} + (1-x_i)^{-\alpha/2-2/r}) \\ & + h^2 \begin{cases} |\frac{1}{2} - x_{i-1}|^{1-\alpha}, & 1 \leq i \leq N \\ |\frac{1}{2} - x_{i+1}|^{1-\alpha}, & N < i \leq 2N-1 \end{cases} \end{aligned}$$

THEOREM 3.2 (Convergence). *The discrete equation (2.7) has solution U , and there exists a positive constant $C = C(\alpha, r, \|f\|_{C^2(\Omega)})$ such that the error between the numerical solution U with the exact solution $u(x_i)$ satisfies*

$$(3.2) \quad \max_{1 \leq i \leq 2N-1} |U_i - u(x_i)| \leq Ch^{\min\{\frac{r\alpha}{2}, 2\}}$$

That means the numerical method has convergence order $\min\{\frac{r\alpha}{2}, 2\}$.

4. Algorithm. Sed gravida lectus ut purus. Morbi laoreet magna. Pellentesque eu wisi. Proin turpis. Integer sollicitudin augue nec dui. Fusce lectus. Vivamus faucibus nulla nec lacus. Integer diam. Pellentesque sodales, enim feugiat cursus volutpat, sem mauris dignissim mauris, quis consequat sem est fermentum ligula. Nullam justo lectus, condimentum sit amet, posuere a, fringilla mollis, felis. Morbi nulla nibh, pellentesque at, nonummy eu, sollicitudin nec, ipsum. Cras neque. Nunc augue. Nullam vitae quam id quam pulvinar blandit. Nunc sit amet orci. Aliquam erat elit, pharetra nec, aliquet a, gravida in, mi. Quisque urna enim, viverra quis, suscipit quis, tincidunt ut, sapien. Cras placerat consequat sem. Curabitur ac diam. Curabitur diam tortor, mollis et, viverra ac, tempus vel, metus.

Our analysis leads to the algorithm in Algorithm 4.1.

Algorithm 4.1 Build tree

```

Define  $P := T := \{\{1\}, \dots, \{d\}\}$ 
while  $\#P > 1$  do
  Choose  $C' \in \mathcal{C}_p(P)$  with  $C' := \operatorname{argmin}_{C \in \mathcal{C}_p(P)} \varrho(C)$ 
  Find an optimal partition tree  $T_{C'}$ 
  Update  $P := (P \setminus C') \cup \{\bigcup_{t \in C'} t\}$ 
  Update  $T := T \cup \{\bigcup_{t \in \tau} t : \tau \in T_{C'} \setminus \mathcal{L}(T_{C'})\}$ 
end while
return  $T$ 

```

Curabitur ac lorem. Vivamus non justo in dui mattis posuere. Etiam accumsan ligula id pede. Maecenas tincidunt diam nec velit. Praesent convallis sapien ac est. Aliquam ullamcorper euismod nulla. Integer mollis enim vel tortor. Nulla sodales placerat nunc. Sed tempus rutrum wisi. Duis accumsan gravida purus. Nunc nunc. Etiam facilisis dui eu sem. Vestibulum semper. Praesent eu eros. Vestibulum tellus nisl, dapibus id, vestibulum sit amet, placerat ac, mauris. Maecenas et elit ut erat placerat dictum. Nam feugiat, turpis et sodales volutpat, wisi quam rhoncus neque, vitae aliquam ipsum sapien vel enim. Maecenas suscipit cursus mi.

5. Experimental results. Quisque facilisis auctor sapien. Pellentesque gravida hendrerit lectus. Mauris rutrum sodales sapien. Fusce hendrerit sem vel lorem. Integer pellentesque massa vel augue. Integer elit tortor, feugiat quis, sagittis et, ornare non, lacus. Vestibulum posuere pellentesque eros. Quisque venenatis ipsum dictum nulla. Aliquam quis quam non metus eleifend interdum. Nam eget sapien ac mauris malesuada adipiscing. Etiam eleifend neque sed quam. Nulla facilisi. Proin a ligula. Sed id dui eu nibh egestas tincidunt. Suspendisse arcu.

Figure 1 shows some example results. Additional results are available in the supplement in Table 1.




FIG. 1. Example figure using external image files.

Table 1 shows additional supporting evidence.

TABLE 1
Example table.

Species	Mean	Std. Dev.
1	3.4	1.2
2	5.4	0.6
3	7.4	2.4
4	9.4	1.8

Maecenas dui. Aliquam volutpat auctor lorem. Cras placerat est vitae lectus. Curabitur massa lectus, rutrum euismod, dignissim ut, dapibus a, odio. Ut eros erat, vulputate ut, interdum non, porta eu, erat. Cras fermentum, felis in porta congue, velit leo facilisis odio, vitae consectetur lorem quam vitae orci. Sed ultrices, pede eu placerat auctor, ante ligula rutrum tellus, vel posuere nibh lacus nec nibh. Maecenas laoreet dolor at enim. Donec molestie dolor nec metus. Vestibulum libero. Sed quis erat. Sed tristique. Duis pede leo, fermentum quis, consectetur eget, vulputate sit amet, erat.

6. Discussion of $Z = X \cup Y$. Curabitur nunc magna, posuere eget, venenatis eu, vehicula ac, velit. Aenean ornare, massa a accumsan pulvinar, quam lorem laoreet purus, eu sodales magna risus molestie lorem. Nunc erat velit, hendrerit quis, malesuada ut, aliquam vitae, wisi. Sed posuere. Suspendisse ipsum arcu, scelerisque nec, aliquam eu, molestie tincidunt, justo. Phasellus iaculis. Sed posuere lorem non ipsum. Pellentesque dapibus. Suspendisse quam libero, laoreet a, tincidunt eget, consequat at, est. Nullam ut lectus non enim consequat facilisis. Mauris leo. Quisque pede ligula, auctor vel, pellentesque vel, posuere id, turpis. Cras ipsum sem, cursus et, facilisis ut, tempus euismod, quam. Suspendisse tristique dolor eu orci. Mauris

106 mattis. Aenean semper. Vivamus tortor magna, facilisis id, varius mattis, hendrerit
 107 in, justo. Integer purus.

108 **7. Conclusions.** Some conclusions here.

109 **Appendix A. An example appendix.** Aenean tincidunt laoreet dui. Vestibu-
 110 lum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Integer
 111 ipsum lectus, fermentum ac, malesuada in, eleifend ut, lorem. Vivamus ipsum turpis,
 112 elementum vel, hendrerit ut, semper at, metus. Vivamus sapien tortor, eleifend id,
 113 dapibus in, egestas et, pede. Pellentesque faucibus. Praesent lorem neque, dignissim
 114 in, facilisis nec, hendrerit vel, odio. Nam at diam ac neque aliquet viverra. Morbi da-
 115 pibus ligula sagittis magna. In lobortis. Donec aliquet ultricies libero. Nunc dictum
 116 vulputate purus. Morbi varius. Lorem ipsum dolor sit amet, consectetur adipiscing
 117 elit. In tempor. Phasellus commodo porttitor magna. Curabitur vehicula odio vel
 118 dolor.

119 **LEMMA A.1.** *Test Lemma.*

120 **Acknowledgments.** We would like to acknowledge the assistance of volunteers
 121 in putting together this example manuscript and supplement.