

AN EXAMPLE ARTICLE*

DIANNE DOE[†], PAUL T. FRANK[‡], AND JANE E. SMITH[‡]

Abstract. This is an example SIAM L^AT_EX article. This can be used as a template for new articles. Abstracts must be able to stand alone and so cannot contain citations to the paper's references, equations, etc. An abstract must consist of a single paragraph and be concise. Because of online formatting, abstracts must appear as plain as possible. Any equations should be inline.

Key words. example, L^AT_EX

MSC codes. 68Q25, 68R10, 68U05

1. Introduction. The introduction introduces the context and summarizes the manuscript. It is importantly to clearly state the contributions of this piece of work.

For $\Omega = (0, 2T)$, $1 < \alpha < 2$, suppose $f \in C^\beta(\Omega) \cap L^\infty(\Omega)$, $\beta > 4 - \alpha$, $\|f\|_\beta^{\alpha/2} < \infty$

$$(1.1) \quad \begin{cases} (-\Delta)^{\frac{\alpha}{2}} u(x) = f(x), & x \in \Omega \\ u(x) = 0, & x \in \mathbb{R} \setminus \Omega \end{cases}$$

where

$$(1.2) \quad (-\Delta)^{\frac{\alpha}{2}} u(x) = -\frac{\partial^\alpha u}{\partial |x|^\alpha} = -\kappa_\alpha \frac{d^2}{dx^2} \int_\Omega \frac{|x-y|^{1-\alpha}}{\Gamma(2-\alpha)} u(y) dy$$

$$(1.3) \quad \kappa_\alpha = -\frac{1}{2 \cos(\alpha\pi/2)} > 0$$

2. Regularity.

Remark 2.1. 1. $C^k(U)$ is the set of all k -times continuously differentiable functions on open set U .

2. $C^\beta(U)$ is the collection of function f which for any $V \subset\subset U$ $f|_V \in C^\beta(\bar{V})$.

THEOREM 2.2. If $f \in C^\beta(\Omega)$, $\beta > 2$ and $\|f\|_\beta^{(\alpha/2)} < \infty$, then for $l = 0, 1, 2$

$$(2.1) \quad |f^{(l)}(x)| \leq \|f\|_\beta^{(\alpha/2)} \begin{cases} x^{-l-\alpha/2}, & \text{if } 0 < x \leq T \\ (2T-x)^{-l-\alpha/2}, & \text{if } T \leq x < 2T \end{cases}$$

THEOREM 2.3 (Regularity up to the boundary [1]).

$$(2.2) \quad \|u\|_{\beta+\alpha}^{(-\alpha/2)} \leq C \left(\|u\|_{C^{\alpha/2}(\mathbb{R})} + \|f\|_\beta^{(\alpha/2)} \right)$$

*Submitted to the editors DATE.

Funding: This work was funded by the Fog Research Institute under contract no. FRI-454.

[†]Imagination Corp., Chicago, IL (ddoe@imag.com, <http://www.imag.com/~ddoe/>).

[‡]Department of Applied Mathematics, Fictional University, Boise, ID (ptfrank@fictional.edu, jesmith@fictional.edu).

28 **COROLLARY 2.4.** *Let u be a solution of (1.1) on Ω . Then, for any $x \in \Omega$ and*
 29 *$l = 0, 1, 2, 3, 4$*

$$30 \quad (2.3) \quad |u^{(l)}(x)| \leq C \begin{cases} x^{\alpha/2-l}, & \text{if } 0 < x \leq T \\ (2T-x)^{\alpha/2-l}, & \text{if } T \leq x < 2T \end{cases}$$

31 The paper is organized as follows. Our main results are in section 4, experimental
 32 results are in section 7, and the conclusions follow in section 8.

3. Numeric Format.

$$33 \quad (3.1) \quad x_i = \begin{cases} T \left(\frac{i}{N} \right)^r, & 0 \leq i \leq N \\ 2T - T \left(\frac{2N-i}{N} \right)^r, & N \leq i \leq 2N \end{cases}$$

34 where $r \geq 1$. And let

$$35 \quad (3.2) \quad h_j = x_j - x_{j-1}, \quad 1 \leq j \leq 2N$$

36 Let $\{\phi_j(x)\}_{j=1}^{2N-1}$ be standard hat functions, which are basis of the piecewise linear
 37 function space.

$$38 \quad (3.3) \quad \phi_j(x) = \begin{cases} \frac{1}{h_j}(x - x_{j-1}), & x_{j-1} \leq x \leq x_j \\ \frac{1}{h_{j+1}}(x_{j+1} - x), & x_j \leq x \leq x_{j+1} \\ 0, & \text{otherwise} \end{cases}$$

39 And then, we can approximate $u(x)$ with

$$40 \quad (3.4) \quad u_h(x) := \sum_{j=1}^{2N-1} u(x_j) \phi_j(x)$$

41 For convience, we denote

$$42 \quad (3.5) \quad I_h^{2-\alpha}(x_i) := \frac{1}{\Gamma(2-\alpha)} \int_{\Omega} |x_i - y|^{1-\alpha} u_h(y) dy$$

43 And now, we can approximate the operator (1.2) at x_i with

$$44 \quad (3.6) \quad \begin{aligned} D_h^\alpha u_h(x_i) &:= D_h^2 I_h^{2-\alpha}(x_i) \\ &= \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} I_h^{2-\alpha}(x_{i-1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) I_h^{2-\alpha}(x_i) + \frac{1}{h_{i+1}} I_h^{2-\alpha}(x_{i+1}) \right) \end{aligned}$$

45 Finally, we approximate the equation (1.1) with

$$46 \quad (3.7) \quad -\kappa_\alpha D_h^\alpha u_h(x_i) = f(x_i), \quad 1 \leq i \leq 2N-1$$

47 The discrete equation (3.7) can be written in matrix form

$$48 \quad (3.8) \quad AU = F$$

where U is unknown, $F = (f(x_1), \dots, f(x_{2N-1}))$. The matrix A is constructed as follows: Since

(3.9)

$$\begin{aligned}
 I_h^{2-\alpha}(x_i) &= \frac{1}{\Gamma(2-\alpha)} \int_{\Omega} |x_i - y|^{1-\alpha} u_h(y) dy \\
 &= \sum_{j=1}^{2N-1} \frac{1}{\Gamma(2-\alpha)} \int_{\Omega} |x_i - y|^{1-\alpha} u(x_j) \phi_j(y) dy \\
 &= \sum_{j=1}^{2N-1} u(x_j) \frac{1}{\Gamma(2-\alpha)} \int_{x_{j-1}}^{x_{j+1}} |x_i - y|^{1-\alpha} \phi_j(y) dy \\
 &= \sum_{j=1}^{2N-1} \frac{u(x_j)}{\Gamma(4-\alpha)} \left(\frac{|x_i - x_{j-1}|^{3-\alpha}}{h_j} - \frac{h_j + h_{j+1}}{h_j h_{j+1}} |x_i - x_j|^{3-\alpha} + \frac{|x_i - x_{j+1}|^{3-\alpha}}{h_{j+1}} \right) \\
 &=: \sum_{j=1}^{2N-1} \tilde{a}_{ij} u(x_j), \quad 0 \leq i \leq 2N
 \end{aligned}$$

Then, substitute in (3.6), we have

$$(3.10) \quad -\kappa_{\alpha} D_h^{\alpha} u_h(x_i) = \sum_{j=1}^{2N-1} a_{ij} u(x_j)$$

where

$$(3.11) \quad a_{ij} = -\kappa_{\alpha} \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} \tilde{a}_{i-1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) \tilde{a}_{i,j} + \frac{1}{h_{i+1}} \tilde{a}_{i+1,j} \right)$$

4. Main results. Here we state our main results; the proof is deferred to section 5 and section 6.

Let's denote $h = \frac{1}{N}$, we have

THEOREM 4.1 (Truncation Error). *If $f \in C^2(\Omega)$ and $\alpha \in (1, 2)$, and $u(x)$ is a solution of the equation (1.1), then there exists a constant $C_1, C_2 = C_1(T, \alpha, r, \|f\|_{C^2(\Omega)}), C_2(T, \alpha, r, \|f\|_{C^2(\Omega)})$, such that the truncation error of the discrete format satisfies*

$$\begin{aligned}
 |-\kappa_{\alpha} D_h^{\alpha} u_h(x_i) - f(x_i)| &\leq C_1 (h^{r\alpha/2+r} (x_i^{-1-\alpha} + (2T - x_i)^{-1-\alpha}) \\
 &\quad + h^2 (x_i^{-\alpha/2-2/r} + (2T - x_i)^{-\alpha/2-2/r})) \\
 &\quad + C_2 h^2 \begin{cases} |T - x_{i-1}|^{1-\alpha}, & 1 \leq i \leq N \\ |T - x_{i+1}|^{1-\alpha}, & N < i \leq 2N - 1 \end{cases}
 \end{aligned}$$

where $C_2 = 0$ if $r = 1$.

THEOREM 4.2 (Convergence). *The discrete equation (3.7) has solution U , and there exists a positive constant $C = C(T, \alpha, r, \|f\|_{C^2(\Omega)})$ such that the error between the numerical solution U with the exact solution $u(x_i)$ satisfies*

$$(4.2) \quad \max_{1 \leq i \leq 2N-1} |U_i - u(x_i)| \leq Ch^{\min\{\frac{r\alpha}{2}, 2\}}$$

That means the numerical method has convergence order $\min\{\frac{r\alpha}{2}, 2\}$.

5. Proof of Theorem 4.1. For convience, let's denote

$$(5.1) \quad I^{2-\alpha}(x) = \frac{1}{\Gamma(2-\alpha)} \int_{\Omega} |x-y|^{1-\alpha} u(y) dy$$

Then, the truncation error of the discrete format can be written as

$$(5.2) \quad \begin{aligned} -\kappa_{\alpha} D_h^{\alpha} u_h(x_i) - f(x_i) &= -\kappa_{\alpha} (D_h^2 I_h^{2-\alpha}(x_i) - \frac{d^2}{dx^2} I^{2-\alpha}(x_i)) \\ &= -\kappa_{\alpha} D_h^2 (I_h^{2-\alpha} - I^{2-\alpha})(x_i) - \kappa_{\alpha} (D_h^2 - \frac{d^2}{dx^2}) I^{2-\alpha}(x_i) \end{aligned}$$

5.1. Estimate of $-\kappa_{\alpha} (D_h^2 - \frac{d^2}{dx^2}) I^{2-\alpha}(x_i)$.

THEOREM 5.1. *There exists a constant $C = C(T, \alpha, r, \|f\|_{\beta}^{(\alpha/2)})$ such that*

$$(5.3) \quad \left| -\kappa_{\alpha} (D_h^2 - \frac{d^2}{dx^2}) I^{2-\alpha}(x_i) \right| \leq Ch^2 (x_i^{-\alpha/2-2/r} + (2T-x_i)^{-\alpha/2-2/r})$$

Proof. Since $f \in C^2(\Omega)$ and

$$(5.4) \quad \frac{d^2}{dx^2} (-\kappa_{\alpha} I^{2-\alpha}(x)) = f(x), \quad x \in \Omega,$$

we have $I^{2-\alpha} \in C^4(\Omega)$. Therefore, using equation (A.3) of Lemma A.1, for $1 \leq i \leq 2N-1$, we have

$$(5.5) \quad -\kappa_{\alpha} (D_h^2 - \frac{d^2}{dx^2}) I^{2-\alpha}(x_i) = \frac{h_{i+1} - h_i}{3} f'(x_i) + \frac{1}{4!} \frac{2}{h_i + h_{i+1}} (h_i^3 f''(\eta_1) + h_{i+1}^3 f''(\eta_2))$$

where $\eta_1 \in [x_{i-1}, x_i]$, $\eta_2 \in [x_i, x_{i+1}]$. By Lemma B.2 and Theorem 2.2 we have 1.

$$(5.6) \quad \left| \frac{h_{i+1} - h_i}{3} f'(x_i) \right| \leq \frac{\|f\|_{\beta}^{(\alpha/2)}}{3} Ch^2 \begin{cases} x_i^{-\alpha/2-2/r}, & 1 \leq i \leq N-1 \\ 0, & i = N \\ (2T-x_i)^{-\alpha/2-2/r}, & N < i \leq 2N-1 \end{cases}$$

2. See Proof 7, there is a constant $C = C(T, \alpha, r, \|f\|_{\beta}^{(\alpha/2)})$ such that

$$(5.7) \quad \begin{aligned} &\left| \frac{1}{4!} \frac{2}{h_i + h_{i+1}} (h_i^3 f''(\eta_1) + h_{i+1}^3 f''(\eta_2)) \right| \\ &\leq Ch^2 \begin{cases} x_i^{-\alpha/2-2/r}, & 1 \leq i \leq N \\ (2T-x_i)^{-\alpha/2-2/r}, & N \leq i \leq 2N-1 \end{cases} \end{aligned}$$

Summarizes, we get the result. \square

5.2. Estimate of R_i . Now, we study the first part of (5.2)

$$(5.8) \quad D_h^2 (I^{2-\alpha} - I_h^{2-\alpha})(x_i) = D_h^2 \left(\int_0^{2T} (u(y) - u_h(y)) \frac{|y-x_i|^{1-\alpha}}{\Gamma(2-\alpha)} dy \right)$$

For convience, let's denote

$$(5.9) \quad T_{ij} = \int_{x_{j-1}}^{x_j} (u(y) - u_h(y)) \frac{|y-x_i|^{1-\alpha}}{\Gamma(2-\alpha)} dy$$

91 And define

$$92 \quad (5.10) \quad \begin{aligned} R_i &:= D_h^2(I^{2-\alpha} - I_h^{2-\alpha})(x_i) \\ &= \frac{2}{h_i + h_{i+1}} \sum_{j=1}^{2N} \left(\frac{1}{h_i} T_{i-1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_{i+1}} T_{i+1,j} \right) \end{aligned}$$

93 We have some results about the estimate of R_i

94 **THEOREM 5.2.** *For $1 \leq i < N/2$, there exists a constant C such that*

$$95 \quad (5.11) \quad R_i \leq C(h^{r\alpha/2+r} x_i^{-1-\alpha} + h^2 x_i^{-\alpha/2-2/r})$$

96

97 **THEOREM 5.3.** *For $N/2 \leq i \leq N$, there exists constant C, C_2 such that*

$$98 \quad (5.12) \quad R_i \leq C h^2 x_i^{-\alpha/2-2/r} + C_2 h^2 |T - x_{i-1}|^{1-\alpha}$$

99 *where $C_2 = 0$ if $r = 1$.*

100 And for $N < i \leq 2N - 1$, it is symmetric to the previous case.

101 To prove these results, we need some utils. Also for simplicity, we denote

$$102 \quad (5.13) \quad S_{ij} = \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} T_{i-1,j} - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) T_{i,j} + \frac{1}{h_{i+1}} T_{i+1,j} \right)$$

103 then

$$104 \quad (5.14) \quad R_i = \sum_{j=1}^{2N} S_{ij}$$

105 **LEMMA 5.4.** *While $0 \leq i < N/2$, By Lemma A.3*

$$106 \quad (5.15) \quad \begin{aligned} |T_{i1}| &\leq C \int_0^{x_1} x_1^{\alpha/2} \frac{|x_i - y|^{1-\alpha}}{\Gamma(2-\alpha)} dy \\ &= C \frac{1}{\Gamma(3-\alpha)} x_1^{\alpha/2} |x_i^{2-\alpha} - |x_i - x_1|^{2-\alpha}| \\ &\leq C \frac{1}{\Gamma(3-\alpha)} x_1^{\alpha/2+2-\alpha} = C \frac{1}{\Gamma(3-\alpha)} x_1^{2-\alpha/2} \quad 0 < 2-\alpha < 1 \end{aligned}$$

107 *For $2 \leq j \leq N$, by Lemma A.2*

$$108 \quad (5.16) \quad \begin{aligned} |T_{ij}| &\leq \frac{C}{4} \int_{x_{j-1}}^{x_j} h_j^2 x_{j-1}^{\alpha/2-2} \frac{|y - x_i|^{1-\alpha}}{\Gamma(2-\alpha)} dy \\ &\leq \frac{C}{4\Gamma(3-\alpha)} h_j^2 x_{j-1}^{\alpha/2-2} ||x_j - x_i|^{2-\alpha} - |x_{j-1} - x_i|^{2-\alpha}| \end{aligned}$$

109

110 **LEMMA 5.5.** *While $1 \leq i < N/2$,*

$$111 \quad (5.17) \quad \sum_{j=\max\{2i+1, i+3\}}^N S_{ij} \leq C h^2 x_i^{-\alpha/2-2/r}$$

Proof. For $\max\{2i+1, i+3\} \leq j \leq N$, by Lemma C.1 and Lemma C.2

$$\begin{aligned} S_{ij} &= \int_{x_{j-1}}^{x_j} (u(y) - u_h(y)) D_h^2 \left(\frac{|y - \cdot|^{1-\alpha}}{\Gamma(2-\alpha)} \right) (x_i) dy \\ &\leq Ch^2 \int_{x_{j-1}}^{x_j} y^{\alpha/2-2/r} \frac{y^{-1-\alpha}}{\Gamma(-\alpha)} dy \\ &= Ch^2 \int_{x_{j-1}}^{x_j} y^{-\alpha/2-2/r-1} dy \end{aligned}$$

Therefore,

$$\begin{aligned} \sum_{j=\max\{2i+1, i+3\}}^N S_{ij} &\leq Ch^2 \int_{x_{2i}}^{x_N} y^{-\alpha/2-2/r-1} dy \\ &= \frac{C}{\alpha/2+2/r} h^2 (2^{r(-\alpha/2-2/r)} x_i^{-\alpha/2-2/r} - T^{-\alpha/2-2/r}) \end{aligned}$$

□

6. Proof of Theorem 4.2.

7. Experimental results.

8. Conclusions. Some conclusions here.

Appendix A. Approximate of difference quotients.

LEMMA A.1. *If $g(x)$ is twice differentiable continuous function on open set Ω , there exists $\xi \in [x_{i-1}, x_{i+1}]$ such that*

$$\begin{aligned} D_h^2 g(x_i) &:= \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g(x_i) + \frac{1}{h_i} g(x_{i-1}) \right) \\ &= g''(\xi), \quad \xi \in [x_{i-1}, x_{i+1}] \end{aligned}$$

$$\begin{aligned} &\frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g(x_i) + \frac{1}{h_i} g(x_{i-1}) \right) \\ &= \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_i} \int_{x_{i-1}}^{x_i} g''(y) (y - x_{i-1}) dy + \frac{1}{h_{i+1}} \int_{x_i}^{x_{i+1}} g''(y) (x_{i+1} - y) dy \right) \end{aligned}$$

And if $g(x) \in C^4(\Omega)$, then

$$\begin{aligned} &\frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g(x_i) + \frac{1}{h_i} g(x_{i-1}) \right) \\ &= g''(x_i) + \frac{h_{i+1} - h_i}{3} g'''(x_i) + \frac{1}{4!} \frac{2}{h_i + h_{i+1}} (h_i^3 g''''(\eta_1) + h_{i+1}^3 g''''(\eta_2)) \end{aligned}$$

where $\eta_1 \in [x_{i-1}, x_i]$, $\eta_2 \in [x_i, x_{i+1}]$.

Proof.

$$g(x_{i-1}) = g(x_i) - (x_i - x_{i-1}) g'(x_i) + \frac{(x_i - x_{i-1})^2}{2} g''(\xi_1), \quad \xi_1 \in [x_{i-1}, x_i]$$

$$g(x_{i+1}) = g(x_i) + (x_{i+1} - x_i) g'(x_i) + \frac{(x_{i+1} - x_i)^2}{2} g''(\xi_2), \quad \xi_2 \in [x_i, x_{i+1}]$$

130 Subsitute them in the left side of (A.1), we have

$$131 \quad \frac{2}{h_i + h_{i+1}} \left(\frac{1}{h_{i+1}} g(x_{i+1}) - \left(\frac{1}{h_i} + \frac{1}{h_{i+1}} \right) g(x_i) + \frac{1}{h_i} g(x_{i-1}) \right) \\ = \frac{h_i}{h_i + h_{i+1}} g''(\xi_1) + \frac{h_{i+1}}{h_i + h_{i+1}} g''(\xi_2)$$

132 Now, using intermediate value theorem , there exists $\xi \in [\xi_1, \xi_2]$ such that

$$133 \quad \frac{h_i}{h_i + h_{i+1}} g''(\xi_1) + \frac{h_{i+1}}{h_i + h_{i+1}} g''(\xi_2) = g''(\xi)$$

134 For the second equation, similarly

$$135 \quad g(x_{i-1}) = g(x_i) - (x_i - x_{i-1})g'(x_i) + \int_{x_{i-1}}^{x_i} g''(y)(y - x_{i-1})dy \\ 136 \quad g(x_{i+1}) = g(x_i) + (x_{i+1} - x_i)g'(x_i) + \int_{x_i}^{x_{i+1}} g''(y)(x_{i+1} - y)dy$$

137 And the last equation can be obtained by

$$138 \quad g(x_{i-1}) = g(x_i) - h_i g'(x_i) + \frac{h_i^2}{2} g''(x_i) - \frac{h_i^3}{3!} g'''(x_i) + \frac{h_i^4}{4!} g''''(\eta_1) \\ 139 \quad g(x_{i+1}) = g(x_i) + h_{i+1} g'(x_i) + \frac{h_{i+1}^2}{2} g''(x_i) + \frac{h_{i+1}^3}{3!} g'''(x_i) + \frac{h_{i+1}^4}{4!} g''''(\eta_2)$$

140 where $\eta_1 \in [x_{i-1}, x_i]$, $\eta_2 \in [x_i, x_{i+1}]$. Expecially,

$$141 \quad (A.4) \quad \frac{h_i^4}{4!} g''''(\eta_1) = \int_{x_{i-1}}^{x_i} g''''(y) \frac{(y - x_{i-1})^3}{3!} dy \\ \frac{h_{i+1}^4}{4!} g''''(\eta_2) = \int_{x_i}^{x_{i+1}} g''''(y) \frac{(x_{i+1} - y)^3}{3!} dy$$

142 Subsitute them to the left side of (A.3), we can get the result. □

143 LEMMA A.2. If $y \in [x_{j-1}, x_j]$, denote $y = \theta x_{j-1} + (1 - \theta)x_j$, $\theta \in [0, 1]$,

$$144 \quad (A.5) \quad u(y_j^\theta) - u_h(y_j^\theta) = -\frac{\theta(1-\theta)}{2} h_j^2 u''(\xi), \quad \xi \in [x_{j-1}, x_j]$$

$$145 \quad (A.6) \quad u(y_j^\theta) - u_h(y_j^\theta) = -\frac{\theta(1-\theta)}{2} h_j^2 u''(y_j^\theta) + \frac{\theta(1-\theta)}{3!} h_j^3 (\theta^2 u'''(\eta_1) - (1-\theta)^2 u'''(\eta_2))$$

147 where $\eta_1 \in [x_{j-1}, y_j^\theta]$, $\eta_2 \in [y_j^\theta, x_j]$.

148 *Proof.* By Taylor expansion, we have

$$149 \quad u(x_{j-1}) = u(y_j^\theta) - \theta h_j u'(y_j^\theta) + \frac{\theta^2 h_j^2}{2!} u''(\xi_1), \quad \xi_1 \in [x_{j-1}, y_j^\theta] \\ 150 \quad u(x_j) = u(y_j^\theta) + (1-\theta) h_j u'(y_j^\theta) + \frac{(1-\theta)^2 h_j^2}{2!} u''(\xi_2), \quad \xi_2 \in [y_j^\theta, x_j]$$

Thus

$$\begin{aligned} u(y_j^\theta) - u_h(y_j^\theta) &= u(y_j^\theta) - (1 - \theta)u(x_{j-1}) - \theta u(x_j) \\ &= -\frac{\theta(1 - \theta)}{2} h_j^2 (\theta u''(\xi_1) + (1 - \theta)u''(\xi_2)) \\ &= -\frac{\theta(1 - \theta)}{2} h_j^2 u''(\xi), \quad \xi \in [\xi_1, \xi_2] \end{aligned}$$

The second equation is similar,

$$\begin{aligned} u(x_{j-1}) &= u(y_j^\theta) - \theta h_j u'(y_j^\theta) + \frac{\theta^2 h_j^2}{2!} u''(y_j^\theta) - \frac{\theta^3 h_j^3}{3!} u'''(\eta_1) \\ u(x_j) &= u(y_j^\theta) + (1 - \theta) h_j u'(y_j^\theta) + \frac{(1 - \theta)^2 h_j^2}{2!} u''(y_j^\theta) + \frac{(1 - \theta)^3 h_j^3}{3!} u'''(\eta_2) \end{aligned}$$

where $\eta_1 \in [x_{j-1}, y_j^\theta]$, $\eta_2 \in [y_j^\theta, x_j]$. Thus

$$\begin{aligned} u(y_j^\theta) - u_h(y_j^\theta) &= u(y_j^\theta) - (1 - \theta)u(x_{j-1}) - \theta u(x_j) \\ &= -\frac{\theta(1 - \theta)}{2} h_j^2 u''(y_j^\theta) + \frac{\theta(1 - \theta)}{3!} h_j^3 (\theta^2 u'''(\eta_1) - (1 - \theta)^2 u'''(\eta_2)) \end{aligned}$$

LEMMA A.3. For $x \in [x_{j-1}, x_j]$

$$\begin{aligned} |u(x) - u_h(x)| &= \left| \frac{x_j - x}{h_j} \int_{x_{j-1}}^x u'(y) dy - \frac{x - x_{j-1}}{h_j} \int_x^{x_j} u'(y) dy \right| \\ &\leq \int_{x_{j-1}}^{x_j} |u'(y)| dy \end{aligned}$$

If $x \in [0, x_1]$, with Corollary 2.4, we have

$$|u(x) - u_h(x)| \leq \int_0^{x_1} |u'(y)| dy \leq \int_0^{x_1} C y^{\alpha/2-1} dy \leq C \frac{2}{\alpha} x_1^{\alpha/2}$$

Similarly, if $x \in [x_{2N-1}, 1]$, we have

$$|u(x) - u_h(x)| \leq C \frac{2}{\alpha} (2T - x_{2N-1})^{\alpha/2} = C \frac{2}{\alpha} x_1^{\alpha/2}$$

Appendix B. Inequality.

LEMMA B.1.

$$h_i \leq r T^{1/r} h \begin{cases} x_i^{1-1/r}, & 1 \leq i \leq N \\ (2T - x_{i-1})^{1-1/r}, & N < i \leq 2N - 1 \end{cases}$$

Proof. For $1 \leq i \leq N$,

$$\begin{aligned} h_i &= T \left(\left(\frac{i}{N} \right)^r - \left(\frac{i-1}{N} \right)^r \right) \\ &\leq r T \frac{1}{N} \left(\frac{i}{N} \right)^{r-1} = r T^{1/r} h x_i^{1-1/r} \end{aligned}$$

168 For $N < i \leq 2N - 1$,

$$\begin{aligned} h_i &= T \left(\left(\frac{2N-i+1}{N} \right)^r - \left(\frac{2N-i}{N} \right)^r \right) \\ &\leq rT \frac{1}{N} \left(\frac{2N-i+1}{N} \right)^{r-1} = rT^{1/r} h(2T - x_{i-1})^{1-1/r} \end{aligned}$$

169

170

171 LEMMA B.2. *There is a constant $C = 2^{\lfloor r-2 \rfloor} r(r-1)T^{2/r}$ such that for all $i \in$*
 172 *$\{1, 2, \dots, 2N-1\}$*

$$(B.2) \quad |h_{i+1} - h_i| \leq Ch^2 \begin{cases} x_i^{1-2/r}, & 1 \leq i \leq N-1 \\ 0, & i = N \\ (2T - x_i)^{1-2/r}, & N < i \leq 2N-1 \end{cases}$$

Proof.

$$h_{i+1} - h_i = \begin{cases} T \left(\left(\frac{i+1}{N} \right)^r - 2 \left(\frac{i}{N} \right)^r + \left(\frac{i-1}{N} \right)^r \right), & 1 \leq i \leq N-1 \\ 0, & i = N \\ -T \left(\left(\frac{2N-i-1}{N} \right)^r - 2 \left(\frac{2N-i}{N} \right)^r + \left(\frac{2N-i+1}{N} \right)^r \right), & N+1 \leq i \leq 2N-1 \end{cases}$$

175 For $i = 1$,

$$h_2 - h_1 = T(2^r - 2) \left(\frac{1}{N} \right)^r = (2^r - 2)T^{2/r} h^2 x_1^{1-2/r}$$

176 For $2 \leq i \leq N-1$,

$$h_{i+1} - h_i = r(r-1)T N^{-2} \eta^{r-2}, \quad \eta \in \left[\frac{i-1}{N}, \frac{i+1}{N} \right]$$

178 If $r \in [1, 2]$,

$$\begin{aligned} h_{i+1} - h_i &= r(r-1)T N^{-2} \eta^{r-2} \leq r(r-1)T h^2 \left(\frac{i-1}{N} \right)^{r-2} \\ &\leq r(r-1)T h^2 2^{2-r} \left(\frac{i}{N} \right)^{r-2} \\ &= 2^{2-r} r(r-1)T^{2/r} h^2 x_i^{1-2/r} \end{aligned}$$

180

181 else if $r > 2$,

$$\begin{aligned} h_{i+1} - h_i &= r(r-1)T N^{-2} \eta^{r-2} \leq r(r-1)T h^2 \left(\frac{i+1}{N} \right)^{r-2} \\ &\leq r(r-1)T h^2 2^{r-2} \left(\frac{i}{N} \right)^{r-2} \\ &= 2^{r-2} r(r-1)T^{2/r} h^2 x_i^{1-2/r} \end{aligned}$$

182

183 Since

$$2^r - 2 \leq 2^{\lfloor r-2 \rfloor} r(r-1), \quad r \geq 1$$

184

we have

$$h_{i+1} - h_i \leq 2^{|r-2|} r(r-1) T^{2/r} h^2 x_i^{1-2/r}, \quad 1 \leq i \leq N-1$$

For $i = N$, $h_{N+1} - h_N = 0$. For $N < i \leq 2N-1$, it's central symmetric to the first half of the proof, which is

$$h_i - h_{i+1} \leq 2^{|r-2|} r(r-1) T^{2/r} h^2 (2T - x_i)^{1-2/r}$$

Summarizes the inequalities, we can get

$$(B.3) \quad |h_{i+1} - h_i| \leq 2^{|r-2|} r(r-1) T^{2/r} h^2 \begin{cases} x_i^{1-2/r}, & 1 \leq i \leq N-1 \\ 0, & i = N \\ (2T - x_i)^{1-2/r}, & N < i \leq 2N-1 \end{cases} \quad \square$$

Appendix C. Proofs of some technical details.

Additional proof of Theorem 5.1. For $2 \leq i \leq N-1$,

$$\begin{aligned} & \frac{2}{h_i + h_{i+1}} (h_i^3 f''(\eta_1) + h_{i+1}^3 f''(\eta_2)) \\ & \leq C \frac{2}{h_i + h_{i+1}} (h_i^3 x_{i-1}^{-2-\alpha/2} + h_{i+1}^3 x_i^{-2-\alpha/2}) \\ & \leq 2C (h_i^2 x_{i-1}^{-2-\alpha/2} + h_{i+1}^2 x_i^{-2-\alpha/2}) \end{aligned}$$

Since Lemma B.1, we have

$$\begin{aligned} h_i & \leq r T^{1/r} h x_i^{1-1/r}, \quad 1 \leq i \leq N \\ h_{i+1} & \leq r T^{1/r} h x_{i+1}^{1-1/r}, \quad 1 \leq i \leq N-1 \end{aligned}$$

and

$$\begin{aligned} x_{i-1}^{-2-\alpha/2} & \leq 2^{-r(-2-\alpha/2)} x_i^{-2-\alpha/2} \quad 2 \leq i \leq N-1 \\ x_{i+1}^{1-1/r} & \leq 2^{r-1} x_i^{1-1/r} \quad 1 \leq i \leq N-1 \end{aligned}$$

So there is a constant $C = C(T, \alpha, r, \|f\|_{\beta}^{\alpha/2})$ such that

$$\frac{2}{h_i + h_{i+1}} (h_i^3 f''(\eta_1) + h_{i+1}^3 f''(\eta_2)) \leq C h^2 x_i^{-\alpha/2-2/r}, \quad 2 \leq i \leq N-1$$

For $i = 1$, by (A.4)

$$\begin{aligned} & \frac{1}{4!} \frac{2}{h_1 + h_2} (h_1^3 f''(\eta_1) + h_2^3 f''(\eta_2)) \\ & = \frac{2}{h_1 + h_2} \left(\frac{1}{h_1} \int_0^{x_1} f''(y) \frac{y^3}{3!} dy + \frac{1}{4!} h_2^3 f''(\eta_2) \right) \end{aligned}$$

We have proved above that

$$\frac{2}{h_1 + h_2} h_2^3 f''(\eta_2) \leq C h^2 x_1^{-\alpha/2-2/r}$$

207 and we can get

$$\begin{aligned} \int_0^{x_1} f''(y) \frac{y^3}{3!} dy &\leq C \frac{1}{3!} \int_0^{x_1} y^{1-\alpha/2} dy \\ &= C \frac{1}{3!(2-\alpha/2)} x_1^{2-\alpha/2} \end{aligned}$$

209 so

$$\frac{2}{h_1 + h_2} \frac{1}{h_1} \int_0^{x_1} f''(y) \frac{y^3}{3!} dy = \frac{C 2^{1-r}}{3!(2-\alpha/2)} x_1^{-\alpha/2} = \frac{C 2^{1-r}}{3!(2-\alpha/2)} T^{2/r} h^2 x_1^{-\alpha/2-2/r}$$

211 And for $i = N$, we have

$$\begin{aligned} \frac{2}{h_N + h_{N+1}} (h_N^3 f''(\eta_1) + h_{N+1}^3 f''(\eta_2)) \\ &= h_N^2 (f''(\eta_1) + f''(\eta_2)) \\ &\leq r^2 T^{2/r} h^2 x_N^{2-2/r} 2C x_{N-1}^{-2-\alpha/2} \\ &\leq 2r^2 T^{2/r} C 2^{-r(-2-\alpha/2)} h^2 x_N^{-\alpha/2-2/r} \end{aligned}$$

213 Finally, $N + 1 \leq i \leq 2N - 1$ is symmetric to the first half of the proof, so we can
214 conclude that □

$$\frac{2}{h_i + h_{i+1}} (h_i^3 f''(\eta_1) + h_{i+1}^3 f''(\eta_2)) \leq C h^2 \begin{cases} x_i^{-\alpha/2-2/r}, & 1 \leq i \leq N \\ (2T - x_i)^{-\alpha/2-2/r}, & N \leq i \leq 2N - 1 \end{cases}$$

216 LEMMA C.1. *There is a constant C for $2 \leq j \leq N$, if $y \in [x_{j-1}, x_j]$,*

$$(C.1) \quad |u(y) - u_h(y)| \leq C h^2 y^{\alpha/2-2/r}$$

218 *Proof.* For $2 \leq j \leq N$, we have

$$x_j \leq 2^r y, \quad x_{j-1} \geq 2^{-r} y$$

220 And by Lemma A.2 and Lemma B.1, we have

$$\begin{aligned} u(y) - u_h(y) &= -\frac{\theta(1-\theta)}{2} h_j^2 u''(\xi), \quad \xi \in [x_{j-1}, x_j] \\ &\leq \frac{C}{4} h^2 x_j^{2-2/r} x_{j-1}^{\alpha/2-2} \\ &\leq \frac{C}{4} h^2 2^{2r-2} y^{2-2/r} 2^{-r(\alpha/2-2)} y^{\alpha/2-2} \\ &= C 2^{-r\alpha/2-r} h^2 y^{\alpha/2-2/r} \end{aligned}$$

222 symmetricly, for $N < j \leq 2N - 1$, we have

$$(C.2) \quad |u(y) - u_h(y)| \leq C h^2 (2T - y)^{\alpha/2-2/r} \quad \square$$

224 LEMMA C.2. *There is a constant C such that for all $1 \leq i < N/2$,
225 $\max\{2i + 1, i + 3\} \leq j \leq N$ and $y \in [x_{j-1}, x_j]$, we have*

$$(C.3) \quad D_h^2 \left(\frac{|y - \cdot|^{1-\alpha}}{\Gamma(2-\alpha)} \right) (x_i) \leq C y^{-1-\alpha}$$

Proof. Since $y \geq x_{j-1} > x_{i+1}$, by Lemma A.1, if $j - 1 > i + 1$

$$\begin{aligned} D_h^2\left(\frac{|y - \cdot|^{1-\alpha}}{\Gamma(2-\alpha)}\right)(x_i) &= \frac{|y - \xi|^{-1-\alpha}}{\Gamma(-\alpha)}, \quad \xi \in [x_{i-1}, x_{i+1}] \\ &\leq \frac{(y - x_{i+1})^{-1-\alpha}}{\Gamma(-\alpha)} \\ &\leq \left(1 - \left(\frac{2}{3}\right)^r\right)^{-1-\alpha} \frac{y^{-1-\alpha}}{\Gamma(-\alpha)} \end{aligned}$$

But if $i = 1, j = 3, y \in [x_2, x_3]$

□

$$D_h^2\left(\frac{|y - \cdot|^{1-\alpha}}{\Gamma(2-\alpha)}\right)(x_1) = \frac{2}{x_2}()$$

Acknowledgments. We would like to acknowledge the assistance of volunteers in putting together this example manuscript and supplement.

REFERENCES

- [1] X. ROS-OTON AND J. SERRA, *The dirichlet problem for the fractional laplacian: Regularity up to the boundary*, Journal de Mathématiques Pures et Appliquées, 101 (2014), pp. 275–302, <https://doi.org/https://doi.org/10.1016/j.matpur.2013.06.003>, <https://www.sciencedirect.com/science/article/pii/S0021782413000895>.